

MEDICAL POLICY - 8.01.540

Cranial Electrotherapy Stimulation and Auricular Electrostimulation

BCBSA Ref. Policy: 8.01.58

Effective Date: May 1, 2024 RELATED MEDICAL POLICIES:

Last Revised: April 9, 2024 1.01.507 Electrical Stimulation Devices

Replaces: 8.01.58 2.01.526 Transcranial Magnetic Stimulation as a Treatment of Depression and

Other Psychiatric/Neurologic Disorders

7.01.588 Percutaneous Electrical Nerve Stimulation and Percutaneous

Neuromodulation Therapy

Select a hyperlink below to be directed to that section.

POLICY CRITERIA | CODING | RELATED INFORMATION EVIDENCE REVIEW | REFERENCES | HISTORY

Clicking this icon returns you to the hyperlinks menu above.

Introduction

Cranial electrotherapy stimulation (CES) provides weak levels of electrical current to the brain. Electrodes are placed on the skull, earlobes, eyelids, or forehead. A small transmitter then sends weak electrical pulses into the brain. It's believed the current affects particular areas of the brain that play important roles in the body's hormones and emotions. Another use of CES calls for treating pain by placing the electrodes near the site of pain. Auricular stimulation sends electrical pulses to the acupuncture points of the ear. Both of these systems have been used for depression, anxiety, insomnia (sleeplessness), and weight loss. Because there is not enough medical evidence showing that these technologies improve health, both are considered unproven. A small device worn behind the ear that stimulates specific nerves to try to relieve symptoms of opioid withdrawal is unproven. More research is needed to determine if this device is effective.

Note: The Introduction section is for your general knowledge and is not to be taken as policy coverage criteria. The rest of the policy uses specific words and concepts familiar to medical professionals. It is intended for providers. A provider can be a person, such as a doctor, nurse, psychologist, or dentist. A provider also can

be a place where medical care is given, like a hospital, clinic, or lab. This policy informs them about when a service may be covered.

Policy Coverage Criteria

Procedure	Investigational
Cranial electrotherapy	Cranial electrotherapy stimulation (also known as cranial
stimulation	electrostimulation therapy or CES) is investigational in all
	situations.
Auricular electrostimulation	Electrical stimulation of auricular acupuncture points is
	investigational in all situations.
	Electrical stimulation of the ear (also known as auricular
	neurostimulation) for opioid withdrawal is investigational.

Coding

Code	Description
СРТ	
0783T	Transcutaneous auricular neurostimulation, set-up, calibration, and patient education on use of equipment
HCPCS	
A4596	Cranial electrotherapy stimulation (CES) system supplies and accessories, per month
E0732	Cranial electrotherapy stimulation (CES) system, any type (new code effective 1/1/2024)
K1002	Cranial electrotherapy stimulation (CES) system, any type (code termed 1/1/2024)
S8930	Electrical stimulation of auricular acupuncture points; each 15 minutes of personal one-on-one contact with the patient

Note: CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). HCPCS codes, descriptions and materials are copyrighted by Centers for Medicare Services (CMS).

Related Information

N/A

Evidence Review

Description

Cranial electrotherapy stimulation (CES), also known as cranial electrical stimulation, transcranial electrical stimulation, or electrical stimulation therapy, delivers weak pulses of electrical current to the earlobes, mastoid processes, or scalp with devices such as the Alpha-Stim. Auricular electrostimulation involves stimulation of acupuncture points on the ear. Devices, including the P-Stim and E-pulse, provide ambulatory auricular electrical stimulation over a period of several days. CES is being evaluated for a variety of conditions, including pain, insomnia, depression, anxiety, and functional constipation. Auricular electrical stimulation is being evaluated for pain, weight loss, and opioid withdrawal.

Background

CES, also known as cranial electrical stimulation, transcranial electrical stimulation, or electrical stimulation therapy, delivers weak pulses of electrical current to the earlobes, mastoid processes, or scalp with devices such as the Alpha-Stim. Auricular electrostimulation involves stimulation of acupuncture points on the ear. Devices, including the P-Stim and E-pulse, provide ambulatory auricular electrical stimulation over a period of several days. CES and auricular electrostimulation are being evaluated for a variety of conditions, including pain, insomnia, depression, anxiety, weight loss, and opioid withdrawal.

Interest in CES began in the early 1900s on the theory that weak pulses of electrical current have a calming effect on the central nervous system. The technique was further developed in the USS.R. and Eastern Europe in the 1950s as a treatment for anxiety and depression. The use of CES later spread to Western Europe and the United States as a treatment for various psychological and physiological conditions. Presently, the mechanism of action is thought to be the modulation of activity in brain networks by direct action in the hypothalamus, limbic system,



and/or the reticular activating system. One device used in the Unites States is the Alpha-Stim CES, which provides pulsed, low-intensity current via clip electrodes that attach to the earlobes. Other devices place the electrodes on the eyelids, frontal scalp, mastoid processes, or behind the ears. Treatments may be administered once or twice daily for several days to several weeks.

Other devices provide electrical stimulation to auricular acupuncture sites over several days. One device, the P-Stim, is a single-use miniature electrical stimulator for auricular acupuncture points that is worn behind the ear with a self-adhesive electrode patch. A selection stylus that measures electrical resistance is used to identify three auricular acupuncture points. The P-Stim device connects to three inserted acupuncture needles with caps and wires. The device is preprogrammed to be on for 180 minutes, then off for 180 minutes. The maximum battery life of this single-use device is 96 hours.

Summary of Evidence

Cranial Electrotherapy Stimulation

For individuals with acute or chronic pain who receive CES the evidence includes a number of small sham-controlled randomized trials and pooled analyses. The relevant outcomes are symptoms, morbid events, functional outcomes, and treatment-related morbidity. Systematic reviews of randomized trials evaluated CES for headache and chronic pain. Pooled analyses found marginal benefits for a headache with CES and no benefits for chronic pain with CES. A subsequent sham-controlled trial of remotely supervised CES via secure videoconferencing found a significant benefit with CES for pain reduction, but it had important relevance and conduct and design limitations. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with psychiatric, behavioral, or neurologic conditions (e.g., depression and anxiety, Parkinson disease, addiction) who receive CES, the evidence includes a number of small sham-controlled randomized trials and systematic reviews. The relevant outcomes are symptoms, morbid events, functional outcomes, and treatment-related morbidity. Four randomized controlled trials (RCTs) evaluated CES for depression and anxiety. One RCT each found a significant benefit with CES for anxiety or depression, but both had important relevance limitations. Comparisons between these trials cannot be made due to the heterogeneity in study populations and treatment protocols. Studies evaluating CES for Parkinson disease, smoking cessation and tic disorders do not support the use of CES for these conditions. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.



For individuals who have functional constipation who receive CES, the evidence includes an RCT. The relevant outcomes are symptoms, morbid events, functional outcomes, and treatment-related morbidity. The single RCT reported positive results for the treatment of constipation with CES. However, the trial was unblinded and most outcomes were self-reported. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Auricular Electrostimulation

For individuals with acute or chronic pain (e.g., acute pain from surgical procedures, chronic back pain, chronic pain from osteoarthritis or rheumatoid arthritis) who receive auricular electrostimulation, the evidence includes a limited number of trials. The relevant outcomes are symptoms, morbid events, functional outcomes, and treatment-related morbidity. Studies evaluating the effect of electrostimulation technology on acute pain are inconsistent, and the small amount of evidence on chronic pain has methodologic limitations. For example, a comparison of auricular electrostimulation with manual acupuncture for chronic low back pain did not include a sham-control group, and, in a study of rheumatoid arthritis, auricular electrostimulation was compared with autogenic training and resulted in a small improvement in visual analog scale pain scores of unclear clinical significance. Overall, the few published studies have small sample sizes and methodologic limitations. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have obesity who receive auricular electrostimulation, the evidence includes small RCTs and systematic reviews. The relevant outcomes are symptoms, morbid events, functional outcomes, and treatment-related morbidity. The RCTs reported inconsistent results and used different treatment protocols. The systematic reviews are limited by high heterogeneity with respect to the interventions used, participants included, treatment period, and outcome measures. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have opioid withdrawal symptoms who receive auricular electrostimulation, the evidence includes two observational studies. The relevant outcomes are symptoms, morbid events, functional outcomes, and treatment-related morbidity. Both studies report positive outcomes for the use of CES to treat opioid withdrawal symptoms. The studies used different treatment protocols and no comparators, limiting conclusions drawn from the results. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Ongoing and Unpublished Clinical Trials

Table 1 provides a summary of ongoing and unpublished trials that may influence this review.

Table 1. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
Ongoing			
NCT03825471	Effects of Cranial Electrotherapy Stimulation on Anesthetics Consumption, Perioperative Cytokines Response, and Postoperative Pain in Patients Undergoing Colonic Surgery	80	Dec 2020
NCT03896438	Increased Thalamocortical Connectivity in Tdcs-potentiated Generalization of Cognitive Training	85	April 2024
Unpublished			
NCT05384041	Cranial Electrotherapy Stimulation for the Treatment of Major Depressive Disorder in Adults	255	Oct 2022
NCT03815253	Electro-acupuncture for Central Obesity	168	Feb 2021
NCT03060122	The Clinical Feasibility of Combining Cranial Electrotherapy Stimulation (CES Alpha-Stim) and Non-invasive Interactive Neurostimulation (InterX) for Optimized Rehabilitation Following Extremity Immobilization	35	Aug 2019

NCT: national clinical trial

Clinical Input Received from Physician Specialty Societies and Academic Medical Centers

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the policy conclusions.

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

^a Denotes industry sponsorship

2011 Input

In response to requests, input on auricular electrostimulation was received from three physician specialty societies and five academic medical centers while this policy was under review in 2011. There was a consensus that auricular electrostimulation is investigational.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion if they were issued by, or jointly by, a US professional society, an international society with US representation, or the National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

No guidelines or statements were identified.

Medicare National Coverage

There is no national coverage determination.

Regulatory Status

A number of devices for CES have been cleared for marketing by the US Food and Drug Administration's (FDA) 510(k) process. In 1992, the Alpha-Stim CES device (Electromedical Products International) received marketing clearance for the treatment of anxiety, insomnia, and depression. Devices cleared since 2000 are summarized in **Table 2**.

FDA product code: QJQ.

Table 2: Cranial Electrotherapy Stimulation (CES) Devices Cleared by the US Food and Drug Administration

Device Name	Manufacturer	Date Cleared	510(k) No.	Indications
Modius Sleep	Neurovalens Limited	10/27/2023	K230826	Insomnia
Cervella	Innovative Neurological Devices	03/07/2019	K182311	Insomnia, depression, anxiety
Cranial Electrical Nerve Stimulator	Johari Digital Healthcare	05/29/2009	K090052	Insomnia, depression, anxiety
Elexoma Medic	Redplane AG	05/21/2008	K070412	Insomnia, depression, anxiety
CES Ultra	Neuro-Fitness	04/05/2007	K062284	Insomnia, depression, anxiety
Net-2000 Microcurrent Stimulator	Auri-Stim Medical	10/13/2006	K060158	Insomnia, depression, anxiety
Transcranial Electrotherapy Stimulator-A, Model TESA-1	Kalaco Scientific	07/21/2003	K024377	Insomnia, depression, anxiety

FDA: Food and Drug Administration

Several devices for electroacupuncture designed to stimulate auricular acupuncture points have been cleared for marketing by FDA through the 510(k) process. Devices cleared since 2000 are summarized in Table 3.

FDA product code: BWK, PZR.

Table 3: Auricular Electrostimulation Devices Cleared by the US Food and Drug Administration

Device Name	Manufacturer	Date Cleared	510(k) No.	Indications
Needle Stimulator	Wuxi Jiajian Medical Instrument	08/27/2021	K202861	Practice of acupuncture by qualified practitioners of

Device Name	Manufacturer	Date	510(k)	Indications
		Cleared	No.	
				acupuncture as determined by the states
AXUS ES-5 Electro- Acupuncture Device	Lhasa OMS, INC.	02/03/2021	K200636	Practice of acupuncture by qualified practitioners of acupuncture as determined by the states
Drug Relief V1	DyAnsys Inc	11/05/2021	K211971	Reduce symptoms of opioid withdrawal
Sparrow Therapy System	Spark Biomedical, Inc.	01/02/2021	K201873	Reduce symptoms of opioid withdrawal
Drug Relief	DyAnsys	05/02/2018	K173861	Reduce symptoms of opioid withdrawal
Ansistem-Pp	DyAnsys Inc	03/09/2017	K170391	Practice of acupuncture by qualified practitioners of acupuncture as determined by the states
NSS-2 Bridge	Innovative Health Solutions	2017	N/Aª	Substance use disorders
Stivax System	Biegler Gmbh	05/26/2016	K152571	Practice of acupuncture by qualified practitioners of acupuncture as determined by the states
ANSiStim	DyAnsys Inc	05/15/2015	K141168	Practice of acupuncture by qualified practitioners of acupuncture as determined by the states
Pantheon Electrostimulator	Pantheon Research	11/07/2014	K133980	Practice of acupuncture by qualified practitioners as determined by the states
Electro Auricular Device	Navigant Consulting, Inc.	10/02/2014	K140530	Practice of acupuncture by qualified practitioners of acupuncture as determined by the states
P-Stim	Biegler GMBH	06/27/2014	K140788	Practice of acupuncture by qualified practitioners of acupuncture as determined by the states



Device Name	Manufacturer	Date	510(k)	Indications
		Cleared	No.	
Jiajian Cmn Stimulator	Wuxi Jiajian Medical Instrument Co., Ltd.	08/16/2013	K130768	Practice of acupuncture by qualified practitioners of acupuncture as determined by the states
JiaJian Electro- Acupuncture Stimulators	Wuxi Jiajian Medical Instrument Co., Ltd.	04/11/2013	K122812	Practice of acupuncture by qualified practitioners of acupuncture as determined by the states
Multi-Purpose Health Device	UPC Medical Supplies, Inc. DBA United Pacific Co.	08/05/2010	K093322	Unknown - Summary not provided
Electro-Acupuncture: Aculife/Model ADOC-01	Inno-Health Technology, Inc.	04/02/2010	K091933	Practice of acupuncture by qualified practitioners of acupuncture as determined by the states
e-Pulse	Medevice Corporation	12/07/2009	K091875	Practice of acupuncture by qualified practitioners of acupuncture as determined by the states
Model ES-130	Ito Co., Ltd.	11/24/2008	K081943	Practice of acupuncture by qualified practitioners of acupuncture as determined by the states
P-Stim	NeuroScience Therapy Corp	03/30/2006	K050123	Practice of acupuncture by qualified practitioners of acupuncture as determined by the states
Aculife	Inno-Health Technology, Inc.	03/28/2006	К051197	Practice of acupuncture by qualified practitioners of acupuncture as determined by the states
AcuStim	S.H.P. Intl. Pty., Ltd.	06/12/2002	K014273	As an electroacupuncture device

^a "FDA cleared the NSS-2 Bridge Device for Substance Use Disorders through the de novo premarket review pathway, a regulatory pathway for some low- to moderate-risk devices that are novel and for which there is no legally marketed predicate device to which the device can claim substantial equivalence"¹ N/A: Not applicable



References

- US Food & Drug Administration. FDA News Release: FDA grants marketing authorization of the first device for use in helping to reduce the symptoms of opioid withdrawal. https://www.fda.gov/news-events/press-announcements/fda-grants-marketing-authorization-first-device-use-helping-reduce-symptoms-opioid-withdrawal. Accessed March 25, 2024.
- 2. Klawansky S, Yeung A, Berkey C, et al. Meta-analysis of randomized controlled trials of cranial electrostimulation. Efficacy in treating selected psychological and physiological conditions. J Nerv Ment Dis. Jul 1995; 183(7): 478-84. PMID 7623022
- 3. Bronfort G, Nilsson N, Haas M, et al. Non-invasive physical treatments for chronic/recurrent headache. Cochrane Database Syst Rev. 2004: (3): CD001878. PMID 15266458
- 4. Brønfort G, Haas M, Evans RL, et al. WITHDRAWN: Non-invasive physical treatments for chronic/recurrent headache. Cochrane Database Syst Rev. Aug 26 2014; 2014(8): CD001878. PMID 25157618
- 5. O'Connell NE, Wand BM, Marston L, et al. Non-invasive brain stimulation techniques for chronic pain. Cochrane Database Syst Rev. Apr 11 2014; (4): CD008208. PMID 24729198
- 6. O'Connell NE, Marston L, Spencer S, et al. Non-invasive brain stimulation techniques for chronic pain. Cochrane Database Syst Rev. Mar 16 2018; 3(3): CD008208. PMID 29547226
- 7. Ahn H, Galle K, Mathis KB, et al. Feasibility and efficacy of remotely supervised cranial electrical stimulation for pain in older adults with knee osteoarthritis: A randomized controlled pilot study. J Clin Neurosci. Jul 2020; 77: 128-133. PMID 32402609
- 8. Price L, Briley J, Haltiwanger S, et al. A meta-analysis of cranial electrotherapy stimulation in the treatment of depression. J Psychiatr Res. Mar 2021; 135: 119-134. PMID 33477056
- Ching PY, Hsu TW, Chen GW, et al. Efficacy and Tolerability of Cranial Electrotherapy Stimulation in the Treatment of Anxiety: A Systemic Review and Meta-Analysis. Front Psychiatry. 2022; 13: 899040. PMID 35757229
- 10. Patel S, Boutry C, Patel P, et al. A randomised controlled trial investigating the clinical and cost-effectiveness of Alpha-Stim AID cranial electrotherapy stimulation (CES) in patients seeking treatment for moderate severity depression in primary care (Alpha-Stim-D Trial). Trials. Apr 04 2022; 23(1): 250. PMID 35379314
- 11. Morriss R, Patel S, Boutry C, et al. Clinical effectiveness of active Alpha-Stim AlD versus sham Alpha-Stim AlD in major depression in primary care in England (Alpha-Stim-D): a multicentre, parallel group, double-blind, randomised controlled trial. Lancet Psychiatry. Mar 2023; 10(3): 172-183. PMID 36724796
- 12. Kim J, Kim H, Kim DH, et al. Effects of cranial electrotherapy stimulation with novel in-ear electrodes on anxiety and restingstate brain activity: A randomized double-blind placebo-controlled trial. J Affect Disord. Dec 01 2021; 295: 856-864. PMID 34706456
- 13. Barclay TH, Barclay RD. A clinical trial of cranial electrotherapy stimulation for anxiety and comorbid depression. J Affect Disord. Aug 2014; 164: 171-7. PMID 24856571
- 14. Shekelle PG, Cook IA, Miake-Lye IM, et al. Benefits and Harms of Cranial Electrical Stimulation for Chronic Painful Conditions, Depression, Anxiety, and Insomnia: A Systematic Review. Ann Intern Med. Mar 20 2018; 168(6): 414-421. PMID 29435567
- 15. Mischoulon D, De Jong MF, Vitolo OV, et al. Efficacy and safety of a form of cranial electrical stimulation (CES) as an add-on intervention for treatment-resistant major depressive disorder: A three week double blind pilot study. J Psychiatr Res. Nov 2015; 70: 98-105. PMID 26424428
- 16. Lyon D, Kelly D, Walter J, et al. Randomized sham controlled trial of cranial microcurrent stimulation for symptoms of depression, anxiety, pain, fatigue and sleep disturbances in women receiving chemotherapy for early-stage breast cancer. Springerplus. 2015; 4: 369. PMID 26435889
- 17. Shill HA, Obradov S, Katsnelson Y, et al. A randomized, double-blind trial of transcranial electrostimulation in early Parkinson's disease. Mov Disord. Jul 2011; 26(8): 1477-80. PMID 21538515



- 18. Pickworth WB, Fant RV, Butschky MF, et al. Evaluation of cranial electrostimulation therapy on short-term smoking cessation. Biol Psychiatry. Jul 15 1997; 42(2): 116-21. PMID 9209728
- 19. Wu WJ, Wang Y, Cai M, et al. A double-blind, randomized, sham-controlled study of cranial electrotherapy stimulation as an add-on treatment for tic disorders in children and adolescents. Asian J Psychiatr. Jun 2020; 51: 101992. PMID 32145674
- Gong BY, Ma HM, Zang XY, et al. Efficacy of Cranial Electrotherapy Stimulation Combined with Biofeedback Therapy in Patients with Functional Constipation. J Neurogastroenterol Motil. Jul 30 2016; 22(3): 497-508. PMID 26932836
- 21. Sator-Katzenschlager SM, Michalek-Sauberer A. P-Stim auricular electroacupuncture stimulation device for pain relief. Expert Rev Med Devices. Jan 2007; 4(1): 23-32. PMID 17187468
- 22. Holzer A, Leitgeb U, Spacek A, et al. Auricular acupuncture for postoperative pain after gynecological surgery: a randomized controlled trail. Minerva Anestesiol. Mar 2011; 77(3): 298-304. PMID 21441884
- 23. Sator-Katzenschlager SM, Scharbert G, Kozek-Langenecker SA, et al. The short- and long-term benefit in chronic low back pain through adjuvant electrical versus manual auricular acupuncture. Anesth Analg. May 2004; 98(5): 1359-64, table of contents. PMID 15105215
- 24. Sator-Katzenschlager SM, Szeles JC, Scharbert G, et al. Electrical stimulation of auricular acupuncture points is more effective than conventional manual auricular acupuncture in chronic cervical pain: a pilot study. Anesth Analg. Nov 2003; 97(5): 1469-1473. PMID 14570667
- 25. Bernateck M, Becker M, Schwake C, et al. Adjuvant auricular electroacupuncture and autogenic training in rheumatoid arthritis: a randomized controlled trial. Auricular acupuncture and autogenic training in rheumatoid arthritis. Forsch Komplementmed. Aug 2008; 15(4): 187-93. PMID 18787327
- 26. Kim SY, Shin IS, Park YJ. Effect of acupuncture and intervention types on weight loss: a systematic review and meta-analysis. Obes Rev. Nov 2018; 19(11): 1585-1596. PMID 30180304
- 27. Yeh TL, Chen HH, Pai TP, et. al. The Effect of Auricular Acupoint Stimulation in Overweight and Obese Adults: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. Evidence-Based Complementary and Alternative Medicine. 2017; vol. 2017, Article ID 3080547, 16 pages, 2017. https://doi.org/10.1155/2017/3080547.
- 28. Schukro RP, Heiserer C, Michalek-Sauberer A, et al. The effects of auricular electroacupuncture on obesity in female patients--a prospective randomized placebo-controlled pilot study. Complement Ther Med. Feb 2014; 22(1): 21-5. PMID 24559812
- 29. Yeh ML, Chu NF, Hsu MY, et al. Acupoint Stimulation on Weight Reduction for Obesity: A Randomized Sham-Controlled Study. West J Nurs Res. Dec 2015; 37(12): 1517-30. PMID 25183702
- 30. Kroening RJ, Oleson TD. Rapid narcotic detoxification in chronic pain patients treated with auricular electroacupuncture and naloxone. Int J Addict. Sep 1985; 20(9): 1347-60. PMID 2867052
- 31. Miranda A, Taca A. Neuromodulation with percutaneous electrical nerve field stimulation is associated with reduction in signs and symptoms of opioid withdrawal: a multisite, retrospective assessment. Am J Drug Alcohol Abuse. 2018; 44(1): 56-63. PMID 28301217US

History

Date	Comments
11/08/11	New policy; add to Therapy section. Policy created with literature search through April 2011; clinical input reviewed; considered investigational.



Date	Comments	
11/13/12	Replace policy. Policy updated with literature review through June 2012, references 1-7 added; cranial electrotherapy stimulation (CES) added as investigational. "Cranial Electrotherapy Stimulation (CES)" added to policy title.	
01/22/13	Update Related Policies. 2.01.50 has been replaced with 2.01.526.	
02/15/13	Update Related Policies. Change title to policy 2.01.526.	
10/14/13	Replace policy. Policy updated with literature review through July 10, 2013; policy statement unchanged.	
11/20/14	Annual Review. Policy updated with literature review through July 16, 2014. References 4-5, 7, and 14 added; others renumbered. Policy statement unchanged.	
10/13/15	Annual Review. Policy updated with literature review through July 6, 2015; no references added. Policy statements unchanged. Related policies updated; 2.01.526 removed.	
05/01/16	Annual Review, approved April 12, 2016. Policy updated with literature review through December 10, 2015; references 6-7 added. Policy statement unchanged.	
06/01/17	Annual Review, approved May 2, 2017. Policy updated with literature review through December 22, 2016; references 8 and 12 added. Policy statements unchanged.	
10/17/17	Coding update; added CPT code 20974.	
08/01/18	Annual Review, approved July 10, 2018. Policy updated with literature review through April 2018; 17-19 references added. Added electrical stimulation of the ear (also known as auricular neurostimulation) for opioid withdrawal is investigational. Removed CPT code 20974.	
05/01/19	Annual Review, approved April 18, 2019. Policy updated with literature review through January 2019; reference 16 added. Policy statements unchanged.	
01/01/20	Coding update, added HCPCS code K1002 (new code effective 1/1/20).	
05/01/20	Annual Review, approved April 7, 2020. Policy updated with literature review through December 2019; references added. Policy statements unchanged. Removed CPT codes 97813 and 97814.	
05/01/21	Annual Review, approved April 1, 2021. Policy updated with literature review through December 14, 2020; references added. Policy statements unchanged.	
05/01/22	Annual Review, approved April 11, 2022. Policy updated with literature review through December 10, 2021; references added. Policy statements unchanged.	
10/01/22	Coding update. Updated description of HCPCS code K1002.	
11/01/22	Coding update. Added HCPCS code A4596.	
01/01/23	Coding update. Added new CPT code 0783T. Added HCPC code L8692.	
05/01/23	Annual Review, approved April 10, 2023. Policy updated with literature review through December 27, 2022; references added. Policy statements unchanged.	



Date	Comments
10/04/23	Updated related policy. Policy 7.01.29 Percutaneous Electrical Nerve Stimulation and Percutaneous Neuromodulation Therapy was renumbered to 7.01.588 Percutaneous Electrical Nerve Stimulation and Percutaneous Neuromodulation Therapy.
01/01/24	Coding update. Added new HCPCS code E0732 and added term date to HCPCS code K1002.
05/01/24	Policy renumbered from 8.01.58 to 8.01.540 Cranial Electrotherapy Stimulation and Auricular Electrostimulation, approved April 9, 2024. Policy updated with literature review through January 3, 2024; reference added. Policy statements unchanged. Removed HCPCS L8692.

Disclaimer: This medical policy is a guide in evaluating the medical necessity of a particular service or treatment. The Company adopts policies after careful review of published peer-reviewed scientific literature, national guidelines and local standards of practice. Since medical technology is constantly changing, the Company reserves the right to review and update policies as appropriate. Member contracts differ in their benefits. Always consult the member benefit booklet or contact a member service representative to determine coverage for a specific medical service or supply. CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). ©2024 Premera All Rights Reserved.

Scope: Medical policies are systematically developed guidelines that serve as a resource for Company staff when determining coverage for specific medical procedures, drugs or devices. Coverage for medical services is subject to the limits and conditions of the member benefit plan. Members and their providers should consult the member benefit booklet or contact a customer service representative to determine whether there are any benefit limitations applicable to this service or supply. This medical policy does not apply to Medicare Advantage.



PREMERA . HMO

Discrimination is Against the Law

Premera Blue Cross HMO (Premera HMO) complies with applicable Federal and Washington state civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, sex, gender identity, or sexual orientation. Premera HMO does not exclude people or treat them differently because of race, color, national origin, age, disability, sex, gender identity, or sexual orientation. Premera HMO provides free aids and services to people with disabilities to communicate effectively with us, such as qualified sign language interpreters and written information in other formats (large print, audio, accessible electronic formats, other formats). Premera HMO provides free language services to people whose primary language is not English, such as qualified interpreters and information written in other languages. If you need these services, contact the Civil Rights Coordinator. If you believe that Premera HMO has failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, sex, gender identity, or sexual orientation, you can file a grievance with: Civil Rights Coordinator — Complaints and Appeals, PO Box 91102, Seattle, WA 98111, Toll free: 855-332-4535, Fax: 425-918-5592, TTY: 711, Email AppealsDepartmentInquiries@Premera.com. You can file a grievance in person or by mail, fax, or email. If you need help filing a grievance, the Civil Rights Coordinator is available to help you. You can also file a civil rights complaint with the U.S. Department of Health and Human Services, Office for Civil Rights, electronically through the Office for Civil Rights Complaint Portal, available at https://ocrportal.hhs.gov/ocr/portal/lobby.jsf, or by mail or phone at: U.S. Department of Health and Human Services, 200 Independence Ave SW, Room 509F, HHH Building, Washington, D.C. 20201, 1-800-368-1019, 800-537-7697 (TDD). Complaint forms are available at http://www.hhs.gov/ocr/office/file/index.html. You can also file a civil rights complaint with the Washington State Office of the Insurance Commissioner, electronically through the Office of the Insurance Commissioner Complaint Portal available at https://www.insurance.wa.gov/file-complaint-or-check-your-complaint-status, or by phone at 800-562-6900, 360-586-0241 (TDD). Complaint forms are available at https://fortress.wa.gov/oic/onlineservices/cc/pub/complaintinformation.aspx.

Language Assistance

ATENCIÓN: si habla español, tiene a su disposición servicios gratuitos de asistencia lingüística. Llame al 844-722-4661 (TTY: 711). 注意:如果您使用繁體中文,您可以免費獲得語言援助服務。請致電 844-722-4661 (TTY: 711)。 CHÚ Ý: Nếu bạn nói Tiếng Việt, có các dịch vụ hỗ trợ ngôn ngữ miễn phí dành cho bạn. Gọi số 844-722-4661 (TTY: 711). 조의: 한국어를 사용하시는 경우, 언어 지원 서비스를 무료로 이용하실 수 있습니다. 844-722-4661 (TTY: 711) 번으로 전화해 주십시오. ВНИМАНИЕ: Если вы говорите на русском языке, то вам доступны бесплатные услуги перевода. Звоните 844-722-4661 (телетайп: 711). РАЦИАША: Кипд падзазавіта ка пд Тадаюд, тадагі капд дитаті пд тра serbisyo ng tulong sa wika nang walang bayad. Титаwад sa 844-722-4661 (ТТҮ: 711). УВАГА! Якщо ви розмовляєте українською мовою, ви можете звернутися до безкоштовної служби мовної підтримки. Телефонуйте за номером 844-722-4661 (телетайп: 711).

<u>المحوظة</u>؛ إذا كنت تتحدث اذكر اللغة، فإن خدمات المساعدة اللغوية تتوافر لك بالمجان. اتصل برقم 844-722-4661 (رقم هاتف الصم والبكم: 711). <u>ਧਿਆਨ ਦਿਓ</u>: ਜੇ ਤੁਸੀਂ ਪੰਜਾਬੀ ਬੋਲਦੇ ਹੋ, ਤਾਂ ਭਾਸ਼ਾ ਵਿੱਚ ਸਹਾਇਤਾ ਸੇਵਾ ਤੁਹਾਡੇ ਲਈ ਮੁਫਤ ਉਪਲਬਧ ਹੈ। 844-722-4661 (TTY: 711) 'ਤੇ ਕਾਲ ਕਹੋ। <u>ACHTUNG</u>: Wenn Sie Deutsch sprechen, stehen Ihnen kostenlos sprachliche Hilfsdienstleistungen zur Verfügung. Rufnummer: 844-722-4661 (TTY: 711). <u>ਪਿਨਕ੍ਰਾ</u>ਹ: ຖ້າວ່າ ທ່ານເວົ້າພາສາ ລາວ, ການບໍລິການຊ່ວຍເຫຼືອດ້ານພາສາ, ໂດຍບໍ່ເສັງຄ່າ, ແມ່ນມີພ້ອມໃຫ້ທ່ານ. ໂທຣ 844-722-4661 (TTY: 711). <u>ATANSYON</u>: Si w pale Kreyòl Ayisyen, gen sèvis èd pou lang ki disponib gratis pou ou. Rele 844-722-4661 (TTY: 711).

<u>ATTENTION</u>: Si vous parlez français, des services d'aide linguistique vous sont proposés gratuitement. Appelez le 844-722-4661 (ATS : 711). <u>UWAGA</u>: Jeżeli mówisz po polsku, możesz skorzystać z bezpłatnej pomocy językowej. Zadzwoń pod numer 844-722-4661 (TTY: 711). <u>ATENÇÃO</u>: Se fala português, encontram-se disponíveis serviços linguísticos, grátis. Ligue para 844-722-4661 (TTY: 711).

ATTENZIONE: In caso la lingua parlata sia l'italiano, sono disponibili servizi di assistenza linguistica gratuiti. Chiamare il numero 844-722-4661 (TTY: 711). منايد، توجه: اگر به زبان فارسی گفتگو می کنید، تسهیلات زبانی بصورت رایگان برای شما فراهم می باشد. با (TTY: 711) 844-722-4661 تماس بگیرید.