

Five things to know about COVID-19, flu, and RSV symptoms

... from Premera Blue Cross Blue Shield



With COVID-19, the flu, and some allergies having similar symptoms, it can be difficult to know if you should get a test, quarantine, or just take it easy. Here are five things to know that will help you identify the cause of your symptoms.

1 What are common flu symptoms?

According to the Centers for Disease Control and Prevention (CDC), flu symptoms usually appear suddenly after exposure. Here are some of the possible signs you may have the flu:

- Fever or chills
- Cough
- Sore throat
- Runny or stuffy nose
- Muscle or body aches
- Headaches
- Fatigue

Flu can cause serious illness and sometimes death. Because this virus changes each year, it's recommended that everyone who is eligible get a flu shot before the start of the flu season, which typically spans from November through February. The flu is mostly contagious during the first three to four days of the illness, but sometimes it can be contagious even longer. Visit the CDC website to [find out more about the flu](#).

2

How are flu symptoms different than COVID symptoms?

What used to be a clear sign of COVID is not so clear-cut anymore. Dulled senses of taste and smell are no longer considered good indicators of a COVID infection. This makes it even harder to tell the difference between a cold, a flu, and a COVID infection. The new **COVID variants** like Eris (EG.5) and Omicron BA.2.86, tend to have the following **symptoms**:

- Fever or chills
- Cough
- Shortness of breath or difficulty breathing
- Fatigue
- Muscle or body aches
- Headache
- New loss of taste or smell
- Sore throat
- Congestion or runny nose
- Diarrhea
- Nausea or vomiting

These symptoms may be a sign of a COVID infection or another illness.

Those sick with COVID can spread the virus that causes the disease for two to three days before symptoms appear and you can still be infectious eight days after you no longer have symptoms. If symptoms do appear, it's best to keep clear of big gatherings and notify those you've been in contact with even if they aren't showing symptoms. Find out more **about COVID** here.

3

When should I take a COVID test?

If you have symptoms or have been exposed to COVID, it's recommended that you self-test with an at-home COVID test. You can also visit your primary care provider or your local pharmacy for a test. Many pharmacies offer "test to treat" meaning if you do test positive, they may be able to prescribe treatments for COVID.

You can get free at-home COVID tests from the government at covidtests.gov. If you still have unused at-home test kits, check the expiration dates as many of the expiration dates have been extended. Testing not only helps you, but it also keeps those around you safe. Don't wait to test if you think you have contracted COVID.

4

What should I do if I test positive for COVID or if I've been exposed to someone with COVID?

The CDC recommends that if you test positive for COVID, you should stay home and isolate yourself from others for at least five days from the date you tested positive. If you have no symptoms, you can end isolation after five days. If you do have symptoms, the CDC recommends that you wait until you are fever free for 24 hours, without the use of fever-reducing medicines, to end your isolation.

It is also recommended that when you do end isolation, you continue to wear a mask when around others for at least 10 days. You can discover more about exposure and recommendations **from the CDC here**.

5

What is RSV?

Respiratory Syncytial Virus (RSV) is a potentially dangerous, highly contagious virus that can cause respiratory disease. Anyone can get RSV, but providers know it can be more serious for babies as well as adults over 60, including those with underlying medical conditions.

RSV usually causes mild, cold-like symptoms that typically show up within four to six days after getting infected. Those symptoms include the following:

- Runny nose
- Decrease in appetite
- Coughing
- Sneezing
- Fever
- Wheezing

Anyone infected with RSV is usually contagious for three to eight days and may be contagious for one to two days before symptoms appear. There is a new vaccine available, and the CDC recommends that you check with your provider about whether the RSV vaccine is right for you.

Because symptoms are similar between all three viruses, it's important to get tested and get right treatments or medications as soon as possible. Check with your healthcare provider for more information about testing and treatment. The more education you have about these infections, the safer you, your family, and your community will be.

Check out our previous 'Five things to know' blogs to discover more [about getting vaccinated](#) and for more information about [how the latest vaccines can help](#) prevent you from becoming ill or lessen your symptoms

Notice of availability and nondiscrimination 800-508-4722 | TTY: 711

Call for free language assistance services and appropriate auxiliary aids and services.

Tumawag para sa mga libreng serbisyo ng tulong sa wika at angkop na mga karagdagang tulong at serbisyo.

Llame para obtener servicios gratuitos de asistencia lingüística, y ayudas y servicios auxiliares apropiados.

무료 언어 지원 서비스와 적절한 보조 도구 및 서비스를 신청하십시오.

Hu thov kev pab txhais lus pub dawb thiab lwm yam khoom pab dawb thiab kev pab cuam ua tsim nyog.

Звоните для получения бесплатных услуг по переводу и других вспомогательных средств и услуг.

呼吁提供免费的语言援助服务和适当的辅助设备及服务。

呼籲提供免費的語言援助服務和適當的輔助設備及服務。

Vala'au mo auaunaga tau fesoasoani mo gagana e leai ni totogi ma fesoasoani fa'aopo'opo talafeagai ma auaunaga.

ໂທເພື່ອຮັບການບໍລິການຊ່ວຍເຫຼືອດ້ານພາສາ ແລະ ການບໍລິການ ແລະ ການຊ່ວຍເຫຼືອພິເສດທີ່ເໝາະສົມແບບບໍ່ເສຍຄ່າ.

無料言語支援サービスと適切な補助器具及びサービスをお求めください。

Tumawag para kadagiti libre a serbisio iti tulong iti pagsasao ken dagiti nakanada nga aid ken serbisio iti komunikasion.

Gọi cho các dịch vụ hỗ trợ ngôn ngữ miễn phí và các hỗ trợ và dịch vụ phụ trợ thích hợp.

Звертайтеся за безкоштовною мовною підтримкою та відповідними додатковими послугами.

ติดต่อขอบริการช่วยเหลือด้านภาษาฟรีพร้อมความช่วยเหลือและบริการอื่นๆ เพิ่มเติม

Fordern Sie kostenlose Sprachunterstützungsdienste und geeignete Hilfsmittel und Dienstleistungen an.

Zadzwoń, aby uzyskać bezpłatną pomoc językową oraz odpowiednie wsparcie i usługi pomocnicze.

Rele pou w jwenn sèvis asistans lengwistik gratis ak èd epi sèvis oksilyè ki apwopriye.

Appelez pour obtenir des services gratuits d'assistance linguistique et des aides et services auxiliaires appropriés.

Ligue para serviços gratuitos de assistência linguística e auxiliares e serviços auxiliares adequados.

Chiama per i servizi di assistenza linguistica gratuiti e per gli ausili e i servizi ausiliari appropriati.

اتصل للحصول على خدمات المساعدة اللغوية المجانية والمساعدات والخدمات المناسبة.

برای خدمات کمک زبانی رایگان و کمک‌ها و خدمات امدادی مقتضی، تماس بگیرید.

Discrimination is against the law. Premera Blue Cross Blue Shield of Alaska (Premera) complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex, including sex characteristics, intersex traits, pregnancy or related conditions, sexual orientation, gender identity, and sex stereotypes. Premera does not exclude people or treat them less favorably because of race, color, national origin, age, disability, or sex. Premera provides people with disabilities reasonable modifications and free appropriate auxiliary aids and services to communicate effectively with us, such as qualified sign language interpreters and written information in other formats (large print, audio, accessible electronic formats, other formats). Premera provides free language assistance services to people whose primary language is not English, which may include qualified interpreters and information written in other languages. If you need reasonable modifications, appropriate auxiliary aids and services, or language assistance services, contact our Civil Rights Coordinator. If you believe that Premera has failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, or sex, you can file a grievance with: Civil Rights Coordinator — Complaints and Appeals, PO Box 91102, Seattle, WA 98111, Toll free: 855-332-4535, TTY: 711, Fax: 425-918-5592, Email AppealsDepartmentInquiries@Premera.com. You can file a grievance in person or by mail, fax, or email. If you need help filing a grievance, our Civil Rights Coordinator is available to help you. You can also file a civil rights complaint with the U.S. Department of Health and Human Services, Office for Civil Rights, electronically through the Office for Civil Rights Complaint Portal, available at <https://ocrportal.hhs.gov/ocr/portal/lobby.jsf>, or by mail or phone at: U.S. Department of Health and Human Services, 200 Independence Ave SW, Room 509F, HHH Building, Washington, D.C. 20201, 1-800-368-1019, 800-537-7697 (TDD). Complaint forms are available at <http://www.hhs.gov/ocr/office/file/index.html>.