

PHARMACY / MEDICAL POLICY – 5.01.647

Medical Necessity Criteria for Custom Open Formulary

Effective Date: **Jan. 1, 2027***
 Last Revised: Sep. 8, 2026
 Replaces: N/A

*View current version here.

RELATED MEDICAL POLICIES:

5.01.550 Pharmacotherapy of Arthropathies
 5.01.555 Pharmacologic Treatment of Interstitial Lung Disease
 5.01.556 Rituximab: Non-oncologic and Miscellaneous Uses
 5.01.563 Pharmacotherapy of Inflammatory Bowel Disease
 5.01.564 Pharmacotherapy of Miscellaneous Autoimmune Diseases
 5.01.575 Dupixent (dupilumab)
 5.01.607 Continuity of Coverage for Maintenance Medications
 5.01.628 Pharmacologic Treatment of Atopic Dermatitis
 5.01.645 Pharmacologic Treatment of Psoriatic Arthritis
 5.01.629 Pharmacologic Treatment of Psoriasis

Select a hyperlink below to be directed to that section.

[POLICY CRITERIA](#) | [DOCUMENTATION REQUIREMENTS](#) | [CODING](#)
[RELATED INFORMATION](#) | [EVIDENCE REVIEW](#) | [REFERENCES](#) | [HISTORY](#)

 Clicking this icon returns you to the hyperlinks menu above.

Introduction

Prior authorization and step therapy are a way to provide safe and effective drugs. In step therapy, at least one drug on the health plan's list of covered drugs (the formulary) needs to be tried first. This policy describes the plan's prior authorization and step therapy criteria for specific drugs in the plan's formulary. This policy applies only to one custom Open formulary plan. This policy contains separate criteria to be used only for members with this specific custom Open formulary plan.

Note: The Introduction section is for your general knowledge and is not to be taken as policy coverage criteria. The rest of the policy uses specific words and concepts familiar to medical professionals. It is intended for providers. A provider can be a person, such as a doctor, nurse, psychologist, or dentist. A provider also can be a place where medical care is given, like a hospital, clinic, or lab. This policy informs them about when a service may be covered.

Policy Coverage Criteria

This policy applies only to one Open formulary (Formulary ID: 6062; Rx Plan F1). This policy contains separate criteria to be used only for members with this custom Open formulary plan. Please check the member Plan booklet or member ID card to determine whether this policy criteria applies.

Site of Service (SOS) Medical Necessity criteria applies ONLY to medical benefit reviews. SOS Medical Necessity criteria does NOT apply to Alaska fully-insured members; refer to the infusion and injection drug Medical Necessity criteria only. Please contact Customer Service for more information.

We will review specific intravenous (IV) and injectable drugs for medical necessity for all ages.

For those aged 13 and older, we also will review the site of service for medical necessity. Site of service is defined as the location where the drug is administered, such as a hospital-based outpatient setting, an infusion center, a physician's office, or at home. Click [here](#) to be directed to the site of service review criteria.

Drugs subject to site of service review addressed in this policy are:

- Actemra (tocilizumab) IV
- Avsola (infliximab-axxq) IV
- Avtozma (tocilizumab-anoh) IV
- Cosentyx (secukinumab) IV
- Inflectra (infliximab-dyyb) IV
- Infliximab (Janssen – unbranded) IV
- Orencia (abatacept) IV
- Remicade (infliximab)
- Renflexis (infliximab-abda) IV
- Riabni (rituximab-arrx) IV
- Rituxan (rituximab) IV



- Ruxience (rituximab-pvvr) IV
- Tofidence (tocilizumab-bavi) IV
- Truxima (rituximab-abbs) IV
- Tyenne (tocilizumab-aazg) IV

Site of Service Administration	Medical Necessity
Medically necessary sites of service • Physician's office • Infusion center • Home infusion	IV infusion and injection therapy of various medical or biologic agents will be covered in the most appropriate, safe and cost-effective site: • These are the preferred medically necessary sites of service for specified drugs.
Hospital-based outpatient setting • Outpatient hospital IV infusion department • Hospital-based outpatient clinical level of care	IV infusion and injection therapy of various medical or biologic agents will be covered in the most appropriate, safe and cost-effective site. This site is considered medically necessary for the first 90 days for the following: • The initial course of infusion or injection of a pharmacologic or biologic agent OR • Re-initiation of an agent after 6 months or longer following discontinuation of therapy* *This does not include when standard dosing between infusions is 6 months or longer. This site is considered medically necessary when there is no outpatient infusion center within 50 miles of the individual's home and there is no contracted home infusion agency that will travel to their home, or a hospital is the only place that offers infusions or injections of this drug. This site is considered medically necessary only when the individual has a clinical condition which puts him or her at



Site of Service Administration	Medical Necessity
	increased risk of complications for infusions or injections, including any 1 of the following: • Known cardiac condition (e.g., symptomatic cardiac arrhythmia) or pulmonary condition (e.g., significant respiratory disease, serious obstructive airway disease, %FVC less than or equal to 40%) that may increase the risk of an adverse reaction • Unstable renal function which decreases the ability to respond to fluids • Difficult or unstable vascular access • Acute mental status changes or cognitive conditions that impact the safety of infusion or injection therapy • A known history of severe adverse drug reactions and/or anaphylaxis to prior treatment with a related or similar drug This site is considered medically necessary when the individual has cytokine release syndrome (CRS) and all the following are met: • CRS is grade 3 or 4 as evidenced by ALL the following: ○ Temperature at least 38 °C ○ Hypotension that requires 1 or more vasopressors ○ Hypoxia that requires oxygen through a high-flow nasal cannula, face mask, non-rebreather mask, or Venturi mask OR positive pressure (continuous positive airway pressure [CPAP], bilevel positive airway pressure [BiPAP], intubation, or mechanical ventilation) AND • The individual will be admitted into an inpatient setting as soon as possible
Hospital-based outpatient setting • Outpatient hospital IV infusion department • Hospital-based outpatient clinical level of care	These sites are considered not medically necessary for infusion and injectable therapy services of various medical and biologic agents when the site-of-service criteria in this policy are not met.



Step therapy tiers are listed below; please refer to the Policy section for details.

First-line Agents					
TNF- α Inhibitors	IL-6 Inhibitors		CD20-Directed Antibodies		
• Adalimumab-aaty (SC) • Adalimumab-adaz (SC) • Adalimumab-adbm [NDCs starting with 00597] (SC) • Humira (AbbVie) [NDCs starting with 00074] (SC)	• Actemra (IV/SC) • Avtozma (IV/SC) • Tofidence (IV) • Tyenne (IV/SC)		• Riabni (IV) • Ruxience (IV)		
Second-line Agents					
TNF- α Inhibitors	α -4 Integrin Inhibitor	IL-1 Inhibitor	IL-6 Inhibitor	IL-17 Inhibitors	Sphingosine 1-Phosphate Receptor Modulator
• Abrilada (SC) • Adalimumab-aacf (SC) • Adalimumab-adbm [NDCs starting with 82009] (SC) • Adalimumab-bwwd (SC) • Adalimumab-fkjp (SC) • Adalimumab-ryvk (SC) • Amjevita (SC) • Cimzia (SC) • Cyltezo (SC) • Hadlima (SC) • Hulio (SC) • Humira (Cordavis) [NDCs starting	• Entyvio (SC)	• Kineret (SC)	• Kevzara (SC)	• Bimzelx (SC) • Cosentyx (IV/SC) • Siliq (SC)	• Zeposia (oral)

with 83457] (SC) • Hyrimoz (SC) • Immgolis (SC) • Infliximab (Janssen – unbranded) (IV) • Remicade (IV) • Simlandi (SC) • Simponi (SC) • Yusimry (SC)					
Second-line Agents					
Janus Kinase Inhibitor	IL-23 Inhibitor	T-Cell Costimulation Modulator	CD20-Directed Antibodies		
• Olumiant (oral)	• Ilumya (SC) • Icotyde (oral)	• Orencia (IV/SC)	• Rituxan (IV) • Rituxan Hycela (SC) • Truxima (IV)		

Drug	Medical Necessity
First-line CD20-Directed Antibodies	
• Riabni (rituximab-arrx) IV • Ruxience (rituximab-pvvr) IV	Riabni (rituximab-arrx) IV and Ruxience (rituximab-pvvr) IV are subject to review for site of service administration. Riabni (rituximab-arrx) IV and Ruxience (rituximab-pvvr) may be considered medically necessary for the treatment of rheumatoid arthritis when: • Treating moderately to severely active rheumatoid arthritis (e.g., at least 8 swollen and at least 8 tender joints) AND • Is administered in combination with methotrexate AND • Is used as a second-line therapy when either: ○ The individual has tried and failed any one of the first line therapies listed below: ▪ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597]



Drug	Medical Necessity
	▪ Enbrel (etanercept) ▪ Avsola (infliximab-axxq) OR Inflectra (infliximab-dyyb) OR Renflexis (infliximab-abda) ▪ Actemra (tocilizumab) OR Avtozma (tocilizumab-anoh) OR Tyenne (tocilizumab aazg) ▪ Tofacitinib OR tofacitinib extended-release OR Xeljanz (tofacitinib) OR Xeljanz XR (tofacitinib extended release) ▪ Rinvoq (upadacitinib) OR ○ The individual has had an inadequate response to methotrexate or other conventional synthetic disease-modifying anti-rheumatic drug (DMARD) AND ○ Is not a suitable candidate for treatment with TNF inhibitors (e.g., due to a recent [i.e., within 5 years] history of lymphoma or other malignancy; latent tuberculosis, and contraindication to chemoprophylaxis; or previous demyelinating disease)
First-line IL-6 Inhibitors	
• Actemra (tocilizumab) IV/SC • Avtozma (tocilizumab-anoh) IV/SC • Tofidence (tocilizumab-bavi) IV • Tyenne (tocilizumab-aazg) IV/SC	Actemra (tocilizumab) IV, Avtozma (tocilizumab-anoh) IV, Tofidence (tocilizumab-bavi) IV, and Tyenne (tocilizumab-aazg) IV are subject to review for site of service administration. Actemra (tocilizumab) IV/SC, Avtozma (tocilizumab-anoh) IV/SC, Tofidence (tocilizumab-bavi) IV, and Tyenne (tocilizumab-aazg) IV/SC may be considered medically necessary for the treatment of polyarticular juvenile idiopathic arthritis when: • The individual is aged 2 years or older AND • Has aggressive disease (Related Information) AND • Has had an inadequate response or intolerance to leflunomide, methotrexate, or sulfasalazine AND • Meets 1 of the following:



Drug	Medical Necessity
	- Has had an inadequate response or intolerance to Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] OR - Has heart failure or a previously treated lymphoproliferative disorder AND - Medication is being prescribed by or in consultation with a rheumatologist Actemra (tocilizumab) IV/SC, Avtozma (tocilizumab-anoh) IV/SC, Tofidence (tocilizumab-bavi) IV, and Tyenne (tocilizumab-aazg) IV/SC may be considered medically necessary for the treatment of moderate to severe rheumatoid arthritis when: - The individual is aged 18 years or older AND - Has had an inadequate response or intolerance to hydroxychloroquine, leflunomide, methotrexate, or sulfasalazine AND - Meets 1 of the following: - Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] OR - Has heart failure or a previously treated lymphoproliferative disorder AND - Medication is being prescribed by or in consultation with a rheumatologist
First-line TNF-α Inhibitors	
- Adalimumab-aaty SC - Adalimumab-adaz SC	Adalimumab-aaty, adalimumab-adaz, adalimumab-adbm [NDCs starting with 00597], and Humira (adalimumab) (AbbVie) [NDCs starting with 00074] may be considered



Drug	Medical Necessity
• Adalimumab-adbm [NDCs starting with 00597] SC • Humira (adalimumab) (AbbVie) [NDCs starting with 00074] SC	medically necessary for the treatment of ankylosing spondylitis when: • The individual is aged 18 years or older AND • Medication is being prescribed by or in consultation with a rheumatologist Adalimumab-aaty, adalimumab-adaz, adalimumab-adbm [NDCs starting with 00597], and Humira (adalimumab) (AbbVie) [NDCs starting with 00074] may be considered medically necessary for the treatment of Crohn's disease when: • The individual is aged 6 years or older AND • Meets 1 of the following: ○ Has tried 1 corticosteroid (e.g., methylprednisolone, prednisone, prednisolone, dexamethasone, budesonide, etc.) or is currently taking a corticosteroid medication OR ○ Has tried 1 other agent for Crohn's disease (e.g., azathioprine, 6-mercaptopurine, methotrexate, mesalamine extended-release [Pentasa formulation], etc.) OR ○ Has enterocutaneous (perianal or abdominal) or rectovaginal fistulas OR ○ Has had ileocolonic resection (to reduce the chance of Crohn's disease recurrence) AND • Medication is being prescribed by or in consultation with a gastroenterologist Adalimumab-aaty, adalimumab-adaz, adalimumab-adbm [NDCs starting with 00597], and Humira (adalimumab) (AbbVie) [NDCs starting with 00074] may be considered



Drug	Medical Necessity
	medically necessary for the treatment of hidradenitis suppurativa when: • The individual is aged 12 years or older AND • Has tried at least 1 other therapy (e.g., intralesional or oral corticosteroids, systemic antibiotics) AND • The medication is prescribed by or in consultation with a dermatologist Adalimumab-aaty, adalimumab-adaz, adalimumab-adbm [NDCs starting with 00597], and Humira (adalimumab) (AbbVie) [NDCs starting with 00074] may be considered medically necessary for the treatment of plaque psoriasis when: • The individual is aged 18 years or older AND • Has a diagnosis of chronic plaque psoriasis involving at least 10% of the body surface area (BSA) ○ Exception: This may be granted when ANY of the following are true: ▪ There is extensive recalcitrant facial involvement OR ▪ There is pustular involvement of the hands and feet OR ▪ There is genital involvement which interferes with normal sexual function AND • Has a history of an adequate trial and treatment failure with at least 1 approved systemic therapy (e.g., methotrexate, cyclosporine, acitretin or psoralen plus ultraviolet A light [PUVA]) unless contraindicated or not tolerated AND • Medication is being prescribed by or in consultation with a dermatologist

Drug	Medical Necessity
	Adalimumab-aaty, adalimumab-adaz, adalimumab-adbm [NDCs starting with 00597], and Humira (adalimumab) (AbbVie) [NDCs starting with 00074] may be considered medically necessary for the treatment of polyarticular juvenile idiopathic arthritis when: • The individual is aged 2 years or older AND • Has aggressive disease (Related Information) AND • Meets 1 of the following: ○ Has had an inadequate response or intolerance to leflunomide, methotrexate, or sulfasalazine OR ○ Is being started on adalimumab concurrently with leflunomide, methotrexate, or sulfasalazine AND • Medication is being prescribed by or in consultation with a rheumatologist Adalimumab-aaty, adalimumab-adaz, adalimumab-adbm [NDCs starting with 00597], and Humira (adalimumab) (AbbVie) [NDCs starting with 00074] may be considered medically necessary for the treatment of active psoriatic arthritis when: • The individual is aged 18 years or older AND • Medication is being prescribed by or in consultation with a rheumatologist or dermatologist Adalimumab-aaty, adalimumab-adaz, adalimumab-adbm [NDCs starting with 00597], and Humira (adalimumab) (AbbVie) [NDCs starting with 00074] may be considered medically necessary for the treatment of pyoderma gangrenosum when:



Drug	Medical Necessity
	- The individual is aged 18 years or older AND - Has not responded to 1 standard non-biologic therapy (e.g., oral corticosteroids, systemic cyclosporine, topical tacrolimus, etc.) AND - The medication is prescribed by or in consultation with a dermatologist Adalimumab-aaty, adalimumab-adaz, adalimumab-adbm [NDCs starting with 00597], and Humira (adalimumab) (AbbVie) [NDCs starting with 00074] may be considered medically necessary for the treatment of moderate to severe rheumatoid arthritis when: - The individual is aged 18 years or older AND - Has not responded to or does not tolerate hydroxychloroquine, leflunomide, methotrexate, or sulfasalazine AND - Medication is being prescribed by or in consultation with a rheumatologist Adalimumab-aaty, adalimumab-adaz, adalimumab-adbm [NDCs starting with 00597], and Humira (adalimumab) (AbbVie) [NDCs starting with 00074] may be considered medically necessary for the treatment of sarcoidosis when: - The individual is aged 18 years or older AND - Has tried and had an inadequate response or intolerance to 1 corticosteroid AND - Has tried and had an inadequate response or intolerance to 1 immunosuppressive medication (e.g., methotrexate, leflunomide, azathioprine, mycophenolate, cyclosporine, chlorambucil, cyclophosphamide, thalidomide, or chloroquine)



Drug	Medical Necessity
	AND - The medication is prescribed by or in consultation with a pulmonologist, ophthalmologist, or dermatologist Adalimumab-aaty, adalimumab-adaz, adalimumab-adbm [NDCs starting with 00597], and Humira (adalimumab) (AbbVie) [NDCs starting with 00074] may be considered medically necessary for the treatment of ulcerative colitis when: - The individual is aged 5 years or older AND - Medication is being prescribed by or in consultation with a gastroenterologist Adalimumab-aaty, adalimumab-adaz, adalimumab-adbm [NDCs starting with 00597], and Humira (adalimumab) (AbbVie) [NDCs starting with 00074] may be considered medically necessary for the treatment of non-infectious intermediate uveitis, posterior uveitis, or panuveitis when: - The individual is aged 2 years or older AND - Has tried 1 of the following therapies: - Periocular, intraocular, or systemic corticosteroids - Immunosuppressives AND - Medication is prescribed by or in consultation with an ophthalmologist
Avsola (infliximab-axxq) IV Inflectra (infliximab-dyyb) IV Renflexis (infliximab-abda) IV	Avsola (infliximab-axxq), Inflectra (infliximab-dyyb), and Renflexis (infliximab-abda) are subject to review for site of service administration. Avsola (infliximab-axxq), Inflectra (infliximab-dyyb), and Renflexis (infliximab-abda) may be considered medically necessary for the treatment of non-infectious intermediate uveitis, posterior uveitis, or panuveitis when:

Drug	Medical Necessity
	- The individual is aged 2 years or older AND - Has tried 1 of the following therapies: - Periocular, intraocular, or systemic corticosteroids - Immunosuppressives AND - Has had an inadequate response or intolerance to 1 of the following agents: - Adalimumab-aaty - Adalimumab-adaz - Adalimumab-adbm [NDCs starting with 00597] - Humira (adalimumab) (AbbVie) [NDCs starting with 00074] AND - The medication is prescribed by or in consultation with an ophthalmologist
Second-line α-4 Integrin Inhibitors	
Entyvio (vedolizumab) SC	Entyvio (vedolizumab) SC may be considered medically necessary for the treatment of adult individuals with Crohn's disease when: - The individual is aged 18 years or older AND - Meets 1 of the following: - Has tried one corticosteroid (e.g., methylprednisolone, prednisone, prednisolone, dexamethasone, budesonide, etc.) or is currently taking a corticosteroid medication OR - Has tried one other agent for Crohn's disease (e.g., azathioprine, 6-mercaptopurine, methotrexate, mesalamine extended-release [Pentasa formulation], etc.) OR - Has enterocutaneous (perianal or abdominal) or rectovaginal fistulas OR - Has had ileocolonic resection (to reduce the chance of Crohn's disease recurrence) AND - Meets 1 of the following:

Drug	Medical Necessity
	○ Has already received induction therapy with Entyvio (vedolizumab) IV AND has had a trial and treatment failure with 2 of the following: ▪ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] ▪ Cimzia (certolizumab pegol) ▪ Omvoh (mirikizumab-mrkz) SC ▪ Rinvoq (upadacitinib) ▪ Steqeyma (ustekinumab-stba) SC OR Yesintek (ustekinumab-kfce) SC ▪ Skyrizi (risankizumab-rzaa) SC on-body injector ▪ Tremfya (guselkumab) SC OR ○ Has already started on or is currently undergoing induction therapy with Entyvio (vedolizumab) IV AND • Entyvio (vedolizumab) SC is being prescribed by or in consultation with a gastroenterologist Entyvio (vedolizumab) SC may be considered medically necessary for the treatment of adult individuals with ulcerative colitis when: • The individual is aged 18 years or older AND • Meets 1 of the following: ○ Has already received induction therapy with Entyvio (vedolizumab) AND has had a trial and treatment failure with 2 of the following: ▪ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDC starting with 00597] ▪ Omvoh (mirikizumab-mrkz) SC ▪ Rinvoq (upadacitinib) ▪ Immgolis (golimumab-sldi) SC OR Simponi (golimumab) SC ▪ Skyrizi (risankizumab-rzaa) SC

Drug	Medical Necessity
	▪ Steqeyma (ustekinumab-stba) SC OR Yesintek (ustekinumab-kfce) SC ▪ Tremfya (guselkumab) SC ▪ Velsipity (etrasimod) ▪ Tofacitinib OR tofacitinib extended-release OR Xeljanz (tofacitinib) OR Xeljanz XR (tofacitinib extended-release) OR ○ Has already started on or is currently undergoing induction therapy with Entyvio (vedolizumab) IV AND • Entyvio (vedolizumab) SC is being prescribed by or in consultation with a gastroenterologist
Second-line CD20-Directed Antibodies	
• Rituxan (rituximab) IV • Rituxan Hycela (rituximab and hyaluronidase human) SC • Truxima (rituximab-abbs)	Rituxan (rituximab) IV and Truxima (rituximab-abbs) are subject to review for site of service administration. Rituxan (rituximab) IV, Rituxan Hycela (rituximab and hyaluronidase human) SC, and Truxima (rituximab-abbs) may be considered medically necessary for the treatment of rheumatoid arthritis when: • Treating moderately to severely active rheumatoid arthritis (e.g., at least 8 swollen and at least 8 tender joints) AND • Is administered in combination with methotrexate AND • Is used as a second-line therapy when either: ○ The individual has tried and failed any one of the first line therapies listed below: ▪ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] ▪ Enbrel (etanercept) ▪ Avsola (infliximab-axxq) OR Inflectra (infliximab-dyyb) OR Renflexis (infliximab-abda) ▪ Actemra (tocilizumab) OR Avtozma (tocilizumab-anoh) OR Tyenne (tocilizumab aazg)



Drug	Medical Necessity
	▪ Tofacitinib OR tofacitinib extended-release OR Xeljanz (tofacitinib) OR Xeljanz XR (tofacitinib extended release) ▪ Rinvoq (upadacitinib) OR ○ The individual has had an inadequate response to methotrexate or other conventional synthetic disease-modifying anti-rheumatic drug (DMARD) AND ○ Is not a suitable candidate for treatment with TNF inhibitors (e.g., due to a recent [i.e., within 5 years] history of lymphoma or other malignancy; latent tuberculosis, and contraindication to chemoprophylaxis; or previous demyelinating disease) AND • Has had an inadequate response or intolerance to Riabni (rituximab-arrx) AND Ruxience (rituximab-pvvr) ○ Exception: An exception may be granted for Rituxan Hycela (rituximab and hyaluronidase human) when documentation is provided of difficult venous access.
Second-line IL-1 Inhibitors	
Kineret (anakinra) SC	Kineret (anakinra) may be considered medically necessary for the treatment of moderate to severe rheumatoid arthritis when: • The individual is aged 18 years or older AND • Has not responded to or does not tolerate hydroxychloroquine, leflunomide, methotrexate, or sulfasalazine AND • Has had an inadequate response or intolerance to 2 of the following agents: ○ Actemra (tocilizumab) SC OR Avtozma (tocilizumab-anoh) SC OR Tyenne (tocilizumab-aazg) SC ○ Enbrel (etanercept) ○ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] ○ Rinvoq (upadacitinib)



Drug	Medical Necessity
	○ Tofacitinib OR tofacitinib extended-release OR Xeljanz (tofacitinib) OR Xeljanz XR (tofacitinib extended-release) AND • Medication is being prescribed by or in consultation with a rheumatologist
Second-line IL-6 Inhibitors	
Kevzara (sarilumab) SC	Kevzara (sarilumab) SC may be considered medically necessary for the treatment of polyarticular juvenile idiopathic arthritis when: • The individual weighs at least 63 kg AND • Has aggressive disease (Related Information) AND • Has had an inadequate response or intolerance to leflunomide, methotrexate, or sulfasalazine AND • Meets 1 of the following: ○ Has had an inadequate response or intolerance to 2 of the following drugs: ▪ Actemra (tocilizumab) SC OR Avtozma (tocilizumab-anoh) SC OR Tyenne (tocilizumab-aazg) SC ▪ Enbrel (etanercept) ▪ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] ▪ Rinvoq (upadacitinib) OR Rinvoq LQ (upadacitinib) ▪ Tofacitinib OR tofacitinib oral solution OR Xeljanz (tofacitinib) OR Xeljanz Oral Solution (tofacitinib) OR ○ Has heart failure, a previously treated lymphoproliferative disorder, a previous serious infection, or a demyelinating disorder AND • Medication is being prescribed by or in consultation with a rheumatologist

Drug	Medical Necessity
	Kevzara (sarilumab) may be considered medically necessary for the treatment of moderate to severe rheumatoid arthritis when: - The individual is aged 18 years or older AND - Has not responded to or does not tolerate hydroxychloroquine, leflunomide, methotrexate, or sulfasalazine AND - Meets 1 of the following: - Has had an inadequate response or intolerance to 2 of the following agents: - Actemra (tocilizumab) SC OR Avtozma (tocilizumab-anoh) SC OR Tyenne (tocilizumab-aazg) SC - Enbrel (etanercept) - Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] - Rinvoq (upadacitinib) - Tofacitinib OR tofacitinib extended-release OR Xeljanz (tofacitinib) OR Xeljanz XR (tofacitinib extended-release) OR - Has heart failure or a previously treated lymphoproliferative disorder AND - Medication is being prescribed by or in consultation with a rheumatologist
Second-line IL-17 Inhibitors	
Bimzelx (bimekizumab-bkzx) SC	Bimzelx (bimekizumab-bkzx) may be considered medically necessary for the treatment of ankylosing spondylitis when: - The individual is aged 18 years or older AND - Has had an inadequate response or intolerance to 1 of the following drugs: - Enbrel (etanercept) - Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597]



Drug	Medical Necessity
	○ Taltz (ixekizumab) AND • Medication is being prescribed by or in consultation with a rheumatologist Bimzelx (bimekizumab-bkzx) may be considered medically necessary for the treatment of hidradenitis suppurativa when: • The individual is aged 18 years or older AND • Has tried at least 1 other therapy (e.g., intralesional or oral corticosteroids, systemic antibiotics) AND • Has had an inadequate response or intolerance to 1 of the following agents: ○ Adalimumab-aaty ○ Adalimumab-adaz ○ Adalimumab-adbm [NDCs starting with 00597] ○ Cosentyx (secukinumab) ○ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] AND • Medication is prescribed by or in consultation with a dermatologist Bimzelx (bimekizumab-bkzx) may be considered medically necessary for the treatment of moderate to severe plaque psoriasis when: • The individual is aged 18 years or older AND • Has a diagnosis of chronic plaque psoriasis involving at least 10% of the body surface area (BSA) ○ Exception: This may be granted when ANY of the following are true: ▪ There is extensive recalcitrant facial involvement OR ▪ There is pustular involvement of the hands and feet



Drug	Medical Necessity
	OR - There is genital involvement which interferes with normal sexual function AND - Has a history of an adequate trial and treatment failure with at least 1 approved systemic therapy (e.g., methotrexate, cyclosporine, acitretin or psoralen plus ultraviolet A light [PUVA]) unless contraindicated or not tolerated AND - Has had an inadequate response or is intolerant to 1 of the following agents: - Enbrel (etanercept) - Adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] OR Humira (adalimumab) (AbbVie) [NDCs starting with 00074] - Otezla (apremilast) OR Otezla XR (apremilast extended-release) - Skyrizi (risankizumab-rzaa) SC - Sotyktu (deucravacitinib) - Steqeyma (ustekinumab-stba) SC OR Yesintek (ustekinumab-kfce) SC - Taltz (ixekizumab) - Tremfya (guselkumab) AND - Medication is being prescribed by or in consultation with a dermatologist Bimzelx (bimekizumab-bkzx) may be considered medically necessary for the treatment of active psoriatic arthritis when: - The individual is aged 18 years or older AND - Has had an inadequate response or intolerance to 1 of the following agents: - Enbrel (etanercept)

Drug	Medical Necessity
	○ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] ○ Otezla (apremilast) OR Otezla XR (apremilast extended-release) ○ Skyrizi (risankizumab-rzaa) ○ Sotyktu (deucravacitinib) ○ Steqeyma (ustekinumab-stba) SC OR Yesintek (ustekinumab-kfce) SC ○ Taltz (ixekizumab) ○ Tremfya (guselkumab) AND • Medication is being prescribed by or in consultation with a dermatologist or a rheumatologist
Cosentyx (secukinumab) IV/SC	Cosentyx (secukinumab) IV is subject to review for site of service administration. Cosentyx (secukinumab) IV/SC may be considered medically necessary for the treatment of ankylosing spondylitis when: • The individual is aged 12 years or older AND • Has had an inadequate response or intolerance to 2 of the following drugs: ○ Enbrel (etanercept) ○ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] ○ Rinvoq (upadacitinib) ○ Taltz (ixekizumab) ○ Tofacitinib OR tofacitinib extended-release OR Xeljanz (tofacitinib) OR Xeljanz XR (tofacitinib extended-release) AND • Medication is being prescribed by or in consultation with a rheumatologist

Drug	Medical Necessity
	Cosentyx (secukinumab) SC may be considered medically necessary for the treatment of moderate to severe plaque psoriasis when: • The individual is aged 6 years or older AND • Has a diagnosis of chronic plaque psoriasis involving at least 10% of the body surface area (BSA) ○ Exception: This may be granted when ANY of the following are true: ▪ There is extensive recalcitrant facial involvement OR ▪ There is pustular involvement of the hands and feet OR ▪ There is genital involvement which interferes with normal sexual function AND • Has a history of an adequate trial and treatment failure with at least 1 approved systemic therapy (e.g., methotrexate, cyclosporine, acitretin or psoralen plus ultraviolet A light [PUVA]) unless contraindicated or not tolerated AND • Has had an inadequate response or intolerance to 2 of the following agents: ○ Enbrel (etanercept) ○ Adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] OR Humira (adalimumab) (AbbVie) [NDCs starting with 00074] ○ Otezla (apremilast) OR Otezla XR (apremilast extended-release) ○ Skyrizi (risankizumab-rzaa) SC ○ Sotyktu (deucravacitinib) ○ Steqeyma (ustekinumab-stba) SC OR Yesintek (ustekinumab-kfce) SC ○ Taltz (ixekizumab) ○ Tremfya (guselkumab) AND



Drug	Medical Necessity
	- Medication is being prescribed by or in consultation with a dermatologist Cosentyx (secukinumab) IV/SC may be considered medically necessary for the treatment of active psoriatic arthritis when: - The individual is aged 2 years or older AND - Has had an inadequate response or intolerance to 2 of the following agents: - Enbrel (etanercept) - Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] - Otezla (apremilast) OR Otezla XR (apremilast extended-release) - Rinvoq (upadacitinib) OR Rinvoq LQ (upadacitinib) - Skyrizi (risankizumab-rzaa) - Sotyktu (deucravacitinib) - Steqeyma (ustekinumab-stba) SC OR Yesintek (ustekinumab-kfce) SC - Taltz (ixekizumab) - Tremfya (guselkumab) - Tofacitinib OR tofacitinib extended-release OR Xeljanz (tofacitinib) OR Xeljanz XR (tofacitinib extended-release) AND - Medication is being prescribed by or in consultation with a dermatologist or a rheumatologist
Siliq (brodalumab) SC	Siliq (brodalumab) may be considered medically necessary for the treatment of moderate to severe plaque psoriasis when: - The individual is aged 18 years or older AND - Has a diagnosis of chronic plaque psoriasis involving at least 10% of the body surface area (BSA) Exception: This may be granted when ANY of the following are true: - There is extensive recalcitrant facial involvement



Drug	Medical Necessity
	OR - There is pustular involvement of the hands and feet OR - There is genital involvement which interferes with normal sexual function AND - Has a history of an adequate trial and treatment failure with at least 1 approved systemic therapy (e.g., methotrexate, cyclosporine, acitretin or psoralen plus ultraviolet A light [PUVA]) unless contraindicated or not tolerated AND - Has had an inadequate response or is intolerant to 2 of the following agents: - Enbrel (etanercept) - Adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] OR Humira (adalimumab) (AbbVie) [NDCs starting with 00074] - Otezla (apremilast) OR Otezla XR (apremilast extended-release) - Skyrizi (risankizumab-rzaa) SC - Sotyktu (deucravacitinib) - Steqeyma (ustekinumab-stba) SC OR Yesintek (ustekinumab-kfce) SC - Taltz (ixekizumab) - Tremfya (guselkumab) AND - Medication is being prescribed by or in consultation with a dermatologist
Second-line IL-23 Inhibitors	
lcotyde (icotrokinra)	lcotyde (icotrokinra) may be considered medically necessary for the treatment of moderate to severe plaque psoriasis when: - The individual is aged 12 years or older AND - Weights at least 40 kg AND

Drug	Medical Necessity
	- Has a diagnosis of chronic plaque psoriasis involving at least 10% of the body surface area (BSA) Exception: This may be granted when ANY of the following are true: - There is extensive recalcitrant facial involvement OR - There is pustular involvement of the hands and feet OR - There is genital involvement which interferes with normal sexual function AND - Has a history of an adequate trial and treatment failure with at least 1 approved systemic therapy (e.g., methotrexate, cyclosporine, acitretin or psoralen plus ultraviolet A light [PUVA]) unless contraindicated or not tolerated AND - Has had an inadequate response or is intolerant to 2 of the following agents: - Enbrel (etanercept) - Adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] OR Humira (adalimumab) (AbbVie) [NDCs starting with 00074] - Otezla (apremilast) OR Otezla XR (apremilast extended-release) - Skyrizi (risankizumab-rzaa) SC - Sotyktu (deucravacitinib) - Steqeyma (ustekinumab-stba) SC OR Yesintek (ustekinumab-kfce) SC - Taltz (ixekizumab) - Tremfya (guselkumab) AND - Medication is being prescribed by or in consultation with a dermatologist AND - The dose is limited to 200 mg daily AND - The quantity is limited to 30 tablets per 30 days

Drug	Medical Necessity
Ilumya (tildrakizumab-asmn) SC	Ilumya (tildrakizumab-asmn) may be considered medically necessary for the treatment of moderate to severe plaque psoriasis when: • The individual is aged 18 years or older AND • Has a diagnosis of chronic plaque psoriasis involving at least 10% of the body surface area (BSA) ○ Exception: This may be granted when ANY of the following are true: ▪ There is extensive recalcitrant facial involvement OR ▪ There is pustular involvement of the hands and feet OR ▪ There is genital involvement which interferes with normal sexual function AND • Has a history of an adequate trial and treatment failure with at least 1 approved systemic therapy (e.g., methotrexate, cyclosporine, acitretin or psoralen plus ultraviolet A light [PUVA]) unless contraindicated or not tolerated AND • Has had an inadequate response or is intolerant to 2 of the following agents: ○ Enbrel (etanercept) ○ Adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] OR Humira (adalimumab) (AbbVie) [NDCs starting with 00074] ○ Otezla (apremilast) OR Otezla XR (apremilast extended-release) ○ Skyrizi (risankizumab-rzaa) SC ○ Sotyktu (deucravacitinib) ○ Steqeyma (ustekinumab-stba) SC OR Yesintek (ustekinumab-kfce) SC ○ Taltz (ixekizumab) ○ Tremfya (guselkumab) AND



Drug	Medical Necessity
	Medication is being prescribed by or in consultation with a dermatologist
Second-line Janus Kinase Inhibitors	
Olumiant (baricitinib) oral	Olumiant (baricitinib) may be considered medically necessary for the treatment of moderate to severe rheumatoid arthritis when: - The individual is aged 18 years or older AND - Has not responded to or does not tolerate hydroxychloroquine, leflunomide, methotrexate, or sulfasalazine AND - Has had an inadequate response or intolerance to 2 of the following agents: - Actemra (tocilizumab) SC OR Avtozma (tocilizumab-anoh) SC OR Tyenne (tocilizumab-aazg) SC - Enbrel (etanercept) - Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] - Rinvoq (upadacitinib) - Tofacitinib OR tofacitinib extended-release OR Xeljanz (tofacitinib) OR Xeljanz XR (tofacitinib extended-release) AND - Medication is being prescribed by or in consultation with a rheumatologist AND - The quantity is limited to 30 tablets per 30 days Note: The use of baricitinib in the setting of alopecia is considered cosmetic and is not covered by this policy.
Second-line Sphingosine 1-Phosphate Receptor Modulators	
Zeposia (ozanimod) oral	Zeposia (ozanimod) may be considered medically necessary for the treatment of adult individuals with ulcerative colitis when: - The individual is aged 18 years or older AND - Has had a trial and treatment failure with 2 of the following:



Drug	Medical Necessity
	○ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] ○ Omvoh (mirikizumab-mrkz) SC ○ Rinvoq (upadacitinib) ○ Immgolis (golimumab-sldi) SC OR Simponi (golimumab) SC ○ Skyrizi (risankizumab-rzaa) SC ○ Steqeyma (ustekinumab-stba) SC OR Yesintek (ustekinumab-kfce) SC ○ Tremfya (guselkumab) SC ○ Velsipity (etrasimod) ○ Tofacitinib OR tofacitinib extended-release OR Xeljanz (tofacitinib) OR Xeljanz XR (tofacitinib) AND • Zeposia (ozanimod) is prescribed by or in consultation with a gastroenterologist AND • The dose is limited to 0.92 mg per day
Second-line T-Cell Costimulation Modulators	
Orencia (abatacept) IV/SC	Orencia (abatacept) IV is subject to review for site of service administration. Orencia (abatacept) IV/SC may be considered medically necessary for the treatment of polyarticular juvenile idiopathic arthritis when: • The individual is aged 2 years or older AND • Has aggressive disease (Related Information) AND • Has had an inadequate response or intolerance to leflunomide, methotrexate, or sulfasalazine AND • Meets 1 of the following: ○ Has had an inadequate response or intolerance to 2 of the following drugs: ▪ Actemra (tocilizumab) SC OR Avtozma (tocilizumab-anoh) SC OR Tyenne (tocilizumab-aazg) SC



Drug	Medical Necessity
	▪ Enbrel (etanercept) ▪ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] ▪ Rinvoq (upadacitinib) OR Rinvoq LQ (upadacitinib) ▪ Tofacitinib OR tofacitinib oral solution OR Xeljanz (tofacitinib) OR Xeljanz Oral Solution (tofacitinib) OR ○ Has heart failure, a previously treated lymphoproliferative disorder, a previous serious infection, or a demyelinating disorder AND • Medication is being prescribed by or in consultation with a rheumatologist Orencia (abatacept) IV/SC may be considered medically necessary for the treatment of adults with active psoriatic arthritis when: • The individual is aged 18 years or older AND • Meets 1 of the following: ○ Has had an inadequate response or intolerance to 2 of the following agents: ▪ Enbrel (etanercept) ▪ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] ▪ Otezla (apremilast) OR Otezla XR (apremilast extended-release) ▪ Rinvoq (upadacitinib) OR Rinvoq LQ (upadacitinib) ▪ Skyrizi (risankizumab-rzaa) ▪ Sotyktu (deucravacitinib) ▪ Steqeyma (ustekinumab-stba) SC OR Yesintek (ustekinumab-kfce) SC ▪ Taltz (ixekizumab) ▪ Tremfya (guselkumab)

Drug	Medical Necessity
	▪ Tofacitinib OR tofacitinib extended-release OR Xeljanz (tofacitinib) OR Xeljanz XR (tofacitinib extended-release) OR ○ Has heart failure, a previously treated lymphoproliferative disorder, a previous serious infection, or a demyelinating disorder AND • Medication is being prescribed by or in consultation with a dermatologist or a rheumatologist Orencia (abatacept) IV/SC may be considered medically necessary for the treatment of moderate to severe rheumatoid arthritis when: • The individual is aged 18 years or older AND • Has not responded to or does not tolerate hydroxychloroquine, leflunomide, methotrexate, or sulfasalazine AND • Meets 1 of the following: ○ Has had an inadequate response or intolerance to 2 of the following agents: ▪ Actemra (tocilizumab) SC OR Avtozma (tocilizumab-anoh) SC OR Tyenne (tocilizumab-aazg) SC ▪ Enbrel (etanercept) ▪ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] ▪ Rinvoq (upadacitinib) ▪ Tofacitinib OR tofacitinib extended-release OR Xeljanz (tofacitinib) OR Xeljanz XR (tofacitinib extended-release) OR ○ Has heart failure, a previously treated lymphoproliferative disorder, a previous serious infection, or a demyelinating disorder AND

Drug	Medical Necessity
	Medication is being prescribed by or in consultation with a rheumatologist
Second-line TNF-α Inhibitors	
- Abrilada (adalimumab-afzb) SC - Adalimumab-aacf SC - Adalimumab-adbm [NDCs starting with 82009] SC - Adalimumab-bwwd SC - Adalimumab-fkjp SC - Adalimumab-ryvk SC - Amjevita (adalimumab-atto) SC - Cyltezo (adalimumab-adbm) SC - Hadlima (adalimumab-bwwd) SC - Hulio (adalimumab-fkjp) SC - Humira (adalimumab) (Cordavis) [NDCs starting with 83457] SC - Hyrimoz (adalimumab-adaz) SC - Simlandi (adalimumab-ryvk) - Yusimry (adalimumab-aqvh) SC	Abrilada (adalimumab-afzb), adalimumab-aacf, adalimumab-adbm [NDCs starting with 82009], adalimumab-bwwd, adalimumab-fkjp, adalimumab-ryvk, Amjevita (adalimumab-atto), Cyltezo (adalimumab-adbm), Hadlima (adalimumab-bwwd), Hulio (adalimumab-fkjp), Humira (adalimumab) (Cordavis) [NDCs starting with 83457], Hyrimoz (adalimumab-adaz), Simlandi (adalimumab-ryvk), and Yusimry (adalimumab-aqvh) may be considered medically necessary for the treatment of ankylosing spondylitis when: - The individual is aged 18 years or older AND - Has had an inadequate response or intolerance to ALL the following agents: - Adalimumab-aaty - Adalimumab-adaz - Adalimumab-adbm [NDCs starting with 00597] - Humira (adalimumab) (AbbVie) [NDCs starting with 00074] AND - Medication is being prescribed by or in consultation with a rheumatologist Abrilada (adalimumab-afzb), adalimumab-aacf, adalimumab-adbm [NDCs starting with 82009], adalimumab-bwwd, adalimumab-fkjp, adalimumab-ryvk, Amjevita (adalimumab-atto), Cyltezo (adalimumab-adbm), Hadlima (adalimumab-bwwd), Hulio (adalimumab-fkjp), Humira (adalimumab) (Cordavis) [NDCs starting with 83457], Hyrimoz (adalimumab-adaz), Simlandi (adalimumab-ryvk), and Yusimry (adalimumab-aqvh) may be considered medically necessary for the treatment of Crohn's disease when: - The individual is aged 6 years or older AND - Meets 1 of the following:

Drug	Medical Necessity
	○ Has tried 1 corticosteroid (e.g., methylprednisolone, prednisone, prednisolone, dexamethasone, budesonide, etc.) or is currently taking a corticosteroid medication OR ○ Has tried 1 other agent for Crohn's disease (e.g., azathioprine, 6-mercaptopurine, methotrexate, mesalamine extended-release [Pentasa formulation], etc.) OR ○ Has enterocutaneous (perianal or abdominal) or rectovaginal fistulas OR ○ Has had ileocolonic resection (to reduce the chance of Crohn's disease recurrence) AND • Has had an inadequate response or intolerance to ALL the following agents: ○ Adalimumab-aaty ○ Adalimumab-adaz ○ Adalimumab-adbm [NDCs starting with 00597] ○ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] AND • Medication is being prescribed by or in consultation with a gastroenterologist Abrilada (adalimumab-afzb), adalimumab-aacf, adalimumab-adbm [NDCs starting with 82009], adalimumab-bwwd, adalimumab-fkjp, adalimumab-ryvk, Amjevita (adalimumab-atto), Cyltezo (adalimumab-adbm), Hadlima (adalimumab-bwwd), Hulio (adalimumab-fkjp), Humira (adalimumab) (Cordavis) [NDCs starting with 83457], Hyrimoz (adalimumab-adaz), Simlandi (adalimumab-ryvk), and Yusimry (adalimumab-aqvh) may be considered medically necessary for the treatment of hidradenitis suppurativa when: • The individual is aged 12 years or older AND



Drug	Medical Necessity
	- Has tried at least 1 other therapy (e.g., intralesional or oral corticosteroids, systemic antibiotics) AND - Has had an inadequate response or intolerance to ALL the following agents: - Adalimumab-aaty - Adalimumab-adaz - Adalimumab-adbm [NDCs starting with 00597] - Humira (adalimumab) (AbbVie) [NDCs starting with 00074] AND - The medication is prescribed by or in consultation with a dermatologist Abrilada (adalimumab-afzb), adalimumab-aacf, adalimumab-adbm [NDCs starting with 82009], adalimumab-bwwd, adalimumab-fkjp, adalimumab-ryvk, Amjevita (adalimumab-atto), Cyltezo (adalimumab-adbm), Hadlima (adalimumab-bwwd), Hulio (adalimumab-fkjp), Humira (adalimumab) (Cordavis) [NDCs starting with 83457], Hyrimoz (adalimumab-adaz), Simlandi (adalimumab-ryvk), and Yusimry (adalimumab-aqvh) may be considered medically necessary for the treatment of plaque psoriasis when: - The individual is aged 18 years or older AND - Has a diagnosis of chronic plaque psoriasis involving at least 10% of the body surface area (BSA) Exception: This may be granted when ANY of the following are true: - There is extensive recalcitrant facial involvement OR - There is pustular involvement of the hands and feet OR - There is genital involvement which interferes with normal sexual function AND

Drug	Medical Necessity
	- Has a history of an adequate trial and treatment failure with at least 1 approved systemic therapy (e.g., methotrexate, cyclosporine, acitretin or psoralen plus ultraviolet A light [PUVA]) unless contraindicated or not tolerated AND - Has had an inadequate response or intolerance to ALL the following agents: - Adalimumab-aaty - Adalimumab-adaz - Adalimumab-adbm [NDCs starting with 00597] - Humira (adalimumab) (AbbVie) [NDCs starting with 00074] AND - Medication is being prescribed by or in consultation with a dermatologist Abrilada (adalimumab-afzb), adalimumab-aacf, adalimumab-adbm [NDCs starting with 82009], adalimumab-bwwd, adalimumab-fkjp, adalimumab-ryvk, Amjevita (adalimumab-atto), Cyltezo (adalimumab-adbm), Hadlima (adalimumab-bwwd), Hulio (adalimumab-fkjp), Humira (adalimumab) (Cordavis) [NDCs starting with 83457], Hyrimoz (adalimumab-adaz), Simlandi (adalimumab-ryvk), and Yusimry (adalimumab-aqvh) may be considered medically necessary for the treatment of polyarticular juvenile idiopathic arthritis when: - The individual is aged 2 years or older AND - Has aggressive disease (Related Information) AND - Has had an inadequate response or intolerance to leflunomide, methotrexate, or sulfasalazine AND - Has had an inadequate response or intolerance to ALL the following agents: - Adalimumab-aaty - Adalimumab-adaz

Drug	Medical Necessity
	○ Adalimumab-adbm [NDCs starting with 00597] ○ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] AND • Medication is being prescribed by or in consultation with a rheumatologist Abrilada (adalimumab-afzb), adalimumab-aacf, adalimumab-adbm [NDCs starting with 82009], adalimumab-bwwd, adalimumab-fkjp, adalimumab-ryvk, Amjevita (adalimumab-atto), Cyltezo (adalimumab-adbm), Hadlima (adalimumab-bwwd), Hulio (adalimumab-fkjp), Humira (adalimumab) (Cordavis) [NDCs starting with 83457], Hyrimoz (adalimumab-adaz), Simlandi (adalimumab-ryvk), and Yusimry (adalimumab-aqvh) may be considered medically necessary for the treatment of active psoriatic arthritis when: • The individual is aged 18 years or older AND • Has had an inadequate response or intolerance to ALL the following agents: ○ Adalimumab-aaty ○ Adalimumab-adaz ○ Adalimumab-adbm [NDCs starting with 00597] ○ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] AND • Medication is being prescribed by or in consultation with a dermatologist or a rheumatologist Abrilada (adalimumab-afzb), adalimumab-aacf, adalimumab-adbm [NDCs starting with 82009], adalimumab-bwwd, adalimumab-fkjp, adalimumab-ryvk, Amjevita (adalimumab-atto), Cyltezo (adalimumab-adbm), Hadlima (adalimumab-bwwd), Hulio (adalimumab-fkjp), Humira (adalimumab) (Cordavis) [NDCs starting with 83457], Hyrimoz (adalimumab-adaz), Simlandi (adalimumab-ryvk), and Yusimry (adalimumab-aqvh) considered medically necessary for the treatment of pyoderma gangrenosum when:

Drug	Medical Necessity
	- The individual is aged 18 years or older AND - Has not responded to 1 standard non-biologic therapy (e.g., oral corticosteroids, systemic cyclosporine, topical tacrolimus, etc.) AND - Has had an inadequate response or intolerance to ALL the following agents: - Adalimumab-aaty - Adalimumab-adaz - Adalimumab-adbm [NDCs starting with 00597] - Humira (adalimumab) (AbbVie) [NDCs starting with 00074] AND - Medication is prescribed by or in consultation with a dermatologist Abrilada (adalimumab-afzb), adalimumab-aacf, adalimumab-adbm [NDCs starting with 82009], adalimumab-bwwd, adalimumab-fkjp, adalimumab-ryvk, Amjevita (adalimumab-atto), Cyltezo (adalimumab-adbm), Hadlima (adalimumab-bwwd), Hulio (adalimumab-fkjp), Humira (adalimumab) (Cordavis) [NDCs starting with 83457], Hyrimoz (adalimumab-adaz), Simlandi (adalimumab-ryvk), and Yusimry (adalimumab-aqvh) may be considered medically necessary for the treatment of moderate to severe rheumatoid arthritis when: - The individual is aged 18 years or older AND - Has not responded to or does not tolerate hydroxychloroquine, leflunomide, methotrexate, or sulfasalazine AND - Has had an inadequate response or intolerance to ALL the following agents: - Adalimumab-aaty - Adalimumab-adaz - Adalimumab-adbm [NDCs starting with 00597]

Drug	Medical Necessity
	○ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] AND • Medication is being prescribed by or in consultation with a rheumatologist Abrilada (adalimumab-afzb), adalimumab-aacf, adalimumab-adbm [NDCs starting with 82009], adalimumab-bwwd, adalimumab-fkjp, adalimumab-ryvk, Amjevita (adalimumab-atto), Cyltezo (adalimumab-adbm), Hadlima (adalimumab-bwwd), Hulio (adalimumab-fkjp), Humira (adalimumab) (Cordavis) [NDCs starting with 83457], Hyrimoz (adalimumab-adaz), Simlandi (adalimumab-ryvk), and Yusimry (adalimumab-aqvh) may be considered medically necessary for the treatment of sarcoidosis when: • The individual is aged 18 years or older AND • Has tried and had an inadequate response or intolerance to 1 corticosteroid AND • Has tried and had an inadequate response or intolerance to 1 immunosuppressive medication (e.g., methotrexate, leflunomide, azathioprine, mycophenolate, cyclosporine, chlorambucil, cyclophosphamide, thalidomide, or chloroquine) AND • Has had an inadequate response or intolerance to ALL the following agents: ○ Adalimumab-aaty ○ Adalimumab-adaz ○ Adalimumab-adbm [NDCs starting with 00597] ○ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] AND • Medication is prescribed by or in consultation with a pulmonologist, ophthalmologist, or dermatologist

Drug	Medical Necessity
	Abrilada (adalimumab-afzb), adalimumab-aacf, adalimumab-adbm [NDCs starting with 82009], adalimumab-bwwd, adalimumab-fkjp, adalimumab-ryvk, Amjevita (adalimumab-atto), Cyltezo (adalimumab-adbm), Hadlima (adalimumab-bwwd), Hulio (adalimumab-fkjp), Humira (adalimumab) (Cordavis) [NDCs starting with 83457], Hyrimoz (adalimumab-adaz), Simlandi (adalimumab-ryvk), and Yusimry (adalimumab-aqvh) may be considered medically necessary for the treatment of ulcerative colitis when: • The individual is aged 5 years or older AND • Has had an inadequate response or intolerance to ALL the following agents: ○ Adalimumab-aaty ○ Adalimumab-adaz ○ Adalimumab-adbm [NDCs starting with 00597] ○ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] AND • Medication is being prescribed by or in consultation with a gastroenterologist Abrilada (adalimumab-afzb), adalimumab-aacf, adalimumab-adbm [NDCs starting with 82009], adalimumab-bwwd, adalimumab-fkjp, adalimumab-ryvk, Amjevita (adalimumab-atto), Cyltezo (adalimumab-adbm), Hadlima (adalimumab-bwwd), Hulio (adalimumab-fkjp), Humira (adalimumab) (Cordavis) [NDCs starting with 83457], Hyrimoz (adalimumab-adaz), Simlandi (adalimumab-ryvk), and Yusimry (adalimumab-aqvh) may be considered medically necessary for the treatment of non-infectious intermediate uveitis, posterior uveitis, or panuveitis when: • The individual is aged 2 years or older AND • Has tried 1 of the following therapies: ○ Periocular, intraocular, or systemic corticosteroids ○ Immunosuppressives

Drug	Medical Necessity
	AND - Has had an inadequate response or intolerance to ALL the following agents: - Adalimumab-aaty - Adalimumab-adaz - Adalimumab-adbm [NDCs starting with 00597] - Humira (adalimumab) (AbbVie) [NDCs starting with 00074] AND - Medication is prescribed by or in consultation with an ophthalmologist
Infliximab (Janssen – unbranded) IV Remicade (infliximab) IV	Infliximab (Janssen – unbranded) and Remicade (infliximab) are subject to review for site of service administration. Infliximab (Janssen – unbranded) and Remicade (infliximab) may be considered medically necessary for the treatment of non-infectious intermediate uveitis, posterior uveitis, or panuveitis when: - The individual is aged 2 years or older AND - Has tried one of the following therapies: - Periocular, intraocular, or systemic corticosteroids - Immunosuppressives AND - Has had an inadequate response or intolerance to 1 of the following agents: - Adalimumab-aaty - Adalimumab-adaz - Adalimumab-adbm [NDCs starting with 00597] - Humira (adalimumab) (AbbVie) [NDCs starting with 00074] AND - Has had an inadequate response or intolerance to Avsola (infliximab-axxq), Inflectra (infliximab-dyyb), AND Renflexis (infliximab-abda)¹ AND - The medication is prescribed by or in consultation with an ophthalmologist

Drug	Medical Necessity
	¹Note: Only applies to individuals not previously treated with requested therapy.
Cimzia (certolizumab pegol) SC	Cimzia (certolizumab pegol) SC may be considered medically necessary for the treatment of ankylosing spondylitis when: - The individual is aged 18 years or older AND - Has had an inadequate response or intolerance to 2 of the following drugs: - Enbrel (etanercept) - Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] - Rinvoq (upadacitinib) - Taltz (ixekizumab) - Tofacitinib OR tofacitinib extended-release OR Xeljanz (tofacitinib) OR Xeljanz XR (tofacitinib extended-release) AND - Medication is being prescribed by or in consultation with a rheumatologist Cimzia (certolizumab pegol) may be considered medically necessary for the treatment of Crohn's disease when: - The individual is aged 18 years or older AND - Meets 1 of the following: - Has tried one corticosteroid (e.g., methylprednisolone, prednisone, prednisolone, dexamethasone, budesonide, etc.) or is currently taking a corticosteroid medication OR - Has tried one other agent for Crohn's disease (e.g., azathioprine, 6-mercaptopurine, methotrexate, mesalamine extended-release [Pentasa formulation], etc.) OR - Has enterocutaneous (perianal or abdominal) or rectovaginal fistulas

Drug	Medical Necessity
	OR ○ Has had ileocolonic resection (to reduce the chance of Crohn's disease recurrence) AND • Has had a trial and treatment failure with 1 the following: ○ Adalimumab-aaty ○ Adalimumab-adaz ○ Adalimumab-adbm [NDCs starting with 00597] ○ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] AND • Medication is being prescribed by or in consultation with a gastroenterologist Cimzia (certolizumab pegol) may be considered medically necessary for the treatment of plaque psoriasis when: • The individual is aged 18 years or older AND • Has a diagnosis of chronic plaque psoriasis involving at least 10% of the body surface area (BSA) ○ Exception: This may be granted when ANY of the following are true: ▪ There is extensive recalcitrant facial involvement OR ▪ There is pustular involvement of the hands and feet OR ▪ There is genital involvement which interferes with normal sexual function AND • Has a history of an adequate trial and treatment failure with at least 1 approved systemic therapy (e.g., methotrexate, cyclosporine, acitretin or psoralen plus ultraviolet A light [PUVA]) unless contraindicated or not tolerated AND • Has had an inadequate response or is intolerant to 2 of the following agents: ○ Enbrel (etanercept)

Drug	Medical Necessity
	○ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] ○ Otezla (apremilast) OR Otezla XR (apremilast extended-release) ○ Skyrizi (risankizumab-rzaa) SC ○ Sotyktu (deucravacitinib) ○ Steqeyma (ustekinumab-stba) SC OR Yesintek (ustekinumab-kfce) SC ○ Taltz (ixekizumab) ○ Tremfya (guselkumab) AND • Medication is being prescribed by or in consultation with a dermatologist Cimzia (certolizumab pegol) SC may be considered medically necessary for the treatment of polyarticular juvenile idiopathic arthritis when: • The individual is aged 2 years or older AND • Has aggressive disease (Related Information) AND • Has had an inadequate response or intolerance to leflunomide, methotrexate, or sulfasalazine AND • Has had an inadequate response or intolerance to 2 of the following drugs: ○ Actemra (tocilizumab) SC OR Avtozma (tocilizumab-anoh) SC OR Tyenne (tocilizumab-aazg) SC ○ Enbrel (etanercept) ○ Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] ○ Rinvoq (upadacitinib) OR Rinvoq LQ (upadacitinib) ○ Tofacitinib OR tofacitinib oral solution OR Xeljanz (tofacitinib) OR Xeljanz Oral Solution (tofacitinib)



Drug	Medical Necessity
	AND - Medication is being prescribed by or in consultation with a rheumatologist Cimzia (certolizumab pegol) may be considered medically necessary for the treatment of active psoriatic arthritis when: - The individual is aged 18 years or older AND - Has had an inadequate response or intolerance to 2 of the following agents: - Enbrel (etanercept) - Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] - Otezla (apremilast) OR Otezla XR (apremilast extended-release) - Rinvoq (upadacitinib) OR Rinvoq LQ (upadacitinib) - Skyrizi (risankizumab-rzaa) SC - Sotyktu (deucravacitinib) - Steqeyma (ustekinumab-stba) SC OR Yesintek (ustekinumab-kfce) SC - Taltz (ixekizumab) - Tremfya (guselkumab) - Tofacitinib OR tofacitinib extended-release OR Xeljanz (tofacitinib) OR Xeljanz XR (tofacitinib extended-release) AND - Medication is being prescribed by or in consultation with a dermatologist or a rheumatologist Cimzia (certolizumab pegol) SC may be considered medically necessary for the treatment of moderate to severe rheumatoid arthritis when: - The individual is aged 18 years or older AND



Drug	Medical Necessity
	- Has not responded to or does not tolerate hydroxychloroquine, leflunomide, methotrexate, or sulfasalazine AND - Has had an inadequate response or intolerance to 2 of the following agents: - Actemra (tocilizumab) SC OR Avtozma (tocilizumab-anoh) SC OR Tyenne (tocilizumab-aazg) SC - Enbrel (etanercept) - Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] - Rinvoq (upadacitinib) - Tofacitinib OR tofacitinib extended-release OR Xeljanz (tofacitinib) OR Xeljanz XR (tofacitinib extended-release) AND - Medication is being prescribed by or in consultation with a rheumatologist
- Immgolis (golimumab-sldi) SC - Simponi (golimumab) SC	Immgolis (golimumab-sldi) SC and Simponi (golimumab) SC may be considered medically necessary for the treatment of ankylosing spondylitis when: - The individual is aged 18 years or older AND - Has had an inadequate response or intolerance to 2 of the following drugs: - Enbrel (etanercept) - Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] - Rinvoq (upadacitinib) - Taltz (ixekizumab) - Tofacitinib OR tofacitinib extended-release OR Xeljanz (tofacitinib) OR Xeljanz XR (tofacitinib extended-release) AND - Medication is being prescribed by or in consultation with a rheumatologist

Drug	Medical Necessity
	Immgolis (golimumab-sldi) SC and Simponi (golimumab) SC may be considered medically necessary for the treatment of active psoriatic arthritis when: - The individual is aged 18 years or older AND - Has had an inadequate response or intolerance to 2 of the following agents: - Enbrel (etanercept) - Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] - Otezla (apremilast) OR Otezla XR (apremilast extended-release) - Rinvoq (upadacitinib) OR Rinvoq LQ (upadacitinib) - Skyrizi (risankizumab-rzaa) SC - Sotyktu (deucravacitinib) - Steqeyma (ustekinumab-stba) SC OR Yesintek (ustekinumab-kfce) SC - Taltz (ixekizumab) - Tremfya (guselkumab) - Tofacitinib OR tofacitinib extended-release OR Xeljanz (tofacitinib) OR Xeljanz XR (tofacitinib extended-release) AND - Medication is being prescribed by or in consultation with a dermatologist or a rheumatologist Immgolis (golimumab-sldi) SC and Simponi (golimumab) SC may be considered medically necessary for the treatment of moderate to severe rheumatoid arthritis when: - The individual is aged 18 years or older AND - Has not responded to or does not tolerate hydroxychloroquine, leflunomide, methotrexate, or sulfasalazine AND - Has had an inadequate response or intolerance to 2 of the following agents:

Drug	Medical Necessity
	- Actemra (tocilizumab) SC OR Avtozma (tocilizumab-anoh) SC OR Tyenne (tocilizumab-aazg) SC - Enbrel (etanercept) - Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] - Rinvoq (upadacitinib) - Tofacitinib OR tofacitinib extended-release OR Xeljanz (tofacitinib) OR Xeljanz XR (tofacitinib extended-release) AND - Medication is being prescribed by or in consultation with a rheumatologist Immgolis (golimumab-sldi) SC and Simponi (golimumab) may be considered medically necessary for the treatment of ulcerative colitis when: - The individual weighs at least 15 kg AND - Has had a trial and treatment failure with Humira (adalimumab) (AbbVie) [NDCs starting with 00074] OR adalimumab-aaty OR adalimumab-adaz OR adalimumab-adbm [NDCs starting with 00597] AND - Medication is being prescribed by or in consultation with a gastroenterologist

Drug	Investigational
As listed	The medications listed in this policy are subject to the product's US Food and Drug Administration (FDA) dosage and administration prescribing information. All other uses of the above-named agents when used in combination with each other or for conditions not outlined in this policy are considered investigational.



Drug	Not Medically Necessary
As listed	All other uses of the drugs for approved conditions listed in this policy are considered not medically necessary.

Length of Approval	
Approval	Criteria
Initial authorization	All drugs listed in policy may be approved up to 12 months.
Re-authorization criteria	All drugs listed in policy may be approved up to 12 months as long as the drug-specific coverage criteria are met, and chart notes demonstrate that the individual continues to show a positive clinical response to therapy.

Documentation Requirements
The patient's medical records submitted for review for all conditions should document that medical necessity criteria are met. The record should include the following: Office visit notes that contain the diagnosis, relevant history, physical evaluation and medication history

Coding

Code	Description
HCPs	
J0129	Injection, abatacept (Orencia), 10 mg
J0139	Injection, adalimumab (Humira) 1 mg
J0717	Injection, certolizumab pegol (Cimzia), 1 mg
J1745	Injection, infliximab, excludes biosimilar (Remicade or Janssen unbranded), 10 mg
J3245	Injection, tildrakizumab (Ilumya), 1 mg
J3247	Injection, secukinumab (Cosentyx), IV, 1 mg
J3262	Injection, tocilizumab (Actemra), 1 mg
J3590	Unclassified biologics (use to report: Amjevita, Bimzelx, Cyltezo, Hadlima, Hulio, Hyrimoz, Immogolis, Kevzara, Kineret, Siliq, Simlandi, Simponi SC, Yuflyma and Yusimry)



Code	Description
J9311	Injection, rituximab 10 mg and hyaluronidase (Rituxan Hycela)
J9312	Injection, rituximab (Rituxan), 10 mg
Q5103	Injection, infliximab-dyyb, biosimilar, (Inflectra), 10 mg
Q5104	Injection, infliximab-abda, biosimilar, (Renflexis), 10 mg
Q5115	Injection, rituximab-abbs, biosimilar, (Truxima), 10 mg
Q5119	Injection, rituximab-pvvr, biosimilar, (Ruxience), 10 mg
Q5121	Injection, infliximab-axxq, biosimilar, (Avsola), 10 mg
Q5123	Injection, rituximab-arrx, biosimilar, (Riabni), 10 mg
Q5133	Injection, tocilizumab-bavi (Tofidence), biosimilar, 1 mg
Q5135	Injection, tocilizumab-aazg (Tyenne), biosimilar, 1 mg
Q5140	Injection, adalimumab-fkjp (Hulio), biosimilar, 1 mg
Q5141	Injection, adalimumab-aaty, biosimilar (Yuflyma), 1 mg
Q5142	Injection, adalimumab-ryvk biosimilar (Simlandi), 1 mg
Q5143	Injection, adalimumab-adbm, biosimilar (Cyltezo), 1 mg
Q5144	Injection, adalimumab-aacf (Idacio), biosimilar, 1 mg
Q5145	Injection, adalimumab-afzb (Abrilada), biosimilar, 1 mg
Q5156	Injection, tocilizumab-anoh (Avtozma), biosimilar, 1 mg

Note: CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). HCPCS codes, descriptions and materials are copyrighted by Centers for Medicare Services (CMS).

Related Information

Consideration of Age

Age limits specified in this policy are determined according to FDA-approved indications, where applicable.



Medical Benefit

Actemra (tocilizumab) IV, Avsola (infliximab-axxq), Avtozma (tocilizumab-anoh) IV, Cosentyx (secukinumab) IV, Inflectra (infliximab-dyyb), Infliximab (Janssen – unbranded), Orencia (abatacept) IV, Remicade (infliximab), Renflexis (infliximab-abda), Riabni (rituximab-arrx), Rituxan (rituximab), Rituxan Hycela (rituximab and hyaluronidase human), Ruxience (rituximab-pvvr), Tofidence (tocilizumab-bavi) IV, Tyenne (tocilizumab-aazg) IV, and Truxima (rituximab-abbs) are managed through the medical benefit.

Medical / Pharmacy Benefit

Abrilada (adalimumab-afzb), Actemra (tocilizumab) SC, adalimumab-aacf (Idacio unbranded), adalimumab-aaty (Yuflyma unbranded), adalimumab-adaz (Hyrimoz unbranded), adalimumab-adbm (Cyltezo unbranded), adalimumab-fkjp (Hulio unbranded), adalimumab-ryvk (Simlandi unbranded), Amjevita (adalimumab-atto), Avtozma (tocilizumab-anoh) SC, Bimzelx (bimekizumab-bkzx), Cimzia (certolizumab pegol), Cosentyx (secukinumab SC, Cyltezo (adalimumab-adbm), Entyvio (vedolizumab) SC, Hadlima (adalimumab-bwwd), Hulio (adalimumab-fkjp), Humira (adalimumab) (AbbVie) [NDCs starting with 00074], Humira (adalimumab) (Cordavis) [NDCs starting with 83457], Hyrimoz (adalimumab-adaz), Ilumya (tildrakizumab-asmn), Immgolis (golimumab-sldi), Kevzara (sarilumab), Kineret (anakinra), Orencia (abatacept) SC, Siliq (brodalumab), Simlandi (adalimumab-ryvk), Simponi (golimumab), Tyenne (tocilizumab-aazg) SC, Yuflyma (adalimumab-aaty), and Yusimry (adalimumab-aqvh) are managed through both the pharmacy and medical benefit.

Pharmacy Benefit

Icotyde (icotrokinra), Olumiant (baricitinib) and Zeposia (ozanimod) are managed through the pharmacy benefit.

Polyarticular Juvenile Idiopathic Arthritis: Aggressive Disease

Polyarticular juvenile idiopathic arthritis (PJIA) can become relentlessly progressive in children aged less than 10 years. Older children with rheumatoid factor (RF) are more likely to have more severe disease, mimicking their adult counterparts with rheumatoid arthritis. Some of these



children may be HLA-DR4 positive and/or have anti-CCP antibodies, which may put them at risk for an aggressive disease course in the absence of appropriate therapy.

Evidence Review

Crohn's Disease (CD)

The American College of Gastroenterology indicates current therapeutic recommendations depend on disease location, disease severity, and the presence of disease-associated complications. Pharmacologic approaches include various 5-aminosalicylates (5-ASAs), corticosteroids, and immunosuppressants. While the effectiveness of the 5-ASAs is less than corticosteroids, their side effect profile is more favorable. Azathioprine and sulfasalazine are also associated with clinically significant long-term toxicity, according to the National Cooperative Crohn's Disease Study. Azathioprine, sulfasalazine, and prednisone have not been demonstrated to prevent recurrence of disease flares.

Surgical resection is a common occurrence in CD, with up to 57% of individuals requiring at least one surgery in any given year. Within 10 years of disease onset, 71% of individuals undergo this therapy.

The safety and efficacy of adalimumab (Humira) for the induction and/or maintenance of remission in individuals with moderately to severely active CD (Crohn's Disease Activity Index [CDAI] ≥ 220 and ≤ 450) was evaluated in four randomized placebo-controlled studies. Two of these studies evaluated Humira for induction of remission (defined as a CDAI < 150), one study in individuals who were TNF antagonist naïve (CLASSIC-I) and the other in individuals who had lost response or were intolerant to Remicade (GAIN). Two of these studies evaluated Humira for maintenance of remission, both studies in individuals who were TNF antagonist naïve (CLASSIC-II and CHARM).

In CLASSIC-I, 299 individuals with moderately to severely active CD, including individuals with draining fistulas, were randomized to two subcutaneous injections at Weeks 0 and 2 with Humira 40 mg/20 mg, 80 mg/40 mg, or 160 mg/80 mg or placebo. Enrollees were also able to maintain existing therapy with immunomodulatory agents, corticosteroids, and/or aminosalicylates. The primary efficacy endpoint was induction of remission (CDAI < 150) at Week 4. The rate of remission was significantly higher in the 160 mg/80 mg group (36%, $p=0.001$), but not for the 40 mg/20 mg (18%, $p=0.36$) or 80 mg/40 mg (24%, $p=0.06$) groups compared with

placebo (12%). Injection site reactions occurred more frequently in Humira-treated individuals; otherwise, adverse events occurred at similar frequencies in all four treatment groups.

In GAIN, 325 individuals with moderately to severely active CD who were intolerant of, who had lost response, or who had an inadequate response to Remicade were randomized to two subcutaneous injections at Weeks 0 and 2 with Humira 160 mg/80 mg or placebo. Primary efficacy endpoint was induction of remission (CDAI <150) at Week 4. Clinical response (decrease in CDAI score ≥ 70 or 100) at Week 4 was also assessed. More Humira-treated individuals (21%, $p < 0.001$) achieved clinical remission compared to those treated with placebo (7%). More Humira-treated individuals (52%, $p < 0.01$) achieved a clinical response-70 compared with the placebo group (34%).

A total of 276 individuals participating in CLASSIC-I enrolled in CLASSIC-II and received open-label Humira 40 mg subcutaneously at Weeks 0 (Week 4 of CLASSIC-I) and 2. Those individuals ($n=55$) in remission at both Week 0 and Week 4 were re-randomized to Humira 40 mg QOW, 40 mg QW, or placebo for 52 additional weeks. Individuals who were not in remission at both Weeks 0 and 4 were treated with open-label Humira 40 mg QOW. These individuals were allowed to have their dose increase to 40 mg QW for non-response or disease flare. The re-randomized individuals were also allowed to "escape" into this open-label arm with disease flare. The primary efficacy endpoint was maintenance of remission (CDAI <150) in randomized individuals through week 56. Of the 55 individuals randomized at Week 4, a greater proportion receiving Humira (79% of the Humira 40 mg QOW group and 83% of the 40 mg QW group, both $p < 0.05$) were in remission compared to the placebo group (44%). Of 204 individuals entering the open-label arm, 46% were in remission at Week 56. Humira was generally well-tolerated.

In CHARM, a total of 854 individuals with moderately to severely active CD were treated with open-label Humira 80 mg at Week 0 followed by 40 mg at Week 2 as induction therapy. At Week 4, individuals were stratified by clinical response (decrease of CDAI ≥ 70) and randomized to double-blind treatment with subcutaneous Humira 40 mg QOW, Humira 40 mg QW, or placebo weekly for 52 additional weeks. The proportion of randomized clinical responders achieving clinical remission at Weeks 26 and 56 were coprimary endpoints. At Week 4, 499/854 (58%) of individuals achieved a clinical response-70 and were randomized to Humira or placebo. The percentage of randomized responders in remission was significantly greater in the Humira 40 mg QOW and 40 mg QW groups compared to the placebo group at Week 26 (40%, 47%, and 17%, respectively; $p < 0.001$) and at Week 56 (36%, 41%, and 12%, respectively; $p < 0.001$). No significant differences in efficacy were observed between the two active treatment groups. Individuals who did not achieve clinical response after 12 weeks were unlikely to achieve response. The safety profile for Humira was consistent with previous experience with the drug.

More individuals receiving placebo (13.4%) discontinued treatment for an adverse event than those receiving Humira (6.9% in the 40 mg QOW and 4.7% in the 40 mg QW group).

Hidradenitis Suppurativa

Hidradenitis Suppurativa (HS) is an inflammatory skin disease affecting an estimated 1 to 4% of the world population. The main features of HS include painful and chronically recurring, deep-seated follicular nodules, papules, pustules, and abscesses, scarring, sinus tracts, and recurrent discharge. The area's most commonly affected are the under the arms, groin, buttocks, and under the breasts. The disease is variable and recurrent. It may occur as solitary or multiple lesions in one area, or in many areas. In more severe cases, there may be large areas of skin affected by recurrent, draining lesions.

The FDA approved Humira (adalimumab) to treat individuals with HS.

Two randomized, double-blind, placebo-controlled studies (Studies HS-I and II) evaluated the safety and efficacy of Humira in a total of 633 adult subjects with moderate to severe hidradenitis suppurativa (HS) with Hurley Stage II or III disease and with at least 3 abscesses or inflammatory nodules. In both studies, subjects received placebo or Humira at an initial dose of 160 mg at Week 0, 80 mg at Week 2, and 40 mg every week starting at Week 4 and continued through Week 11. Subjects used topical antiseptic wash daily. Concomitant oral antibiotic use was allowed in Study HS-II.

Both studies evaluated Hidradenitis Suppurativa Clinical Response (HiSCR) at Week 12. HiSCR was defined as at least a 50% reduction in total abscess and inflammatory nodule count with no increase in abscess count and no increase in draining fistula count relative to baseline (see Table below). Reduction in HS-related skin pain was assessed using a Numeric Rating Scale in individuals who entered the study with an initial baseline score of 3 or greater on a 11-point scale.

In both studies, a higher proportion of Humira than placebo-treated subjects achieved HiSCR (see [Table 1](#) below).



Table 1. Efficacy Results at 12 Weeks in Subjects with Moderate to Severe Hidradenitis Suppurativa

	HS Study I		HS Study II*	
	Placebo	Humira 40 mg Weekly	Placebo	Humira 40 mg Weekly
Hidradenitis Suppurativa Clinical Response (HiSCR)	N=154, 40 (26%)	N=153, 64 (42%)	N=163, 45 (28%)	N=163, 96 (59%)

*19.3% of subjects in Study HS-II continued baseline oral antibiotic during the study.

In both studies, from Week 12 to Week 35 (Period B), subjects who had received Humira were re-randomized to 1 of 3 treatment groups (Humira 40 mg every week, Humira 40 mg every other week, or placebo). Subjects who had been randomized to placebo were assigned to receive Humira 40 mg every week (Study HS-I) or placebo (Study HS-II).

During Period B, flare of HS, defined as $\geq 25\%$ increase from baseline in abscesses and inflammatory nodule counts and with a minimum of 2 additional lesions, was documented in 22 (22%) of the 100 subjects who were withdrawn from Humira treatment following the primary efficacy time point in two studies.

Juvenile Idiopathic Arthritis

Juvenile Idiopathic Arthritis (JIA) is the most common type of arthritis in children under the age of 17. It causes persistent joint pain, swelling, and stiffness. Some children may experience symptoms for only a few months, while others have symptoms for the rest of their lives. In some cases this disease can cause complications, such as growth problems and eye inflammation. Treatment usually focuses on controlling pain, improving function, and preventing joint damage.

JIA occurs when the body's immune system attacks its own cells and tissues. It is not clear why this happens, however, both heredity and environment seem to play a role. Many different blood tests are used to diagnose JIA. Examples of some are: erythrocyte sedimentation rate (ESR), anti-nuclear antibody, rheumatoid factor, cyclic citrullinated peptide (CCP).

Treatment modalities depend on the extent of the disease, and individual child's needs. Some children benefit from one medication; others may need a combination of a few different medications. Each drug comes with its own side-effect potential which needs to be taken into

consideration based on the child's overall health condition and needs. First-line therapy includes the nonsteroidal anti-inflammatory drugs (NSAIDs)-examples of which are: ibuprofen, naproxen, and others. NSAIDs help to reduce pain and swelling of the joints. Disease-modifying antirheumatic drugs (DMARDs) is another option for drug therapy and include methotrexate, sulfasalazine, and others may be used when NSAIDs alone fail. Their purpose is to slow the progression of JIA. Tumor necrosis factor (TNF) blockers, such as etanercept and adalimumab can help reduce pain, morning stiffness, and swollen joints.

Psoriasis

Psoriasis is a chronic, multifactorial, noncontagious skin disorder that affects about 2.1% of the US population and 1-3% of persons worldwide. About 4.5 million, or 1 in 65, Americans have psoriasis. Onset is typically between the ages of 15 and 35 and prevalence is slightly greater in women. It is also more common in some ethnic groups (Caucasians) than others (African American or Asians). A genetic component has also been identified. There are several forms of psoriasis, but plaque psoriasis (or psoriasis vulgaris) is the most common form of the disease, affecting about 80% of psoriatic individuals.

About 20-30% of people with psoriasis have cases that are considered moderate to severe (covering more than 3% of their body). Although not typically life-threatening, psoriasis can have a large impact on quality of life. Seventy-five percent of people with moderate to severe psoriasis report their disease has a moderate to large impact on their everyday lives. Individuals with palmar-plantar disease may have less than 3% involvement, but often have debilitating and recalcitrant disease. Further, approximately 7% of psoriatic individuals have concurrent arthritis (which may be particularly relevant to one's choice of therapy).

Psoriasis is a chronic immune-mediated inflammatory disease characterized by T-cell activation and accumulation in the epidermis and dermis, leading to abnormal differentiation and hyperproliferation of keratinocytes. Recent advances in the understanding of the cellular mechanisms underlying psoriasis have given rise to a generation of highly targeted biotechnologies for this indication.

As the severity of psoriasis ranges from mild to severe, with or without concurrent arthritis, available treatments lie along a spectrum from minimally invasive with a low risk of systemic side effects, to systemic therapy with a risk of potentially severe side effects. Non-invasive, topical treatments may also have significant side effects; for example, topical corticosteroids applied to large areas of skin may result in significant levels of systemic absorption. Many treatments have



a cumulative toxicity potential, but the benefit of prolonged remissions makes the use of the more potent treatments relatively attractive.

Topical therapy, usually corticosteroids, is recommended as first-line treatment in psoriasis because these products are easy to administer, inexpensive, and safe. However, application to large areas of involvement can be time-consuming, expensive, and messy. Most individuals with moderate to severe disease will not achieve clearance or long-term remission. Tachyphylaxis may also develop with long-term use of topical corticosteroids. In individuals who's moderate to severe psoriasis fails topical therapy, the therapeutic options that remain are systemic agents, phototherapy and biologics.

Approved systemic agents (methotrexate, cyclosporine, and acitretin) are highly effective in the treatment of psoriasis; however, these therapies have limitations due to serious toxicities that require monitoring. Methotrexate can cause hepatotoxicity. Methotrexate is also associated with bone marrow toxicity, severe pulmonary toxicity, and serious drug-drug interactions (e.g., trimethoprim-sulfamethoxazole). Cyclosporine is nephrotoxic and can cause interstitial fibrosis and renal tubular atrophy in individuals treated for more than two years. Hypertension, laboratory abnormalities (electrolytes, liver function tests, lipids), and numerous drug-drug interactions are also among the problems associated with cyclosporine. Because methotrexate and cyclosporine are potent immunosuppressive drugs, individuals are at increased risk of infections and malignancies, including skin cancers and lymphoproliferative disorders. Like all retinoids, acitretin is highly teratogenic, posing a long-lasting risk (up to three years) in women of childbearing potential. Elevation in liver function tests, hyperlipidemia, and mucocutaneous reactions are additional adverse events associated with acitretin. Systemic corticosteroids are generally avoided as they may be associated with severe exacerbations, both during and after treatment.

Phototherapy (e.g., UVB, narrowband UVB, PUVA) is used for individuals who fail topicals or those with disease too extensive for topical therapy. Phototherapy can be effective for many individuals, but may be inconvenient and time-consuming, if frequent office or clinic visits are required and the availability of specialized phototherapy clinics may be limited. Individuals with a durable medical equipment (DME) benefit may purchase a home unit for easier access. Cumulative exposure to PUVA is associated with an increased risk of squamous cell carcinoma and malignant melanoma.

Various other strategies using traditional therapies have also been used to maintain remission and decrease the risk of cumulative end-organ toxicities. Rotational therapy involves the use of a therapy for some time and then switching to another form of therapy. Combination therapy uses low-dosages of different treatments concurrently to minimize toxicity and enhance efficacy.



Traditionally, these strategies usually involve topicals, phototherapy, and systemics in various combinations.

NBUVB continues to appear a very effective therapy in terms of achievement of greater than or equal to 75% response, global assessment ("clear or almost clear"), and length of remission. While the long-term risks of PUVA, methotrexate, and cyclosporine use in psoriatic individuals have become more clearly identified, these data are not available for the biologics in this population.

In the first multicenter, randomized, double-blind, placebo-controlled study, 147 individuals received Humira 80 mg at week 0, then 40 mg every other week beginning week 1, Humira 80 mg at week 0 and 1, then 40 mg every week beginning at week 1, or placebo for 12 weeks, after which placebo individuals were crossed over to Humira 40 mg every other week in a 48-week open label extension trial. At week 12, 53% of individuals taking Humira every other week, 80% of individuals taking Humira weekly, and 4% of individuals taking placebo achieved 75% improvement in Psoriasis Area and Severity Index score (Pless than 0 .001). Responses were sustained for 60 weeks. Humira was safe and well tolerated in this population.

In the Phase III REVEAL study (Randomized Controlled Evaluation of adalimumab Every Other Week Dosing in Moderate to Severe Psoriasis TriAL), 1,212 individuals with moderate to severe chronic plaque psoriasis were randomized to treatment with Humira 80 mg at week 0, then 40 mg every other week beginning at week 1 or placebo. The trial was comprised of 3 periods, a 16-week, double-blind period for assessment of initial response; a 17-week open-label sustained response period, in which responders to either treatment (those achieving a PASI-75) received Humira 40 mg every other week; and a final 19-week, double-blind loss of response period, in which individuals receiving Humira throughout the previous 2 study periods were re-randomized to either Humira every other week or placebo. In the initial response phase, more Humira-treated individuals achieved a PASI-75 compared to those receiving placebo beginning at week 4 and at every visit throughout the 16-week evaluation period. At week 16, 71% of Humira- and 6.5% of placebo-treated individuals achieved a PASI-75 (Pless than 0.001). In Humira responders, mean PASI scores were maintained throughout the subsequent maintenance of response period (weeks 16-33) of the study. In the last period of the study examining loss of response, 28.4% of individuals re-randomized to placebo lost response by week 52 compared to 4.9% of individuals maintaining Humira (Pless than 0.001). Humira was generally well tolerated, and no unexpected adverse events were observed over the 52 weeks of the trial.

In a second Phase III trial, CHAMPION (Comparative Study of HUMIRA vs. Methotrexate vs. Placebo In PsOriasias Patients), 271 individuals were randomized to treatment with Humira 80 mg at week 0, then 40 mg every other week beginning at week 1 (n=108), methotrexate 7.5 mg x 2



weeks, 10 mg x 2 weeks, then 15 mg orally (n=110), or placebo (n=53) for a total of 16 weeks. At week 16, more Humira-treated individuals achieved a PASI-75 response (80%) than individuals receiving either methotrexate (36%, P less than 0.001) or placebo (19%, P less than 0.001). Similar results were observed for PASI-90 response and PGA "clear" or "minimal" response. Humira was generally well-tolerated, with a safety profile similar to that known for an arthritis population.

Psoriatic Arthritis

Psoriatic Arthritis (PsA) is characterized as a spondyloarthropathy associated with psoriasis. The true incidence is unknown and is variably reported to occur in 6-42% (25% is considered a reasonable estimate) of individuals with psoriasis, an immunologic skin disease which occurs in 2-3% of the general population. There is similarity in the histopathogenesis of PsA and RA, including the role of cytokines such as tumor necrosis factor alpha (TNF- α), although there are important differences as well. Several subsets of PsA have also been described. PsA is characterized by stiffness - both peripheral and spine inflammation and pain - joint deformities related to joint destruction, dactylitis, enthesitis (inflammation at insertion sites of tendons, ligaments, and joint capsule fibers), and psoriasis skin plaques. The course of PsA is variable, but the majority of individuals develop a chronic progressive form of the disease resulting in joint destruction, unless treated effectively. Although less well characterized than in RA, similar levels of disability, decreased quality of life, increased co-morbidities, and premature mortality are now being noted in long term registry studies.

Pharmacologic therapy combined with a physical rehabilitation program is the most effective available treatment for psoriatic arthritis (PsA). As with RA, early initiation of pharmacologic therapy is needed to avoid joint damage and disability.

NSAIDs have customarily been used in milder disease along with corticosteroids or traditional DMARDs. Moderate to severe disease requires the use of traditional DMARDs such as MTX, sulfasalazine, or the anti-TNF agents. Azathioprine and cyclosporine are rarely used. Retinoids, phototherapy, and topical and systemic corticosteroids have also been used to treat the skin manifestations of PsA. In January 2002, etanercept, a TNF- α inhibitor became the first therapy to be approved for the indication.

Other Spondyloarthropathies

The spondyloarthropathies (SpAs) are a heterogeneous set of disorders characterized by axial skeletal involvement and frequent association with the HLA-B27 antigen. Ankylosing spondylitis



(AS) is probably the most familiar spondyloarthropathy, which is characterized predominantly by progressive vertebral enthesitis and facet joint inflammation of the spine and sacroiliac joints, leading to eventual spine fusion and decreased range of motion, as well as peripheral joint synovitis, although much less than is seen in RA. Variations in incidence among different racial groups support the hypothesis of a genetic role in AS, as is also postulated in other arthropathies. In the United States, AS is believed to affect approximately 1-3 persons/1000, or about 350,000 to of 1 million individuals.

While peripheral arthritis is commonly seen in association with psoriasis, approximately 20-40% of individuals with PsA may have some degree of sacroiliitis with paravertebral ossification. The skin manifestations associated with the arthropathy are not necessarily widespread and may be localized.

About 20% of individuals with inflammatory bowel disease may have evidence of sacroiliitis and some 20% of these individuals may progress to spondylitis. The course of the spondylitis does not necessarily correlate with bowel inflammatory activity.

Treatment of mild spondyloarthropathy may be benefited by symptomatic therapy with NSAIDs, corticosteroids, or sulfasalazine. These agents have shown to have little clinical benefit in individuals with moderate to severe or progressive disease. The paucity of treatment options contrasts with the treatment of RA where there are several different categories of DMARDs (disease-modifying anti-rheumatic drugs) that are used alone or in combination to try and alter the natural history of the disease. Like PsA, etanercept became the first therapy approved by the FDA for the treatment of AS, followed by infliximab and adalimumab.

Pyoderma Gangrenosum

Pyoderma gangrenosum is an inflammatory disease with dermatologic manifestations including painful ulcerations with erythematous borders. It is presumed to be autoimmune in origin, though the mechanism is not well understood. Lesions usually develop at sites of minor skin injury, usually on the lower extremities. These lesions can grow in size and become necrotic. Underlying fasciitis may occasionally develop from them. Some individuals develop pustular, bullous or vegetative lesions. Other common sites are colostomies and paraneoplastic lesions in individuals with hematologic malignancies. Progress of the lesions is highly variable, and individual response to treatment is heterogeneous. Obesity, diabetes or edema may be contributing factors.

Due to the infrequent occurrence and heterogeneity of pyoderma gangrenosum, the treatment approach is empiric and individual specific. First-line options include topical tacrolimus, nicotine,

and 5-ASA, systemic corticosteroids and immunosuppressant agents such as azathioprine, cyclosporine, methotrexate and mycophenolate. When these approaches fail, biologic therapy is usually tried. Successful treatment with TNF inhibitors (etanercept, adalimumab, infliximab) has been reported.

Rheumatoid Arthritis

Rheumatoid Arthritis (RA) is a chronic, progressive, inflammatory, autoimmune disease affecting about 1% of the US adult population and occurs approximately three times more frequently in women than in men (ACR Subcommittee on Rheumatoid Arthritis Guidelines, 2002). Almost 80% of RA cases occur in individuals between 35 and 50 years of age (Kavanaugh and Lipsky, 1996); usually a time of peak social productivity. The underlying cause of RA is unknown, but the disease is characterized by persistent inflammation of the synovium, cartilage loss, and bone erosion in peripheral joints, usually in a symmetric fashion. This inflammation is believed to be mediated by both B- and T-cells and a variety of cytokines (messenger proteins), including tumor necrosis factor-alpha (TNF- α). Research has shown that joint damage occurs within the first two years of symptoms and diagnosis and progresses rapidly if not treated. Although RA primarily affects the joints, it is a systemic disease and does cause systemic and extra-articular clinical features (e.g., fever, fatigue, anorexia, weight loss, and anemia), which contribute to the significant work disability and impaired quality of life which occur. Individuals with RA also have earlier mortality than the general population averaging 7-10 years, primarily due to an increased risk of cardiovascular disease, infection, and lymphoma associated with more severe inflammation.

The American College of Rheumatology (ACR) has established clinical guidelines for the treatment of RA. While both non-pharmacologic (e.g., individual education, exercise, and physical and occupational therapy) and pharmacologic therapies are recommended, the mainstay of RA treatment is pharmacologic therapy. Pharmacologic management often consists of nonsteroidal anti-inflammatory drugs (NSAIDs), disease-modifying antirheumatic drugs (DMARDs) (including biologic response modifiers/cytokine antagonists), and/or corticosteroids. Because of the evidence showing that joint damage can occur early in the disease process, physicians are now encouraged to treat individuals more aggressively earlier by initiating a DMARD (or combinations of DMARDs) within three months of diagnosis.

Emerging evidence also suggests that the DMARD subclass of tumor necrosis factor-alpha (TNF- α) antagonists retard radiographic progression of the disease better than methotrexate (MTX), particularly in individuals with rapidly progressive disease. The predictive risk factor found to be most associated with this subset of individuals was a CRP ≥ 4.1 mg/dl. Other predictors are

currently being investigated. This should lead to improved ability for the clinician to determine the best DMARD for an individual; however, the choice will continue to be influenced by numerous factors, including but not limited to relative efficacy, convenience of administration, adverse effects, monitoring requirements, comorbidities, and cost.

Ulcerative Colitis (UC)

After positive reports in small open-label trials, the safety and efficacy of adalimumab (Humira) was assessed in a multicenter, double-blinded randomized controlled trial in individuals with moderate to severe ulcerative colitis who were anti-TNF naïve and on stable suppressive therapy with oral corticosteroids and/or immunomodulators. A total of 576 individuals were randomized to receive either placebo, high dose (HD), or low dose (LD) adalimumab. HD was 180/60/40/40mg and LD was 80/40/40/40mg of adalimumab at Weeks 0, 2, 4, 6, respectively. Clinical remission was defined as a Mayo score ≤ 2 with subscores no greater than 1. Secondary outcomes included absolute score decrease plus decrease in rectal bleeding subscore, proportion with mucosal healing, and proportion with mild disease (including physician global assessment [PGA], rectal bleeding, and stool frequency subscores). Because the European regulatory authorities wanted to include a LD of adalimumab, there were two parts to the study, a 1:1 with HD (n=186) and a 1:1:1 portion of the study (n=390); results were pulled from the latter.

Twice as many individuals reached clinical remission at Week 8 with HD ($p=0.031$) therapy, while LD individuals were not significantly different versus placebo. Of the secondary outcomes, subscores in rectal bleeding and PGA showed improvement with significance vs. placebo in the HD arm. Individuals with higher baseline CRP levels had less instances of remission, and higher placebo rates were seen in Canadian and Eastern European centers than those in the US. Discontinuation rates were similar in each arm, with UC being the most common reason. Injection site pain was minimal, and infection incidence was similar across groups, and malignancy was only seen in the placebo arm.

Toxicities of TNF- α Antagonists

All TNF- α antagonists have treatment-limiting toxicities. Some of the toxicities associated with these agents include Concomitant use of TNF- α inhibitors and MTX consistently scored better with respect to ACR scores, disease activity in 28 joints (DAS28) scores, radiographical progression and health assessment questionnaire (HAQ) scores compared to TNF- α inhibitor

monotherapy. The ACR70 scores ranged from 15-20% for all agents, with etanercept showing the highest treatment effect over the control group at three years in the TEMPO trial. While infliximab showed high efficacy at both 3mg/kg and 10mg/kg dosing every eight weeks, the ACR50, ACR70 scores, HAQ scores were slightly higher with 10mg/kg dosing. The DAS28 scores and HAQ scores varied from study to study, but over-all showed improvement over controls across the TNF- α inhibitor class at 12 weeks and greater. Radiographical changes are subject to interpretation by the individual investigator, even with standardized scoring, so comparing across the TNF- α inhibitor trials is not practical. However, of the studies that did assess radiographical progression of the disease, the overall rate of radiographical progression was slowed significantly with adalimumab, certolizumab, etanercept and infliximab compared to MTX therapy alone. In the three-year TEMPO trial, the scores for the etanercept + MTX arm showed reversal of radiographical progression, but this is debatable and requires further investigation. There is no radiographical progression data for golimumab, as they did not assess this in their clinical trials.

There have been no prospective trials evaluating safety among the TNF- α inhibitors. The risk of malignancies and serious infections has been studied to some depth retrospectively with the three older agents (adalimumab, etanercept and infliximab). The FDA did a meta-analysis of the available data in 2006 and found that the malignancy rates of individuals on TNF- α inhibitors are no higher than what is to be expected in this individual population. Another study done in 2007 found a higher incidence of cutaneous cancers among the TNF- α inhibitor treated individuals, irrespective of the agent. The newer agents are limited in their data breadth to demonstrate safety with respect to malignancies, but so far, they compare similarly to the older agents. Long-term safety evaluations are necessary to validate this finding.

With regards to serious infections and tuberculosis, there are higher rates of serious infections while on the TNF- α inhibitors, compared to MTX alone. However, the retrospective studies do not come to an agreement on the actual risk.

References

1. Alonso J, Perez A, Castillo J et al. Psoriatic arthritis: a clinical immunological and radiological study of 180 patients. *J Rheum* 1991; 30:245-250.
2. American College of Rheumatology Subcommittee on Rheumatoid Arthritis Guidelines. Guidelines for the management of rheumatoid arthritis: 2002 update. *Arthritis Rheum* 2002; 46:328-346.
3. Braun J et al. Treatment of active ankylosing spondylitis with infliximab: a randomized controlled multicenter trial. *Lancet* 2002; 359:1187-1193.



4. Breathnach S. Psoriatic arthritis: etiology and pathogenesis. In: Klippel JH, Dieppe PA, eds. Rheumatology. London: *Mosby-Yearbook Europe Limited*; 1994; 32.1-32.6.
5. Helliwell P et al. Psoriatic arthritis, In: Klippel JH, Dieppe PA, eds. Rheumatology. London: *Mosby-Yearbook Europe Limited*; 1994; 21.1-21.8.
6. Kavanaugh AF, Lipsky PE. Rheumatoid arthritis. In: Rich RR, Fleisher TA, Schwartz B et al, eds. Clinical Immunology: Principles and Practice. St. Louis, MO: Mosby-Yearbook, 1996; 1093-1116.
7. Plant MJ, Saklatvala J, Borg AA, Jones PW, Dawes PT. Measurement and prediction of radiological progression in early rheumatoid arthritis. *J Rheumatol* 1994; 21:1808-1813.
8. Spondylitis Association of America Website. What is spondylitis? Available at: <https://spondylitis.org/> Accessed February 19, 2026.
9. Van der Heijde DMFM, van Leeuwen MA, van Riel PLCM, et al. Biannual radiographic assessments of hands and feet in a three-year prospective follow-up of patients with early rheumatoid arthritis. *Arthritis Rheum* 1992; 35:26-34.
10. Van der Heijde. Joint erosions and the patient with early rheumatoid arthritis. *Br J Rheumatol*. 1995; 34 (suppl 2):74-78.
11. Baecklund E et al. Disease activity and risk of lymphoma in patients with rheumatoid arthritis: nested case-control study. *BMJ* 1998; 317:180-1.
12. Mease PJ, Gladman DD, Ritchlin CT, et al. Adalimumab for the treatment of patients with moderately to severely active psoriatic arthritis: results of a double-blind, randomized, placebo-controlled trial. *Arthritis Rheum*. 2005; 52(10):3279-3289.
13. Genovese MC, Mease PJ, Thomson GTD, et al. Adalimumab efficacy in patients with psoriatic arthritis who failed prior DMARD therapy (abstract). Presented at EULAR Scientific Meeting 2005, Vienna, Austria, 10 June 2005.
14. Davis J, Kivitz A, Schiff M, et al. The Atlas Study Group. Major clinical response and partial remission in ankylosing spondylitis subjects treated with adalimumab: The ATLAS Trial (abstract). Presented at the Annual Scientific Meeting of the American College of Rheumatology 2005, San Diego, CA, Annual Scientific Meeting of the American College of Rheumatology 2005, San Diego, CA, November 12-17, 2005.
15. Maksymowych WP, Rahman P, Keystone E, et al. The M03-606 Study Group. Efficacy of adalimumab in active ankylosing spondylitis (AS)-results of the Canadian AS study (abstract). Presented at the Annual Scientific Meeting of the American College of Rheumatology 2005, San Diego, CA, Annual Scientific Meeting of the American College of Rheumatology 2005, San Diego, CA, November 12-17, 2005.
16. Online database. Mayo Clinic: Diseases and Conditions: Juvenile Rheumatoid Arthritis. Available at: <https://www.mayoclinic.org/diseases-conditions/juvenile-idiopathic-arthritis/symptoms-causes/syc-20374082> Accessed February 19, 2026.
17. Gherghe AM, Ramiro S, Landewé R, Mihai C, Van der heijde D. Association of the different types of radiographic damage with physical function in patients with rheumatoid arthritis: analysis of the RAPID trials. *RMD Open*. 2016;2(1):e000219.
18. Hirata S, Li W, Kubo S, et al. Association of the multi-biomarker disease activity score with joint destruction in patients with rheumatoid arthritis receiving tumor necrosis factor-alpha inhibitor treatment in clinical practice. *Mod Rheumatol*. 2016;:1-7.
19. Centers for Disease Control and Prevention. (2016). Rheumatoid Arthritis. Available at: <http://www.cdc.gov/arthritis/basics/rheumatoid.htm> Accessed February 19, 2026.
20. Dugowson CE, Koepsell TD, Voigt LF, Bley L, Nelson JL, Daling JR. Rheumatoid arthritis in women. Incidence rates in group health cooperative, Seattle, Washington, 1987-1989. *Arthritis Rheum*. 1991;34(12):1502-1507. PubMed PMID: 1747134. Abstract.
21. Dominick KL, Ahern FM, Gold CH, Heller DA. Health-related quality of life among older adults with arthritis. *Health Qual Life Outcomes*. 2004;2:5. doi:10.1186/1477-7525-2-5. PubMed PMID: 14720300; PubMed Central PMCID: PMC324570.
22. Gabriel SE, Crowson CS, Campion ME, O'Fallon WM. Indirect and nonmedical costs among people with rheumatoid arthritis and osteoarthritis compared with nonarthritic controls. *J.Rheumatol*. 1997;24(1):43-48. PubMed PMID: 9002009. Abstract.



23. Singh, J. A., Saag, K. G., Bridges, S. L., Akl, E. A., Bannuru, R. R., Sullivan, M. C., Vaysbrot, E., McNaughton, C., Osani, M., Shmerling, R. H., Curtis, J. R., Furst, D. E., Parks, D., Kavanaugh, A., O'Dell, J., King, C., Leong, A., Matteson, E. L., Schousboe, J. T., Drevlow, B., Ginsberg, S., Grober, J., St. Clair, E. W., Tindall, E., Miller, A. S. and McAlindon, T. (2016), 2015 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Care Res*, 68: 1–25. doi: 10.1002/acr.22783
24. Hyrimoz (adalimumab-adaz). Prescribing Information. Sandoz Inc; Princeton, NJ. Revised October 2025.
25. Yuflyma (adalimumab-aaty). Prescribing Information. Celltrion USA, Inc; Jersey City, NJ. Revised October 2025.
26. Cyltezo (adalimumab-adbm). Prescribing Information. Boehringer Ingelheim Pharmaceuticals, Inc; Ridgefield, CT. Revised October 2025.
27. Hadlima (adalimumab-bwwd). Prescribing Information. Merck Sharp & Dohme Corp; Whitehouse Station, NJ. Revised October 2025.
28. Abrilada (adalimumab-afzb). Prescribing Information. Pfizer Inc; New York, NY. Revised December 2025.
29. Hulio (adalimumab-fkjp). Prescribing Information. Mylan Pharmaceuticals Inc; Morgantown, WV. Revised February 2025.
30. Yusimry (adalimumab-aqvh). Prescribing Information. Coherus BioSciences, Inc., Redwood City, California. Revised November 2025.
31. Simlandi (adalimumab-ryvk). Prescribing Information. Teva Pharmaceuticals. Parsippany, NJ. Revised October 2025.
32. Humira (adalimumab). Prescribing Information. AbbVie Inc. North Chicago, IL. Revised December 2025.
33. Bernell O, Lapidus A, Hellers G. Risk factors for surgery and postoperative recurrence in Crohn's disease. *Ann Surg* 2000;231:38-45.
34. Hanauer SB, Sandborn W et al. Practice guidelines: management of Crohn's disease in adults. *Am J Gastroenterol* 2001;96:635-643.
35. Loftus EV Jr, Schoenfeld P, Sanborn WJ. The epidemiology and natural history of Crohn's disease in population-based patient cohorts from North America: a systematic review. *Aliment Pharmacol Ther* 2002;16:51-60.
36. Hanauer SB, Sandborn WJ, Rutgeerts P et al. Human anti-tumor necrosis factor monoclonal antibody (adalimumab) in Crohn's disease: the CLASSIC-I trial. *Gastroenterology* 2006;130:323-333.
37. Sandborn WJ, Rutgeerts P, Enns RA et al. Adalimumab rapidly induces clinical remission and response in patients with moderate to severe Crohn's disease who had secondary failure to infliximab therapy: results of the GAIN study (abstract). ACG 2006: The American College of Gastroenterology Annual Scientific Meeting. Abstract No. 750566.
38. Sandborn WJ, Hanauer SB, Rutgeerts P et al. Adalimumab for maintenance treatment of Crohn's disease: results of the CLASSIC II trial. *Gut* 2007. Epub ahead of print February 13, 2007.
39. Colombel JF, Sandborn WJ, Rutgeerts P et al. Adalimumab for maintenance of clinical response and remission in patients with Crohn's disease: the CHARM trial. *Gastroenterology* 2007;132:52-65.
40. Panaccione R, Colombel JF, Sandborn WJ, et al. Adalimumab sustains clinical remission and overall clinical benefit after 2 years of therapy for Crohn's disease. *Aliment Pharmacol Ther*. 2010;31(12):1296-309.
41. Oussalah A, Lacotte C, Chevaux J et al. Long-term outcome of adalimumab therapy for ulcerative colitis with intolerance of lost response to infliximab: a single-centre experience. *Aliment Pharmacol Ther*, 2008: 966-72.
42. Peyrin-Biroulet L, Laclotte C, Roblin X, Bigard M. Adalimumab induction therapy for ulcerative colitis with intolerance or lost response to infliximab: an open-label study. *World J gastroenterol* 2007: 2328-32.
43. Afif W, Leighton J, Hanauer S, et al. Open-label study of adalimumab in patients with ulcerative colitis including those with prior loss of response or intolerance to infliximab. *Inflamm Bowel Dis*. 2009;1302-7.
44. Reinisch W, Sandborn W, Hommes D, et al. Adalimumab for induction of clinical remission in moderately to severely active ulcerative colitis: results of a randomized control trial. *Gut*; 2011:780-787.



45. Russell R, Wilson M, Loganathan S, et al. A British Society of Paediatric Gastroenterology, Hepatology and Nutrition survey of the effectiveness and safety of adalimumab in children with inflammatory bowel disease. *Aliment Pharmacol Ther*; 2011;946-953.
46. Feuerstein JD, Akbari M, Gifford E, et al. Systematic review: the quality of the scientific evidence and conflicts of interest in international inflammatory bowel disease practice guidelines. *Aliment Pharmacol Ther*. 2013 May;37(10):937-46. doi: 10.1111/apt.12290. Epub 2013 Apr 2.
47. Lichtenstein GR, Hanauer SB, Sandborn WJ, et al. Management of Crohn's disease in adults. *Am J Gastroenterol* 6 Jan 2009 [epub ahead of print]; doi: 10.1038/ajg.2008.168.
48. Kornbluth A, Sachar DB, and The Practice Parameters Committee of the American College of Gastroenterology. Ulcerative colitis practice guidelines in adults: American College of Gastroenterology, Practice Parameters Committee. *Am J Gastroenterol* 2010;105:501-523.
49. Dignass A, Van Assche G, Lindsay JO, et al. The second European evidence-based consensus on the diagnosis and management of Crohn's disease: current management. *J Crohn's Colitis* 2010;4:28-62.
50. Allen PB, Peyrin-Biroulet L. Moving towards disease modification in inflammatory bowel disease therapy. *Curr Opin Gastroenterol* 2013;29:397-404.
51. Cheifetz AS. Management of active Crohn's disease. *JAMA* 2013;309:2150-2158.
52. Girardin M, Manz M, Manser C, et al. First-line therapies in inflammatory bowel disease. *Digestion* 2012;86(suppl 1):6-10.
53. Ardizzone S, Cassinotti A, de Franchis R. Immunosuppressives and biologic therapy for ulcerative colitis. *Expert Opin Emerging Drugs* 2012;17:449-467.
54. Rentz AM, Skalicky AM, Burslem K, et al. The content validity of the PSS in patients with plaque psoriasis. *J Patient Rep Outcomes*. 2017;1(1):4
55. FACIT.org. Functional assessment of chronic illness therapy-fatigue. Available from: <https://www.facit.org/measures/FACIT-F>. Accessed February 19, 2026.
56. Actemra (tocilizumab). Prescribing Information. Genentech, Inc.; South San Francisco, CA. Revised December 2025.
57. Bimzelx (bimekizumab-bkzx). Prescribing Information. UCB, Inc.; Smyrna, GA. Revised November 2024.
58. Cimzia (certolizumab pegol). Prescribing Information. UCB, Inc.; Smyrna, GA. Revised September 2025.
59. Cosentyx (secukinumab). Prescribing Information. Novartis Pharmaceuticals Corporation; East Hanover, NJ. Revised April 2026.
60. Entvyio PEN (vedolizumab). Prescribing Information. Takeda Pharmaceuticals U.S.A., Inc.; Lexington, MA. Revised April 2024.
61. Icotyde (icotrokinra). Prescribing Information. Janssen Biotech, Inc.; Horsham, PA. Revised March 2026.
62. Ilumya (tildrakizumab-asmn). Prescribing Information. Sun Pharmaceutical Industries, Inc.; Cranbury, NJ. Revised December 2025.
63. Immgolis (golimumab-sldi). Prescribing Information. Accord BioPharma, Inc.; Raleigh, NC. Revised May 2026.
64. Kevzara (sarilumab). Prescribing Information. Sanofi-Aventis U.S. LLC; Bridgewater, NJ. Revised June 2024.
65. Kineret (anakinra). Prescribing Information. Swedish Orphan Biovitrum AB; Stockholm, Sweden. Revised October 2025.
66. Olumiant (baricitinib). Prescribing Information. Eli Lilly and Company; Indianapolis, IN. Revised June 2022.
67. Orencia (abatacept). Prescribing Information. Bristol-Myers Squibb Company; Princeton, NJ. Revised May 2024.
68. Otezla (apremilast). Prescribing Information. Amgen Inc.; Thousand Oaks, CA. Revised August 2025.
69. Siliq (brodalumab). Prescribing Information. Bausch Health Ireland, Limited; Dublin, Ireland. Revised August 2024.
70. Simponi (golimumab). Prescribing Information. Janssen Biotech, Inc.; Horsham, PA. Revised October 2025.
71. Truxima (rituximab-abbs). Prescribing Information. Teva Pharmaceuticals USA, Inc, Inc; North Wales, PA. Revised June 2025.



History

Date	Comments
01/01/25	New policy, approved December 10, 2024. Added coverage criteria for Abrilada (adalimumab-afzb), adalimumab-aacf (Idacio unbranded), adalimumab-aaty (Yuflyma unbranded), adalimumab-adaz (Hyrimoz unbranded), adalimumab-adbm (Cyltezo unbranded), adalimumab-fkjp (Hulio unbranded), adalimumab-ryvk (Simlandi unbranded), Amjevita (adalimumab-atto), Cyltezo (adalimumab-adbm), Hadlima (adalimumab-bwwd), Hulio (adalimumab-fkjp), Humira (adalimumab) (AbbVie) [NDCs starting with 00074], Humira (adalimumab) (Cordavis) [NDCs starting with 83457], Hyrimoz (adalimumab-adaz) (Cordavis) [NDCs starting with 83457], Hyrimoz (adalimumab-adaz) (Sandoz) [NDCs starting with 61314], Idacio (adalimumab-aacf), Simlandi (adalimumab-ryvk), Yuflyma (adalimumab-aaty), and Yusimry (adalimumab-aqvh). Clarified that the medications listed in this policy are subject to the product's FDA dosage and administration prescribing information. Added HCPCS codes J0135, J3590, Q5131, Q5132.
04/01/25	Annual Review, approved March 11, 2025. Updated Abrilada (adalimumab-afzb), adalimumab-aacf (Idacio unbranded), adalimumab-aaty (Yuflyma unbranded), adalimumab-fkjp (Hulio unbranded), Amjevita (adalimumab-atto), Hadlima (adalimumab-bwwd), Hulio (adalimumab-fkjp), Humira (adalimumab) (Cordavis) [NDCs starting with 83457], Hyrimoz (adalimumab-adaz), Idacio (adalimumab-aacf), Yuflyma (adalimumab-aaty), Yusimry (adalimumab-aqvh), adalimumab-adaz (Hyrimoz unbranded), adalimumab-adbm (Cyltezo unbranded), adalimumab-ryvk (Simlandi unbranded), Cyltezo (adalimumab-adbm), Simlandi (adalimumab-ryvk), and Humira (adalimumab) (AbbVie) [NDCs starting with 00074] to include an age requirement for the ankylosing spondylitis, polyarticular juvenile idiopathic arthritis (PJIA), rheumatoid arthritis, psoriatic arthritis, Crohn's disease, ulcerative colitis, pyoderma gangrenosum, and sarcoidosis coverage criteria.
07/01/25	Interim Review, approved June 23, 2025. Updated title of the policy from "Medical Necessity Criteria for Custom Incentive and Open Formularies" to "Medical Necessity Criteria for Custom Open and Preferred Formularies." Updated re-authorization duration of approval from 3 years to 12 months. Added coverage criteria for the following: Riabni (rituximab-arrx) IV, Rituxan (rituximab) IV, Rituxan Hycela (rituximab and hyaluronidase human) SC, Ruxience (rituximab-pvvr) IV and Truxima (rituximab-abbs) IV for rheumatoid arthritis (RA), Actemra (tocilizumab) IV/SC, Tofidence (tocilizumab-bavi) IV, and Tyenne (tocilizumab-aazg) IV/SC for polyarticular juvenile idiopathic arthritis (PJIA) and RA, Entyvio (vedolizumab) SC for Crohn's disease and ulcerative colitis (UC), Kineret (anakinra) SC for RA, Kevzara (sarilumab) SC for PJIA and RA, Bimzelx (bimekizumab-bkzx) for ankylosing spondylitis (AS), hidradenitis



Date	Comments
	suppurativa (HS), plaque psoriasis (PsO), and psoriatic arthritis (PsA), Cosentyx (secukinumab) IV/SC for AS, PsO, and PsA, Siliq (brodalumab) SC for PsO, Ilumya (tildrakizumab-asmn) SC for PsO, Olumiant (baricitinib) for RA, Zeposia (ozanimod) for UC, Orencia (abatacept) IV/SC for PJIA, PsA in adults, and RA, Cimzia (certolizumab pegol) SC for AS, Crohn's disease, PsO, PJIA, PsA, and RA, Simponi (golimumab) SC for AS, PsA, RA, and UC. Added the following HCPCS codes per the criteria update: Q5119 (Ruxience IV), Q5115 (Truxima IV), J3262 (Actemra), Q5133 (Tofidence IV), Q5135 (Tyenne), J3247 (Cosentyx), J0129 (Orencia), J0717 (Cimzia), and J3245 (Ilumya). Removed termed codes Q5131 and Q5132. Added HCPCS codes Q5140 (Hulio), Q5141 (Yuflyma), Q5142 (Simlandi), Q5143 (Cyltezo), Q5144 (Idacio), Q5145 (Abrilada), Added HCPCS codes Q5123 (Riabni), J9311 (Rituxan Hycela), and J9312 (Rituxan) to match criteria update.
10/01/25	Interim Review, approved September 9, 2025. Removed Idacio (adalimumab-aacf) from the policy as it has been removed from the market. The following policy change is effective January 2, 2026, following 90-day provider notification. Updated Cyltezo (adalimumab-adbm) from a preferred product to a non-preferred product. Updated adalimumab-aaty (Yuflyma unbranded) and Yuflyma (adalimumab-aaty) from a non-preferred product to a preferred product. Updated Zymfentra (infliximab-dyyb) from a preferred product to a non-preferred product in the Entyvio (vedolizumab) IV, Entyvio (vedolizumab) SC, and Zeposia (ozanimod) coverage criteria. Updated Avsola (infliximab-axxq) from a non-preferred product to a preferred product in Riabni (rituximab-arrr), Rituxan (rituximab), Rituxan Hycela (rituximab and hyaluronidase human), Ruxience (rituximab-pvvr), and Truxima (rituximab-abbs) criteria. Updated Riabni (rituximab-arrr) from a non-preferred product to a preferred product.
11/01/25	Interim Review, approved October 14, 2025. Added coverage criteria for Remicade (infliximab), brand infliximab, and Renflexis (infliximab-abda) for the treatment of certain individuals with uveitis. Added HCPCS codes Q5104 and J1745.
12/01/25	Interim Review, approved November 24, 2025. Updated Simponi (golimumab) ulcerative colitis coverage criteria to include treatment of certain pediatric individuals.
01/01/26	Interim Review, approved December 8, 2025. Title changed to Medical Necessity Criteria for Custom Open Formulary. Removed reference to the Preferred formulary (Formulary ID: 6064; Rx Plan G3) as it is no longer available. Removed HCPCS code J0135 that termed effective 01/01/25.
04/01/26	Annual Review, approved March 10, 2026. Added Otezla XR (apremilast extended-release) as a preferred alternative to Bimzelx (bimekizumab) for treatment of psoriasis and psoriatic arthritis, Cosentyx (secukinumab) for treatment of plaque psoriasis and psoriatic arthritis, Siliq (brodalumab) for treatment of plaque psoriasis, Ilumya (tildrakizumab) for treatment of plaque psoriasis, Orencia (abatacept) for treatment of psoriatic arthritis, Cimzia (certolizumab) for treatment of plaque psoriasis and psoriatic arthritis, and Simponi (golimumab) SC for treatment of active psoriatic arthritis. Updated Kevzara (sarilumab) coverage criteria for treatment of PJIA from individual is aged 18 years or older to the individual weighs at least 63 kg, and included Avtozma (tocilizumab-anoh) in list of preferred alternatives. Updated Simponi (golimumab) SC



Date	Comments
	for treatment of active psoriatic arthritis correcting the individual weighs at least 15 kg to the individual is aged 18 years or older. The following was approved December 1, 2025, and is an administrative update for Simponi (golimumab) for treatment of ulcerative colitis changing from the individual is aged 18 years of age to the individual weighs at least 15 kg. Updated criteria for Riabni, Ruxience, Truxima, Rituxan, Rituxan Hycela for treatment of rheumatoid arthritis to include Avtozma (tocilizumab-anoh) and Tyenne (tocilizumab aazg) in list of first line therapies. Added Avtozma (tocilizumab-anoh) IV/SC to the policy with same criteria as Actemra (tocilizumab) IV/SC. Updated criteria for Kineret (anakinra), Kevzara (sarilumab), Olumiant (baricitinib), and Simponi (golimumab) SC for treatment of rheumatoid arthritis to include Avtozma (tocilizumab-anoh) in list of preferred alternatives. Updated criteria for Orencia (abatacept), Cimzia (certolizumab pegol) for treatment of PJIA and rheumatoid arthritis to include Avtozma (tocilizumab-anoh) in list of preferred alternatives. Updated criteria for first line and second line adalimumab products for the treatment of PJIA to include individual has aggressive disease. Added coverage criteria for Avsola (infliximab-axxq) IV and Inflectra (infliximab-dyyb) for the treatment of non-infectious intermediate uveitis, posterior uveitis, or panuveitis. Updated criteria for Infliximab (Janssen-unbranded), Remicade (infliximab), and Renflexis (infliximab-abda) for treatment of uveitis changing trial from ALL preferred adalimumab products to one preferred adalimumab product.
06/01/26	Interim Review, approved May 12, 2026. Updated criteria for Cosentyx (secukinumab) and treatment of ankylosing spondylitis updating age requirement from 18 to 12 years or older. Removed coverage criteria for Yuflyma (adalimumab-aaty) as product has been discontinued. Updated Simlandi (adalimumab-ryvk), adalimumab-ryvk, adalimumab-adbm [NDCs starting with 82009] from preferred products to non-preferred products. Updated criteria for second line adalimumab products, Riabni (rituximab-arrx), Ruxience (rituximab-pvvr), Truxima (rituximab-abbs), Rituxan (rituximab), Rituxan Hycela (rituximab and hyaluronidase human), Actemra (tocilizumab), Avtozma (tocilizumab-anoh), Tofidence (tocilizumab-bavi), Tyenne (tocilizumab-aazg), Infliximab (Janssen – unbranded), Remicade (infliximab), Renflexis (infliximab-abda), Avsola (infliximab-axxq), Inflectra (infliximab-dyyb), Bimzelx (bimekizumab-bkzx), Cimzia (certolizumab pegol), Cosentyx (secukinumab), Entyvio (vedolizumab) SC, Kevzara (sarilumab), Kineret (anakinra), Ilumya (tildrakizumab-asmn), Olumiant (baricitinib), Orencia (abatacept), Siliq (brodalumab), Simponi (golimumab) SC, and Zeposia (ozanimod) updating preferred alternatives to list adalimumab-aaty, adalimumab-adaz, and adalimumab-adbm [NDCs starting with 00597]. Updated criteria for Bimzelx (bimekizumab-bkzx), Cimzia (certolizumab), Cosentyx (secukinumab), Orencia (abatacept), and Simponi (golimumab) SC treatment for psoriatic arthritis updating list of preferred alternatives to include Sotyktu (deucravacitinib). Updated criteria for Olumiant (baricitinib) adding the quantity is limited to 30 tablets per 30 days.
07/01/26	Interim Review, approved June 9, 2026. Added coverage criteria for Icotyde (icotrokinra) for the treatment of psoriasis in individuals aged 12 years or older and weigh at least 40 kg.



Date	Comments
08/01/26	Interim Review, approved July 27, 2026. Updated Cimzia, Cosentyx, Entyvio SC, Kevzara, Kineret, Olumiant, Orencia, Riabni, Rituxan, Rituxan Hycela, Ruxience, Truxima, Simponi SC, and Zeposia to include tofacitinib and tofacitinib extended-release as preferred alternatives where Xeljanz (tofacitinib) and Xeljanz XR (tofacitinib extended-release) are listed as preferred alternatives. Added adalimumab-bwwd to the policy with the same criteria as Hadlima (adalimumab-bwwd) and second-line adalimumab products.
10/01/26	Interim Review, approved September 8, 2026. Added Immgolis (golimumab-sldi) to the policy with the same criteria as Simponi (golimumab) SC. Updated Entyvio (vedolizumab) SC and Zeposia (ozanimod) adding Immgolis (golimumab-sldi) to list of preferred alternatives. Updated Renflexis (infliximab-abda) from a non-preferred product to a preferred product. Updated Riabni, Rituxan, Rituxan Hycela, Ruxience, and Truxima for the treatment of RA adding Renflexis (infliximab-abda) to list of preferred alternatives. Updated non-preferred infliximab products, infliximab (Janssen – unbranded) and Remicade (infliximab), adding Renflexis (infliximab-abda) to list of preferred alternatives. The following policy changes are effective January 1, 2027, following 90-day provider notification. Updated Truxima (rituximab-abbs) from a preferred product to a non-preferred product. Updated non-preferred rituximab products, Rituxan and Rituxan Hycela, removing Truxima from list of preferred rituximab products. Updated criteria for all non-preferred agents removing Stelara (ustekinumab) as a preferred alternative for treatment of Crohn's disease, plaque psoriasis, psoriatic arthritis, and ulcerative colitis.

Disclaimer: This medical policy is a guide in evaluating the medical necessity of a particular service or treatment. The Company adopts policies after careful review of published peer-reviewed scientific literature, national guidelines and local standards of practice. Since medical technology is constantly changing, the Company reserves the right to review and update policies as appropriate. Member contracts differ in their benefits. Always consult the member benefit booklet or contact a member service representative to determine coverage for a specific medical service or supply. CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). ©2026 Premera All Rights Reserved.

Scope: Medical policies are systematically developed guidelines that serve as a resource for Company staff when determining coverage for specific medical procedures, drugs or devices. Coverage for medical services is subject to the limits and conditions of the member benefit plan. Members and their providers should consult the member benefit booklet or contact a customer service representative to determine whether there are any benefit limitations applicable to this service or supply. This medical policy does not apply to Medicare Advantage.

