

MEDICAL POLICY – 7.01.160

Cartilage Implants for Joint Pain

BCBSA Ref. Policy: 7.01.160

Effective Date: **Nov. 6, 2026***
Last Revised: Jul. 14, 2026
Replaces: N/A

*Click here to view the current policy.

RELATED MEDICAL POLICIES:

7.01.48 Autologous Chondrocyte Implantation for Focal Articular Cartilage Lesions
7.01.570 Autografts and Allografts in the Treatment of Focal Articular Cartilage Lesions

Select a hyperlink below to be directed to that section.

[POLICY CRITERIA](#) | [CODING](#) | [RELATED INFORMATION](#)
[EVIDENCE REVIEW](#) | [REFERENCES](#) | [APPENDIX](#) | [HISTORY](#)

 Clicking this icon returns you to the hyperlinks menu above.

Introduction

Cartilage is the smooth, rubbery tissue that covers the ends of bones in a joint. It helps the joint move easily and without pain (articular cartilage). When this tissue is damaged, it does not heal well on its own. This can lead to pain, swelling, and trouble moving the joint. Cartilage implants are devices placed into the joint to replace or repair the damaged tissue. Some implants are made from natural materials, such as AGILI-C, while others are synthetic, such as Cartiva. These treatments are being studied to see if they can restore joint function and reduce symptoms. Cartilage implants (e.g., AGILI-C [natural], Cartiva [synthetic]) are considered investigational (unproven). There's not enough evidence to show they are effective.

Note: The Introduction section is for your general knowledge and is not to be taken as policy coverage criteria. The rest of the policy uses specific words and concepts familiar to medical professionals. It is intended for providers. A provider can be a person, such as a doctor, nurse, psychologist, or dentist. A provider also can be a place where medical care is given, like a hospital, clinic, or lab. This policy informs them about when a service may be covered.

Policy Coverage Criteria

Service	Investigational
Cartilage implants	Cartilage implants (e.g., AGILI-C, Cartiva) are considered investigational for the treatment of articular cartilage damage.

Coding

Code	Description
HCPCS	
L8641	Metatarsal joint implant
L8642	Hallux implant
L8699	Prosthetic Implant, not otherwise specified

Note: CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). HCPCS codes, descriptions and materials are copyrighted by Centers for Medicare Services (CMS).

Related Information

N/A

Evidence Review

Description

Articular cartilage damage, either from a focal lesion or diffuse osteoarthritis (OA), can result in disabling pain. Cartilage is a hydrogel, comprised mostly of water with collagen and glycosaminoglycans, that does not typically heal on its own. There is a need for improved treatment options. In 2016, a synthetic polyvinyl alcohol hydrogel disc received marketing approval by the United States (US) Food and Drug Administration (FDA) for the treatment of

degenerative or posttraumatic arthritis in the first metatarsophalangeal (MTP) joint. In 2022, a cartilage repair implant for knee cartilage and osteochondral defects was approved.

Background

Articular Cartilage Damage

Articular cartilage damage may present as focal lesions or as more diffuse OA. Cartilage is a biological hydrogel that is comprised mostly of water with collagen and glycosaminoglycans and does not typically heal on its own. OA or focal articular cartilage lesions can be associated with substantial pain, loss of function, and disability. OA is most frequently observed in the knees, hips, interphalangeal joints, first carpometacarpal joints, first MTP joint, and apophyseal (facet) joints of the lower cervical and lower lumbar spine. OA less commonly affects the elbow, wrist, shoulder, and ankle. Knee OA is the most common cause of lower-limb disability in adults over age 50, however, OA of the MTP joint with loss of motion (hallux rigidus) can also be severely disabling due to pain in the “toe-off” position of gait. An epidemiologic study found that OA of the first MTP joint may be present in as many as one in 40 people over the age of 50.¹

Treatment

Treatment for articular cartilage damage may include debridement, abrasion arthroplasty, microfracture, osteochondral autografting, and autologous chondrocyte implantation. Debridement involves the removal of the synovial membrane, osteophytes, loose articular debris, and diseased cartilage and is capable of producing symptomatic relief. Subchondral abrasion techniques attempt to restore the articular surface by inducing the growth of fibrocartilage into the chondral defect. Diffuse osteoarthritis of the knee, hip, shoulder or ankle may be treated with joint replacement.

Early-stage osteoarthritis of the first MTP joint is typically treated with conservative management, including pain medication and change in footwear. Failure of conservative management in individuals with advanced osteoarthritis of the MTP joint may be treated surgically. Cheilectomy (removal of bone osteophytes) and interpositional spacers with autograft or allograft have been used as temporary measures to relieve pain. Although partial or total joint replacement have been explored for MTP osteoarthritis, complications from bone loss, loosening, wear debris, implant fragmentation, and transfer metatarsalgia are not uncommon. Also, since the conversion of a failed joint replacement to arthrodesis has greater complications

and worse functional results than a primary arthrodesis (joint fusion), MTP arthrodesis is considered the most reliable and primary surgical option. Arthrodesis can lead to a pain-free foot, but the loss of mobility in the MTP joint alters gait, may restrict participation in running and other sports, and limits footwear options, leading to patient dissatisfaction. Transfer of stress and arthritis in an adjacent joint may also develop over time.

Because of the limitations of MTP arthrodesis, alternative treatments that preserve joint motion are being explored. Synthetic cartilage implants have been investigated as a means to reduce pain and improve function in individuals with hallux rigidus. Some materials such as silastic were found to fragment with use. Other causes of poor performance are the same as those observed with metal and ceramic joint replacement materials and include dislocation, particle wear, osteolysis, and loosening. Synthetic polyvinyl alcohol (PVA) hydrogels have water content and biomechanical properties similar to cartilage and they are biocompatible. Polyvinyl alcohol hydrogels have been used in a variety of medical products including soft contact lenses, artificial tears, hydrophilic nerve guides, and tissue adhesion barriers. This material is being evaluated for cartilage replacement due to the rubber elastic properties and, depending on the manufacturing process, high tensile strength and compressibility.² The Cartiva implant is a 6 to 10 mm PVA disc that is implanted with a slight protrusion to act as a spacer for the first MTP joint. It comes with dedicated reusable instrumentation, which includes a drill bit, introducer, and placer.

For focal cartilage defects of the knee, there are multiple treatment options with patient-specific treatment choice dependent on symptom severity, age, activity level as well as lesion size.³ Conservative medical treatment may be preferred in older individuals with mild symptoms and low-grade lesions; however, surgery is preferred for acute defects, larger lesions, active individuals, or when conservative treatment has failed. There are many surgical repair options. Scaffold-based techniques like Agili-CTM are a type of regenerative repair where the matrix provides a supportive framework for cell growth and tissue repair. Scaffolds are biodegradable and can be implanted in a one-step procedure.

Summary of Evidence

For individuals who have early-stage first MTP joint OA who receive a synthetic cartilage implant, the evidence is lacking. The relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. The pivotal study was performed in individuals with Coughlin stage 2, 3, or 4 hallux rigidus. No evidence was identified in individuals with stage 0 to early-stage 2 hallux rigidus. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.



For individuals who have advanced first MTP joint OA who receive a synthetic cartilage implant, the evidence includes a pivotal non-inferiority trial. The relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Arthrodesis is the established treatment for advanced arthritis of the great toe, although the lack of mobility can negatively impact sports and choice of footwear, and is not a preferred option of individuals. Implants have the potential to reduce pain and maintain mobility in the first MTP joint but have in the past been compromised by fragmentation, dislocation, particle wear, osteolysis, and loosening. A PVA hydrogel implant has shown properties similar to articular cartilage in vitro and was approved by the FDA in 2016 for the treatment of painful degenerative or posttraumatic arthritis in the MTP joint. Results at two years from the pivotal non-inferiority trial showed pain scores that were slightly worse compared to individuals treated with arthrodesis and similar outcomes between the groups for activities of daily living (ADL) and sports. In a non-inferiority trial, some benefit should be observed to justify the non-inferiority margin. However, the benefit of Cartiva with respect to increased range of motion does not appear to translate to improved ADL, sports activities, or individual report of well-being compared to arthrodesis. In addition, the Cartiva group showed a higher rate of adverse outcomes (Moderate Difficulty, Extreme Difficulty, and Unable to Do) compared to the arthrodesis group for walking for 15 min (16% vs 0%), Up Stairs (6% vs 0%) and Squats (19% vs 8%). Some bias in favor of the novel motion preserving implant was also possible, as suggested by the high dropout rate in the arthrodesis group after randomization. Five-year follow-up of both the randomized and run-in individuals who received an implant was reported in 2018 for 135 of 152 individuals. At this time point, 21% of implants had been removed with conversion to arthrodesis. Comparison to arthrodesis at long-term follow-up is needed to determine whether the implant improves function. Corroboration of long-term results in an independent study is also needed to determine the benefits and risks of the implant. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have cartilage and osteochondral defects of the knee who receive a cartilage repair implant, the evidence includes one randomized controlled trial (RCT). Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. There are currently multiple surgical treatment options for individuals with cartilage or osteochondral defects of the knee. The RCT compared the aragonite-based osteochondral implant (Agili-C) compared to either of two standard of care procedures - debridement or microfracture. The study is limited by lack of blinding and a heterogeneous comparator group. Additional, well-designed clinical trials are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.



Ongoing and Unpublished Clinical Trials

Some currently ongoing and unpublished trials that might influence this review are listed in [Table 1](#).

Table 1. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
Unpublished			
NCT03247439^a	A Prospective Study to Evaluate the Safety and Effectiveness of the Cartiva Synthetic Cartilage Implant for CMC in the Treatment of First Carpometacarpal Joint Osteoarthritis as Compared to LRTI Comparator (GRIP2)	74	Mar 2024 (last update Dec 2020)
NCT02391506^a	A Prospective Study to Evaluate the Safety and Effectiveness of the Cartiva Synthetic Cartilage Implant for CMC in the Treatment of First Carpometacarpal Joint Osteoarthritis	50	Mar 2019
NCT03935880	Treatment of Hallux Rigidus With Synthetic Hemiarthroplasty Versus Cheilectomy: A Randomized Controlled Trial	20 (actual)	Sept 2021 (terminated due to difficulty meeting recruitment goals)

NCT: national clinical trial.

^a Denotes industry-sponsored or cosponsored trial.

Practice Guidelines and Position Statements

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the policy conclusions.

Guidelines or position statements will be considered for inclusion if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence. Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

No guidelines or statements were identified.

American Society of Pain and Neuroscience

In 2022, the Consensus Guidelines on Interventional Therapies for Knee Pain (STEP Guidelines) were developed by a multidisciplinary panel of experts, including specialists in pain management, anesthesiology, orthopedic surgery, and rehabilitation medicine and published by the American Society of Pain and Neuroscience (ASPN).²⁷ Although the guidelines include recommendations on interventions such as marrow stimulation, autologous chondrocyte implantation, and osteochondral autograft transplantation, they do not include recommendations for that aragonite-based cartilage repair implant.

National Institute for Health and Care Excellence

The National Institute for Health and Care Excellence (NICE) published recommendations in 2024 for single-step scaffold insertion for repairing symptomatic chondral knee defects (HTG728).²⁶ The guidelines were not specific to a particular scaffold, but NICE recommended, "Single-step scaffold insertion can be used as an option for repairing symptomatic chondral knee defects with standard arrangements in place for clinical governance, consent and audit."

Medicare National Coverage

There is no national coverage determination.

Regulatory Status

The Cartiva implant was approved by the US Food and Drug Administration (FDA) in 2016 for the treatment of arthritis of the MTP joint. It has been distributed commercially since 2002 with approval in Europe, Canada, and Brazil. The Cartiva Synthetic Cartilage Implant (Wright Medical, Alpharetta, GA; now Stryker) was approved by the FDA through the premarket approval process (P150017) for painful degenerative or posttraumatic arthritis in the first MTP joint along with



hallux valgus or hallux limitus, hallux rigidus, and an unstable or painful MTP joint. Lesions greater than 10 mm in size and insufficient quality or quantity of bone are contraindications.

FDA product code: PNW.

In 2020, the FDA granted breakthrough status to Agili-CTM (CartiHeal, Ltd.), a proprietary cell-free biocompatible and biodegradable tapered-shape implant for the treatment of cartilage lesions in arthritic and non-arthritic joints that, when implanted into a pre-prepared osteochondral hole, acts as a 3-dimensional scaffold that potentially supports and promotes the regeneration of the articular cartilage and its underlying subchondral bone. Agili-C was FDA-approved in 2022 for the treatment of knee-joint surface lesions with a treatable area of 1 to 7 cm² without severe osteoarthritis.⁴ FDA product code: QRU.

References

1. Gould N, Schneider W, Ashikaga T. Epidemiological survey of foot problems in the continental United States: 1978-1979. *Foot Ankle*. Jul 1980; 1(1): 8-10. PMID 6115797
2. Baker MI, Walsh SP, Schwartz Z, et al. A review of polyvinyl alcohol and its uses in cartilage and orthopedic applications. *J Biomed Mater Res B Appl Biomater*. Jul 2012; 100(5): 1451-7. PMID 22514196
3. Bulgarelli A, Ruosi L, Conte P, et al. Matrices Used for Cartilage Regeneration. *Clin Sports Med*. Jul 2025; 44(3): 571-586. PMID 40514157
4. US FDA Premarket Approval. Agili-C - P210034. March 30, 2022.
<https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=P210034>. Accessed June 12, 2026.
5. Baumhauer JF, Singh D, Glazebrook M, et al. Prospective, Randomized, Multi-centered Clinical Trial Assessing Safety and Efficacy of a Synthetic Cartilage Implant Versus First Metatarsophalangeal Arthrodesis in Advanced Hallux Rigidus. *Foot Ankle Int*. May 2016; 37(5): 457-69. PMID 26922669
6. Smyth NA, Murawski CD, Hannon CP, et al. The Use of a Synthetic Cartilage Implant for Hallux Rigidus: A Systematic Review. *Foot Ankle Spec*. Aug 2021; 14(4): 366-371. PMID 32618201
7. Cassinelli SJ, Chen S, Charlton TP, et al. Early Outcomes and Complications of Synthetic Cartilage Implant for Treatment of Hallux Rigidus in the United States. *Foot Ankle Int*. Oct 2019; 40(10): 1140-1148. PMID 31195830
8. Butler JJ, Dhillon R, Wingo T, et al. Polyvinyl alcohol hydrogel implant for the treatment of hallux rigidus is associated with a high complication rate and moderate failure rate at short-term follow-up: a systematic review. *Eur J Orthop Surg Traumatol*. May 2024; 34(4): 1765-1778. PMID 38554163
9. U.S. Food and Drug Administration. Cartiva: Summary of Safety and Effectiveness. 2016;
https://www.accessdata.fda.gov/cdrh_docs/pdf15/p150017b.pdf. Accessed June 12, 2026.
10. Glazebrook M, Younger ASE, Daniels TR, et al. Treatment of first metatarsophalangeal joint arthritis using hemiarthroplasty with a synthetic cartilage implant or arthrodesis: A comparison of operative and recovery time. *Foot Ankle Surg*. Oct 2018; 24(5): 440-447. PMID 29409199



11. Goldberg A, Singh D, Glazebrook M, et al. Association Between Patient Factors and Outcome of Synthetic Cartilage Implant Hemiarthroplasty vs First Metatarsophalangeal Joint Arthrodesis in Advanced Hallux Rigidus. *Foot Ankle Int.* Nov 2017; 38(11): 1199-1206. PMID 28820949
12. Baumhauer JF, Singh D, Glazebrook M, et al. Correlation of Hallux Rigidus Grade With Motion, VAS Pain, Intraoperative Cartilage Loss, and Treatment Success for First MTP Joint Arthrodesis and Synthetic Cartilage Implant. *Foot Ankle Int.* Nov 2017; 38(11): 1175-1182. PMID 28992721
13. U.S. Food and Drug Administration. Cartiva: Post approval studies. Updated 2024. https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma_pas.cfm?c_id=4019&t_id=570803. Accessed June 12, 2026.
14. Glazebrook M, Blundell CM, O'Dowd D, et al. Midterm Outcomes of a Synthetic Cartilage Implant for the First Metatarsophalangeal Joint in Advanced Hallux Rigidus. *Foot Ankle Int.* Apr 2019; 40(4): 374-383. PMID 30501401
15. Budde K, Claassen L, Plaass C, et al. Synthetic cartilage implant vs. first metatarsophalangeal arthrodesis for the treatment of hallux rigidus. *Arch Orthop Trauma Surg.* Sep 2024; 144(9): 4343-4354. PMID 39287788
16. Joo PY, Baumhauer JF, Waldman O, et al. Physical Function and Pain Interference Levels of Hallux Rigidus Patients Before and After Synthetic Cartilage Implant vs Arthrodesis Surgery. *Foot Ankle Int.* Oct 2021; 42(10): 1277-1286. PMID 34024138
17. An TW, Cassinelli S, Charlton TP, et al. Radiographic and Magnetic Resonance Imaging of the Symptomatic Synthetic Cartilage Implant. *Foot Ankle Int.* Jan 2020; 41(1): 25-30. PMID 31538827
18. Shi E, Todd N, Rush S, et al. First Metatarsophalangeal Joint Space Area Decreases Within 1 Month After Implantation of a Polyvinyl Alcohol Hydrogel Implant: A Retrospective Radiographic Case Series. *J Foot Ankle Surg.* Nov 2019; 58(6): 1288-1292. PMID 31679683
19. Metikala S, Mahmoud K, O'Connor KM, et al. Adverse Events Related to Cartiva Hemiarthroplasty of First Metatarsal: An Analysis of Reports to the United States Food and Drug Administration. *Foot Ankle Spec.* Apr 2022; 15(2): 113-118. PMID 32723089
20. Glazebrook M, Baumhauer JF, Blundell C, et al. Letter Regarding: Early Outcomes and Complications of Synthetic Cartilage Implant for Treatment of Hallux Rigidus in the United States. *Foot Ankle Int.* Oct 2019; 40(10): 1149-1151. PMID 31600478
21. Thordarson DB, Cassinelli SJ, Charlton TP, et al. Response to "Letter Regarding: Early Outcomes and Complications of Synthetic Cartilage Implant for Treatment of Hallux Rigidus in the United States". *Foot Ankle Int.* Oct 2019; 40(10): 1152-1153. PMID 31600477
22. Altschuler N, Zaslav KR, Di Matteo B, et al. Aragonite-Based Scaffold Versus Microfracture and Debridement for the Treatment of Knee Chondral and Osteochondral Lesions: Results of a Multicenter Randomized Controlled Trial. *Am J Sports Med.* Mar 2023; 51(4): 957-967. PMID 36779614
23. Kon E, De Caro F, Dasa V, et al. Female patients report comparable results to males after the implantation of an aragonite-based scaffold for the treatment of knee chondral and osteochondral defects: a gender-based analysis of a RCT at 4 years' follow-up. *J Orthop Traumatol.* Mar 13 2025; 26(1): 17. PMID 40080289
24. Conte P, Anzillotti G, Crawford DC, et al. Differential analysis of the impact of lesions' location on clinical and radiological outcomes after the implantation of a novel aragonite-based scaffold to treat knee cartilage defects. *Int Orthop.* Dec 2024; 48(12): 3117-3126. PMID 39305313
25. Hunter CW, Deer TR, Jones MR, et al. Consensus Guidelines on Interventional Therapies for Knee Pain (STEP Guidelines) from the American Society of Pain and Neuroscience. *J Pain Res.* 2022; 15: 2683-2745. PMID 36132996
26. National Institute for Health and Care Excellence. Single-step scaffold insertion for repairing symptomatic chondral knee defects [HTG728]. <https://www.nice.org.uk/guidance/htg728/chapter/2-The-condition-current-treatments-and-procedure>. Accessed June 12, 2026.



Appendix

Appendix Table 1. Coughlin Clinical-Radiographic System for Grading Hallux Rigidus

Grade	Dorsiflexion	Radiographic Findings	Clinical Findings
0	40°-60° and/or 10%-20% loss vs normal side	Normal	No pain; only stiffness and loss of motion
1	30°-40° and/or 20%-50% loss vs normal side	Minimal changes	Mild or occasional pain and stiffness
2	10°-30° and/or 50%-75% loss vs normal side	Osteophytes, mild-to-moderate joint-space narrowing	Moderate-to-severe pain and stiffness that may be constant; pain occurs at maximum flexion
3	Greater than or equal to 10° and/or 75%-100% loss vs normal side	Osteophytes, substantial joint space narrowing	Nearly constant pain and substantial stiffness at extremes ROM, not at mid-range
4	Same as grade 3	Same as grade 3	Same as grade 3 but definite pain at mid-ROM

ROM: range of motion.

History

Date	Comments
05/01/19	New policy, approved April 9, 2019, effective August 2, 2019. Policy created with literature review through January 2019. Synthetic cartilage implants are considered investigational for the treatment of articular cartilage damage. Added codes L8641 and L8642.
10/01/19	Interim Review, approved September 5, 2019. Policy updated with literature review through July 2019; no references added. Policy statement unchanged.
10/01/20	Annual Review, approved September 1, 2020. Policy updated with literature review through June 2020; references added. Policy statement unchanged.
10/01/21	Annual Review, approved September 2, 2021. Policy updated with literature review through June 2, 2021; references added. Policy statement unchanged.
09/01/22	Annual Review, approved August 22, 2022. Policy updated with literature review through April 18, 2022; no references added. Policy statement unchanged.

Date	Comments
08/01/23	Minor update to Related Policies. Removed 7.01.569 and replaced with 7.01.48 Autologous Chondrocyte Implantation for Focal Articular Cartilage Lesions.
12/01/23	Annual Review, approved November 6, 2023. Policy updated with literature review through August 3, 2023; no references added. Policy statement unchanged. Changed the wording from "patient" to "individual" throughout the policy for standardization.
10/01/24	Annual Review, approved September 9, 2024. Policy updated with literature review through June 5, 2024; one reference added. Policy statement unchanged.
10/01/25	Annual Review, approved September 8, 2025. Policy updated with literature review through June 18, 2025; one reference added. Policy statement unchanged.
08/01/26	Annual Review, approved July 14, 2026. Effective for dates of service on or after November 6, 2026, following 90-day provider notification. Policy title changed from 7.01.160 Synthetic Cartilage Implants for Joint Pain to 7.01.160 Cartilage Implants for Joint Pain. Policy updated with literature review through March 5, 2026; references added and deleted. Policy statement updated to be inclusive of cartilage implants beyond synthetic cartilage. Removed CPT code 28291.

Disclaimer: This medical policy is a guide in evaluating the medical necessity of a particular service or treatment. The Company adopts policies after careful review of published peer-reviewed scientific literature, national guidelines and local standards of practice. Since medical technology is constantly changing, the Company reserves the right to review and update policies as appropriate. Member contracts differ in their benefits. Always consult the member benefit booklet or contact a member service representative to determine coverage for a specific medical service or supply. CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). ©2026 Premiera All Rights Reserved.

Scope: Medical policies are systematically developed guidelines that serve as a resource for Company staff when determining coverage for specific medical procedures, drugs or devices. Coverage for medical services is subject to the limits and conditions of the member benefit plan. Members and their providers should consult the member benefit booklet or contact a customer service representative to determine whether there are any benefit limitations applicable to this service or supply. This medical policy does not apply to Medicare Advantage.

