

PHARMACY POLICY – 5.01.545


Pharmacologic Treatment of Benign Prostatic Hyperplasia

Effective Date: Jun. 1, 2026
Last Revised: May 25, 2026
Replaces: N/A

RELATED MEDICAL POLICIES:
5.01.522 Advanced Therapies for Pharmacological Treatment of Pulmonary Arterial Hypertension

Select a hyperlink below to be directed to that section.

[POLICY CRITERIA](#) | [DOCUMENTATION REQUIREMENTS](#) | [CODING](#)
[RELATED INFORMATION](#) | [EVIDENCE REVIEW](#) | [REFERENCES](#) | [HISTORY](#)

 Clicking this icon returns you to the hyperlinks menu above.

Introduction

Benign prostatic hyperplasia (BPH) is the medical term for an enlarged prostate. BPH is a noncancerous condition caused by prostate cells that keep multiplying. Over time, the excess prostate tissue puts pressure on the tube that carries urine and semen out of the body. As more pressure is applied, urinary difficulties often develop. These can include a weak or slow stream, a sense that the bladder isn't completely empty, and/or needing to urinate frequently and urgently. This policy discusses the different types of treatment for BPH, and which medications need to be pre-approved by the health plan.

Note: The Introduction section is for your general knowledge and is not to be taken as policy coverage criteria. The rest of the policy uses specific words and concepts familiar to medical professionals. It is intended for providers. A provider can be a person, such as a doctor, nurse, psychologist, or dentist. A provider also can be a place where medical care is given, like a hospital, clinic, or lab. This policy informs them about when a service may be covered.

Policy Coverage Criteria

Documentation in the form of chart notes/medical records must be provided with prior authorization review for the agents described below.

Drug	Medical Necessity
Cialis (tadalafil)	<p>Cialis (tadalafil) may be considered medically necessary as second-line treatment for symptoms of benign prostatic hyperplasia (BPH) when ALL the following conditions are met:</p> <ul style="list-style-type: none"> • The individual has a diagnosis of benign prostatic hyperplasia (BPH) <p>AND</p> <ul style="list-style-type: none"> • Has tried and had an inadequate response or intolerance to a generic alpha blocker (e.g., terazosin, tamsulosin, doxazosin, or alfuzosin) <p>AND</p> <ul style="list-style-type: none"> • Has tried and had an inadequate response or intolerance to generic tadalafil <p>AND</p> <ul style="list-style-type: none"> • Dose is limited to 5 mg once daily <p>Note: Cialis (tadalafil), and generic tadalafil are also available in 10mg and 20mg tablets for use as needed. These are indicated ONLY for treatment of erectile dysfunction, not BPH.</p>
Generic tadalafil	<p>Generic tadalafil may be considered medically necessary as second-line treatment for symptoms of benign prostatic hyperplasia (BPH) when ALL the following conditions are met:</p> <ul style="list-style-type: none"> • The individual has a diagnosis of benign prostatic hyperplasia (BPH) <p>AND</p> <ul style="list-style-type: none"> • Has tried and had an inadequate response or intolerance to a generic alpha blocker (e.g., terazosin, tamsulosin, doxazosin, or alfuzosin) <p>AND</p> <ul style="list-style-type: none"> • Dose is limited to 5 mg once daily. <p>Note: Generic tadalafil are also available in 10mg and 20mg tablets for use as needed. These are indicated ONLY for treatment of erectile dysfunction, not BPH.</p>
Tezruly (terazosin oral solution)	<p>Tezruly (terazosin) may be considered medically necessary for the treatment of symptoms of benign prostatic hyperplasia (BPH) when ALL the following conditions are met:</p>



Drug	Medical Necessity
	<ul style="list-style-type: none"> • The individual has a diagnosis of benign prostatic hyperplasia (BPH) <p>AND</p> <ul style="list-style-type: none"> • Has tried and had an inadequate response or intolerance to ALL of the following: <ul style="list-style-type: none"> ○ One generic alpha blocker (e.g., terazosin, tamsulosin, doxazosin, or alfuzosin) ○ Generic finasteride, dutasteride, or silodosin <p>OR</p> <ul style="list-style-type: none"> • Documentation is provided that the oral solution is medically necessary (e.g., unable to swallow tablets or capsules) <p>AND</p> <ul style="list-style-type: none"> • Dose is limited to 10 mg once daily <p>Tezruly (terazosin) may be considered medically necessary for the treatment of hypertension when ALL the following conditions are met:</p> <ul style="list-style-type: none"> • The individual has a diagnosis of hypertension <p>AND</p> <ul style="list-style-type: none"> • Has tried and had an inadequate response or intolerance to ALL of the following: <ul style="list-style-type: none"> ○ Calcium channel blocker (e.g., amlodipine) ○ Angiotensin receptor blocker (e.g., losartan) OR angiotensin-converting enzyme inhibitor (e.g., lisinopril) ○ Diuretic (e.g., hydrochlorothiazide) <p>OR</p> <ul style="list-style-type: none"> • Documentation is provided that the oral solution is medically necessary (e.g., unable to swallow tablets or capsules) <p>AND</p> <ul style="list-style-type: none"> • Dose is limited to 20 mg once daily

Indication	Contract Exclusion
Erectile dysfunction in individuals without BPH	Use of Cialis (tadalafil) and generic tadalafil for erectile dysfunction in individuals without BPH is considered a contract exclusion.



Indication	Investigational
All other indications	<p>The medications listed in this policy are subject to the product's US Food and Drug Administration (FDA) dosage and administration prescribing information.</p> <p>Use of Cialis (tadalafil), generic tadalafil, and Tezruly (terazosin) for all other indications is considered investigational.</p>

Length of Approval	
Approval	Criteria
Initial authorization	All reviews for Cialis (tadalafil), generic tadalafil, and Tezruly (terazosin) may be approved up to 12 months.
Re-authorization criteria	All reviews for Cialis (tadalafil), generic tadalafil, and Tezruly (terazosin) may be approved up to 12 months as long as the drug-specific coverage criteria are met and chart notes demonstrate that the individual continues to show a positive clinical response to therapy.

Documentation Requirements
<p>The individual's medical records submitted for review for all conditions should document that medical necessity criteria are met. The record should include the following:</p> <ul style="list-style-type: none"> Office visit notes that contain the diagnosis, relevant history, physical evaluation, and medication history

Note: Tadalafil is available as Adcirca in different strengths for the treatment of pulmonary arterial hypertension (PAH). For these individuals, refer to the pulmonary arterial hypertension guidelines contained in a separate medical policy (see [Related Policies](#)).

Coding

N/A



Related Information

Benefit Application

This policy applies to all lines of business. Language pertaining specifically to contractual exclusion of tadalafil is applicable only to benefits that exclude coverage for treatment of erectile dysfunction.

This policy is managed through the pharmacy benefit.

Evidence Review

Description

Benign prostatic hyperplasia (BPH) is a condition characterized by epithelial and stromal cell proliferation (enlargement) of the prostate gland. BPH and BPH-related lower urinary tract symptoms (LUTS) are age-dependent, with BPH being one of the top 10 most commonly diagnosed conditions in men over 50 years of age. The estimated prevalence of LUTS secondary to BPH in Caucasian Americans between the ages of 50 and 79 in 2000 was 42%. In a European cross-sectional study of men >40 years of age attending urology clinics, the prevalence of LUTS secondary to BPH was 56% and erectile dysfunction was 40%. These conditions are also frequently comorbid in the same men. In BPH, as the prostate gland enlarges it may constrict the urethra and/or the bladder wall may thicken and become irritable resulting in lower urinary tract symptoms such as diminished urine stream, intermittency, straining, polyuria, nocturia, urinary urgency, and inability to empty the bladder.

The American Urological Association (AUA) recommends for the treatment of BPH: alpha adrenergic antagonists, 5-alpha reductase inhibitors, anticholinergics, and PDE5 inhibitors. However, the AUA does note the combination of low-dose daily 5mg tadalafil with alpha adrenergic antagonists for the treatment of LUTS/BPH as it offers no advantages in symptom improvement over either agent alone. The mechanism of action of alpha-adrenergic antagonists is blockade of alpha-1 receptors in the prostate, bladder, and urethra which relaxes smooth muscle in these tissues and improves urine flow and symptoms associated with BPH. It is important to note that drugs in this class do not reduce prostate size. 5-Alpha reductase



inhibitors interfere with testosterone's stimulatory effect on prostate enlargement and therefore do reduce prostate size.

Cialis (tadalafil)

The clinical efficacy of tadalafil 5 mg once daily for lower urinary tract symptoms (LUTS) suggestive of benign prostatic hyperplasia (BPH) was primarily established in 3 similarly designed multinational, 12-week, randomized, double-blind, placebo-controlled trials. Two of these studies were conducted in men with LUTS suggestive of BPH (Porst, 2011 and Roehrborn, 2008) and one was specific to men with comorbid BPH and erectile dysfunction (Egerdie, 2011). The primary efficacy endpoint in all 3 trials was change from baseline in total International Prostate Symptom Score (IPSS). The IPSS assesses severity of the spectrum of symptoms associated with BPH. The comorbid ED study had a co-primary endpoint that was change from baseline in the erectile function domain score of International Index of Erectile Function (IIEF-EF). All 3 trials consistently showed a statistically and clinically significant improvement in these surrogate outcomes after 12-weeks. While these studies were not specifically designed to evaluate the following endpoints, nocturia and objective urodynamic measures (peak urinary flow and postvoid residual volume) were consistently not improved.

A 1-year, open-label, safety extension of the registrational Roehrborn (2008) study suggests efficacy for LUTS secondary to BPH is maintained longer-term (Donatucci, 2011). Two other preliminary RCTs also consistently support the short-term (12-week) efficacy of tadalafil 5 mg once daily for LUTS secondary to BPH with or without comorbid ED.

Results from one unpublished, multinational, 12-week, randomized, double-blind, double dummy, placebo-controlled trial with a parallel tamsulosin 0.4 mg once daily arm were recently posted on [Clinicaltrials.gov](https://clinicaltrials.gov) (NCT00970632). While this study was not designed to compare the two active drugs, tadalafil 5 mg once daily appeared to provide similar improvements to that of tamsulosin 0.4 mg in men with LUTS suggestive of BPH. As might be expected, tadalafil appeared superior to the alpha-blocker for treatment of comorbid erectile dysfunction.

One prospective, open-label, uncontrolled study was identified on literature search that examined the efficacy and safety of add-on tadalafil at the approved 5 mg daily dose combined with an alpha-blocker (tamsulosin 0.2 mg or alfuzosin 10 mg) for 12-weeks in 158 Korean men with concurrent LUTS-BPH and ED. Total IPSS and IIEF-5 were significantly ($p < 0.001$) improved with combination therapy, but Qmax and PVR did not, compared to baseline low-dose alpha-blocker monotherapy.



The registrational RCTs show tadalafil 5 mg once daily to be well tolerated for the short-term treatment of LUTS suggestive of BPH with or without comorbid ED. Overall, adverse events were infrequent and the majority (>95%) were mild to moderate in severity. No unexpected adverse events occurred. No clinically significant changes in vitals or electrocardiogram were observed. Rates of myocardial infarction appeared similar to that of placebo in the same population, with the caveat that these trials were of short duration (12 weeks). In short-term (12-week) clinical trials, incidence of a positive orthostatic test was similar in tadalafil-treated and placebo-treated LUTS-BPH individuals, about 20%. No priapism was reported.

A long-term (1-year) safety extension in the target population found no new or greater frequency of any adverse events, serious adverse events, or discontinuation for adverse events compared to double-blind short-term (12-week) treatment (Donatucci, 2011).

Two open-label safety extension trials of tadalafil 5 mg once daily in men with ED also showed the drug to be well tolerated for up to 2 years. No unexpected adverse events were observed. No deaths or serious adverse events, cardiovascular or otherwise, were attributed to study drug.

No sudden hearing or vision losses (non-arteritic anterior ischemic optic neuropathy [NAION]) were reported in short- or long-term clinical trials for LUTS secondary to BPH.

In the unpublished, multinational, 12-week, randomized, double-blind, double dummy, placebo-controlled trial with the parallel tamsulosin 0.4 mg once daily arm (NCT00970632), incidence of adverse events, serious adverse events, and discontinuation for adverse events appeared similar between tadalafil 5 mg, tamsulosin 0.4 mg, and placebo once daily in men with LUTS-BPH with or without ED.

A prospective, multicenter, randomized, open-label, parallel-group study (Yoshida et al. 2017) compared the efficacy and safety of silodosin versus tadalafil in 192 individuals with LUTS-BPH. The primary efficacy endpoint was change in IPSS total symptom score before and after Period 1 treatment (Week 4). Both silodosin and tadalafil demonstrated statistically significant improvement from baseline (Week 0) IPSS total symptom score, with a mean \pm SD change of -10.1 ± 6.4 ($P < 0.0001$) and -8.0 ± 6.3 ($P < 0.0001$), respectively. The difference between silodosin and tadalafil was statistically significant ($P = 0.0277$). In addition, silodosin demonstrated a significantly greater decrease in symptoms of incomplete emptying, weak stream, and nocturia than tadalafil ($P = 0.0254$, $P = 0.0067$, and $P = 0.0387$). Adverse drug reactions occurred more frequently with silodosin (23.4%) than tadalafil (8.4%), but no serious adverse reactions were observed in both arms. Nervous system disorders (i.e., dizziness and headache) and gastrointestinal disorders (i.e., soft feces) were reported in both arms, but orthostatic hypotension ($n = 2$), nasal congestion ($n = 4$), ejaculation disorder ($n = 6$), and retrograde ejaculation ($n = 5$) were only reported in the silodosin arm.



2014 Update

Updated per literature search 7/1/13 to 10/31/14. No changes required.

2015 Update

Updated per literature search 11/1/14 to 9/15/15. There was no information which prompted a change in the policy statements. Rapaflo was added as a qualifier to coverage of Cialis along with Proscar and Avodart. Verbiage also updated to call them out separately from the generic alpha blockers.

2016 Update

A literature search was conducted between 04/01/15 and 12/06/16. No information was found which would prompt a change in the existing policy statements.

2017 Update

A literature search was conducted between 07/01/16 and 11/01/17. No information was found which would prompt a change in the existing policy statements.

2018 Update

A literature search was conducted between 11/01/17 and 10/31/18. No information was found which would prompt a change in the existing policy statements. Tadalafil was studied in a placebo-controlled phase 3 RCT in individuals 7-14 years of age with Duchenne Muscular Dystrophy. Tadalafil failed to show benefit in reducing rate of ambulatory decline.

2019 Update

Reviewed Cialis (tadalafil) prescribing information and conducted a literature search from 11/1/18 through 11/30/19. No new information was identified that would require changes to this policy. Added generic tadalafil name to policy.



2020 Update

Reviewed Cialis (tadalafil) prescribing information and conducted a literature search from 12/1/19 through 9/30/20. No new information was identified that would require changes to this policy. Updated criteria to reflect that Proscar (finasteride), Avodart, (dutasteride), and Rapaflo (silodosin) are now all available as generics and removed reference to the brand name medications from coverage criteria.

2021 Update

Reviewed Cialis (tadalafil) prescribing information and the 2021 on the Management of Benign Prostatic Hyperplasia/ Lower Urinary Tract Symptoms. No new information was identified that would require changes to this policy.

2022 Update

Reviewed Cialis (tadalafil) prescribing information and checked for updated American Urological Association (AUA) Guidelines. The 2021 AUA Guidelines remain the most recent version of guidelines. Added coverage criteria for Entadfi (finasteride and tadalafil) for the treatment of symptoms of BPH. Entadfi is a combination of finasteride, a 5 α -reductase inhibitor, and tadalafil, a phosphodiesterase 5 (PDE5) inhibitor. Entadfi is not recommended for more than 26 weeks because the incremental benefit of tadalafil decreases from 4 weeks until 26 weeks, and the incremental benefit beyond 26 weeks is unknown.

2023 Update

Reviewed Cialis (tadalafil) prescribing information and Entadfi (finasteride and tadalafil) prescribing information. No new information was identified that would require changes to this policy.



2024 Update

Reviewed prescribing information of drugs listed in the policy. Updated title from “Tadalafil Products for Benign Prostatic Hyperplasia” to “Pharmacologic Treatment of Benign Prostatic Hyperplasia”. Added coverage criteria for Avodart (dutasteride), Chewtadzy (tadalafil), Flomax (tamsulosin), and Tezruly (terazosin). Clarified that the quantity limit for Cialis (tadalafil) and generic tadalafil is 5 mg once daily. Updated Cialis (tadalafil) step therapy requirements from tried and had an inadequate response or intolerance to a generic alpha blocker and generic finasteride, dutasteride or silodosin to tried and had an inadequate response or intolerance to a generic alpha blocker and generic tadalafil. Updated generic tadalafil step therapy requirements from tried and had an inadequate response or intolerance to a generic alpha blocker and generic finasteride, dutasteride or silodosin to tried and had an inadequate response or intolerance to a generic alpha blocker.

2025 Update

Reviewed prescribing information of drugs listed in the policy. Clarified that non-formulary exception review authorizations for all drugs listed in this policy may be approved up to 12 months. Clarified that the medications listed in this policy are subject to the product's FDA dosage and administration prescribing information.

2026 Update

Reviewed prescribing information of drugs listed in the policy. Removed coverage criteria for Chewtadzy (tadalafil) as product was discontinued. Removed coverage criteria for Avodart (dutasteride) and Cardura XL (doxazosin extended-release). Removed reference to non-formulary exception reviews.

References

1. Porst H, McVary KT, Montorsi F, et al. Effects of once-daily tadalafil on erectile function in men with erectile dysfunction and signs and symptoms of benign prostatic hyperplasia. *Eur Urol* 2009;56(4):727-735.
2. McVary KT, Roehrborn CG, Kaminetsky JC, et al. Tadalafil relieves lower urinary tract symptoms secondary to benign prostatic hyperplasia. *J Urol* 2007;177:1401-1407.



3. Lee JY, Park SY, Jeong TY, et al. Combined tadalafil and α -blocker therapy for benign prostatic hyperplasia in patients with erectile dysfunction: a multicenter prospective study. *J Andrology* 2012 Aug 25
4. Porst H, Rajfer J, Casabe A, et al. Long-term safety and efficacy of tadalafil 5 mg dosed once daily in men with erectile dysfunction. *J Sex Med* 2008;5:2160-2169.
5. Porst H, Kim ED, Casabe AR, et al. Efficacy and safety of tadalafil once daily in the treatment of men with lower urinary tract symptoms suggestive of benign prostatic hyperplasia: results of an international randomized, double-blind, placebo-controlled trial. *Eur Urol* 2011;60(5):1105-1113.
6. Roehrborn CG, McVary KT, Elion-Mboussa A, et al. Tadalafil administered once daily for lower urinary tract symptoms secondary to benign prostatic hyperplasia: a dose finding study. *J Urol* 2008;180:1228-1234.
7. Donatucci CF, Brock GB, Goldfischer ER, et al. Tadalafil administered once daily for lower urinary tract symptoms secondary to benign prostatic hyperplasia: a 1-year, open-label extension study. *BJUI* 2011;107:1110-1116.
8. Roehrborn CG, Kaminetsky JC, Auerbach SM, et al. Changes in peak urinary flow and voiding efficiency in men with signs and symptoms of benign prostatic hyperplasia during once daily tadalafil administration. *BJUI* 2009;105:502-507.
9. Broderick GA, Brock GB, Roehrborn CG, et al. Effects of tadalafil on lower urinary tract symptoms secondary to benign prostatic hyperplasia in men with or without erectile dysfunction. *Urology* 2010;75:1452-1459.
10. Egerdie RB, Auerbach S, Roehrborn CG, et al. Tadalafil 2.5 or 5 mg administered once daily for 12 weeks in men with both erectile dysfunction and signs and symptoms of benign prostatic hyperplasia: results of a randomized, placebo-controlled, double-blind study. *J Sex Med* 2011 Oct 7
11. National Kidney and Urological Diseases Information Clearinghouse. Prostate enlargement: benign prostatic hyperplasia. Available at: <https://www.niddk.nih.gov/health-information/urologic-diseases/prostate-problems/prostate-enlargement-benign-prostatic-hyperplasia> Accessed April 13, 2026.
12. National Kidney and Urological Diseases Information Clearinghouse. Kidney and urologic diseases statistics for the United States. Available at: <https://www.niddk.nih.gov/health-information/health-statistics> Accessed April 13, 2026.
13. AUA Practice Guidelines Committee. AUA guideline on management of benign prostatic hyperplasia/ lower urinary tract symptoms (updated 2021; amended 2023). Available at [https://www.auanet.org/guidelines/guidelines/benign-prostatic-hyperplasia-\(bph\)-guideline](https://www.auanet.org/guidelines/guidelines/benign-prostatic-hyperplasia-(bph)-guideline) Accessed April 13, 2026.
14. McConnell JD, Roehrborn CG, Bautista OM, et al. The long-term effect of doxazosin, finasteride, and combination therapy on the clinical progression of benign prostatic hyperplasia. *N Engl J Med*. 2003;349:2387-2398.
15. Yoshida M, Origasa H. and Seki N. Comparison of silodosin versus tadalafil in patients with lower urinary tract symptoms associated with benign prostatic hyperplasia. *LUTS*. 2017; 9: 176–186.
16. Víctor RG, Sweeney HL, Finkel R, et al. A phase 3 randomized placebo-controlled trial of tadalafil for Duchenne muscular dystrophy. *Neurology*. 2017 Oct 24;89(17):1811-1820. doi: 10.1212/WNL.0000000000004570. Epub 2017 Sep 29.
17. Tezruly (terazosin). Prescribing Information. ANI Pharmaceuticals, Inc. Baudette, MN. Revised July 2024.
18. Cialis (tadalafil). Prescribing information. Eli Lilly and Company; Indianapolis, IN. Revised March 2026.

History

Date	Comments
08/12/13	New policy. Add to Prescription Drug section.



Date	Comments
12/17/14	Annual Review. Policy updated with literature review. No change in policy statements.
10/13/15	Annual Review. Added Rapaflo as a qualifier to coverage of Cialis along with Proscar and Avodart. Verbiage also updated to call them out separately from the generic alpha blockers. Policy statements unchanged.
01/01/17	Annual Review, approved December 13, 2016. No changes to Policy Criteria.
12/01/17	Annual Review, approved November 21, 2017. No changes to the Policy Criteria. One study was added to the Reference List.
07/01/18	Interim Review, approved June 5, 2018. Added requirement of documentation in the form of chart notes/medical records for medical necessity review of medications within this policy.
12/01/18	Annual Review, approved November 21, 2018. No changes. Added reference to a failed trial in patients with DMD.
01/01/20	Annual Review, approved December 10, 2019. Added generic tadalafil to policy. No changes to policy statement.
12/01/20	Annual Review, approved November 3, 2020. No changes to policy statement. Updated criteria to reflect that Proscar (finasteride), Avodart, (dutasteride), and Rapaflo (silodosin) are now all available as generics and removed reference to the brand name medications from coverage criteria.
10/01/21	Annual Review, approved September 23, 2021. No changes to policy statement.
05/01/22	Annual Review, approved April 12, 2022. Added coverage criteria for Entadfi (finasteride and tadalafil) for the treatment of symptoms of BPH.
05/01/22	Interim Review, approved April 25, 2022. Updated title from "Cialis (tadalafil) for Benign Prostatic Hyperplasia" to "Tadalafil Products for Benign Prostatic Hyperplasia".
06/01/23	Annual Review, approved May 22, 2023. Updated "patient" to "individual" for the process of standardization. Reviewed Cialis (tadalafil) prescribing information and Entadfi (finasteride and tadalafil) prescribing information. No new information was identified that would require changes to this policy.
08/01/24	Annual Review, approved July 8, 2024. No changes to policy statements.
12/01/24	Interim Review, approved November 12, 2024. Updated title from "Tadalafil Products for Benign Prostatic Hyperplasia" to "Pharmacologic Treatment of Benign Prostatic Hyperplasia". Added coverage criteria for Avodart (dutasteride), Chewtadzy (tadalafil), Flomax (tamsulosin), and Tezruly (terazosin). Clarified that the quantity limit for Cialis (tadalafil) and generic tadalafil is 5 mg once daily. Updated Cialis (tadalafil) step therapy requirements from tried and had an inadequate response or intolerance to a generic alpha blocker and generic finasteride, dutasteride or silodosin to tried and had an inadequate response or intolerance to a generic alpha blocker and generic tadalafil. Updated generic tadalafil step therapy requirements from tried and had an inadequate response or intolerance to a generic alpha blocker and generic finasteride, dutasteride



Date	Comments
	or silodosin to tried and had an inadequate response or intolerance to a generic alpha blocker.
03/01/25	Annual Review, approved February 24, 2025. Clarified that non-formulary exception review authorizations for all drugs listed in this policy may be approved up to 12 months. Clarified that the medications listed in this policy are subject to the product's FDA dosage and administration prescribing information.
07/01/25	Interim Review, approved June 10, 2025. Added coverage criteria for Cardura (doxazosin) for the treatment of BPH and hypertension. Added coverage criteria for Cardura XL (doxazosin extended-release) for the treatment of BPH.
01/01/26	Interim Review, approved December 22, 2025. Removed coverage criteria for Cardura (doxazosin), Entadfi (finasteride and tadalafil), Flomax (tamsulosin).
06/01/26	Annual Review, approved May 25, 2026. Removed coverage criteria for Chewtadzy (tadalafil) as product was discontinued. Removed coverage criteria for Avodart (dutasteride) and Cardura XL (doxazosin extended-release). Removed reference to non-formulary exception reviews.

Disclaimer: This medical policy is a guide in evaluating the medical necessity of a particular service or treatment. The Company adopts policies after careful review of published peer-reviewed scientific literature, national guidelines and local standards of practice. Since medical technology is constantly changing, the Company reserves the right to review and update policies as appropriate. Member contracts differ in their benefits. Always consult the member benefit booklet or contact a member service representative to determine coverage for a specific medical service or supply. CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). ©2026 Premera All Rights Reserved.

Scope: Medical policies are systematically developed guidelines that serve as a resource for Company staff when determining coverage for specific medical procedures, drugs or devices. Coverage for medical services is subject to the limits and conditions of the member benefit plan. Members and their providers should consult the member benefit booklet or contact a customer service representative to determine whether there are any benefit limitations applicable to this service or supply. This medical policy does not apply to Medicare Advantage.

