

## MEDICAL POLICY – 2.01.107

# Fractional Carbon Dioxide (CO2) Laser Ablation Treatment of Hypertrophic Scars or Keloids for Functional Improvement

BCBSA Ref. Policy: 2.01.107

Effective Date: April 1, 2024

Last Revised: Mar. 12, 2024


Replaces: N/A

RELATED MEDICAL POLICIES:

2.01.57 Electrostimulation and Electromagnetic Therapy for Treating Wounds

Select a hyperlink below to be directed to that section.

[POLICY CRITERIA](#) | [DOCUMENTATION REQUIREMENTS](#) | [CODING](#)  
[RELATED INFORMATION](#) | [EVIDENCE REVIEW](#) | [REFERENCES](#) | [HISTORY](#)

 Clicking this icon returns you to the hyperlinks menu above.

## Introduction

Fractional carbon dioxide laser ablation is a minimally invasive procedure used to treat skin conditions such as hypertrophic scars or keloids. This treatment involves using a specialized laser to remove layers of scar tissue, promoting the growth of new, healthier skin cells. By targeting specific areas of the scar, the laser helps to improve the appearance and flexibility of the scar tissue, leading to functional improvement.

**Note:** The Introduction section is for your general knowledge and is not to be taken as policy coverage criteria. The rest of the policy uses specific words and concepts familiar to medical professionals. It is intended for providers. A provider can be a person, such as a doctor, nurse, psychologist, or dentist. A provider also can be a place where medical care is given, like a hospital, clinic, or lab. This policy informs them about when a service may be covered.

## Policy Coverage Criteria

Treatment	Investigational
<b>Carbon dioxide (CO2) fractional laser ablation treatment of hypertrophic scars or keloids</b>	<b>Carbon dioxide (CO2) fractional laser ablation treatment of hypertrophic scars or keloids for functional improvement is considered investigational.</b>

## Coding

Code	Description
<b>CPT</b>	
0479T	Fractional ablative laser fenestration of burn and traumatic scars for functional improvement; first 100 cm2 or part thereof, or 1% of body surface area of infants and children
0480T	Fractional ablative laser fenestration of burn and traumatic scars for functional improvement; each additional 100 cm2, or each additional 1% of body surface area of infants and children, or part thereof

**Note:** CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). HCPCS codes, descriptions and materials are copyrighted by Centers for Medicare Services (CMS).

## Related Information

N/A

## Evidence Review

### Description

Hypertrophic scars and keloids are cutaneous lesions resulting from abnormal wound healing. There is no gold standard therapy for hypertrophic scars and keloids, and treatment frequently involves multiple techniques including pharmacotherapy, compression, surgery, radiation, and



light sources. For scars and keloids impairing function, fractional carbon dioxide (CO<sub>2</sub>) ablative laser treatment is proposed to improve abnormal texture, thickness, and stiffness of scars by ablative destruction and resurfacing. The treatment may be used as monotherapy or in combination with other therapies (e.g., sequential treatment with other lasers, sequential treatment with other therapies, or laser-assisted drug delivery).

## Background

### *Hypertrophic Scars and Keloids*

Hypertrophic scars and keloids are cutaneous lesions resulting from abnormal wound healing. Hypertrophic scars present as raised lesions that do not exceed the limits of the original skin injury. They tend to regress spontaneously within one year.<sup>1</sup> Keloids present as raised, firm lesions that extend beyond the margins of original injury. Keloids do not regress spontaneously, are often refractory to treatment, and have a high probability of recurrence after excision. The highest prevalence of keloids is in people of color, with an incidence of up to 16% in Black Africans.<sup>2</sup> Keloids can occur months or years after injury.<sup>3</sup>

Consensus-based clinical recommendations published in 2014 endorsed the use of a scar classification system first developed in 2002.<sup>4</sup> In this system, hypertrophic scars are classified as linear (e.g., surgical, traumatic) or widespread (e.g., burn). Keloids are classified as minor or major. Minor keloids are focally raised, itchy scars extending over normal tissue. Major keloids are large, raised (>0.5 cm) scars, possibly painful or pruritic, and extending over normal tissue. Major keloids are often refractory to treatment and have a high probability of recurrence after excision. Mature scars are light-colored and flat. Immature scars are slightly elevated in the process of remodeling and may be painful or itchy. Immature hypertrophic scars (red, slightly raised) may develop into hypertrophic scars; if they persist for longer than 1 month, the guidelines recommend treating them as a linear hypertrophic scar.

There is no gold standard therapy for hypertrophic scars and keloids, and treatment frequently involves multiple techniques including pharmacotherapy, compression, surgery, radiation, and light sources.<sup>5</sup>

### *Laser Therapy for Scar Treatment*

Carbon dioxide (CO<sub>2</sub>) fractional laser treatment was initially developed for cosmetic purposes (e.g., photoaging, acne scarring). Fractional CO<sub>2</sub> laser ablation works by creating microscopic



thermal wounds, resulting in tissue vaporization and coagulation of surrounding extracellular proteins. The technique has the advantage of reaching the dermis by ablating the epidermis, while avoiding complications associated with nonfractional ablative lasers (no longer in use), such as postoperative pain and infection. For scars and keloids impairing function, CO<sub>2</sub> fractional ablative laser treatment is proposed to improve abnormal texture, thickness, and stiffness of scars by ablative destruction and resurfacing. The treatment may be used as monotherapy or in combination with other therapies (e.g., sequential treatment with other lasers, sequential treatment with other therapies, or laser-assisted drug delivery).

This review focuses on CO<sub>2</sub> fractional ablative laser treatment for functional improvement. Other types of lasers used for hypertrophic scars and keloids include pulsed dye laser and intense pulse light.

## Summary of Evidence

For individuals with hypertrophic scars who receive fractional CO<sub>2</sub> ablative laser treatment as monotherapy for functional improvement, the evidence includes randomized controlled trials (RCTs), nonrandomized studies, and systematic reviews of these studies. Relevant outcomes are functional improvement, quality of life, and adverse effects of treatment. A Cochrane systematic review included 3 RCTs of CO<sub>2</sub> fractional therapy as monotherapy compared to no treatment. None evaluated functional outcomes. For all outcomes reported, the review authors graded the overall evidence as very low certainty, downgraded for very serious imprecision and serious risk of bias. The reviewers concluded that it was unclear whether fractional CO<sub>2</sub> laser impacts scar severity compared with no treatment as measured by commonly used scar scales. Conclusions were limited by study heterogeneity and lack of functional outcome measures. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with keloids who receive fractional CO<sub>2</sub> ablative laser treatment as monotherapy for functional improvement, the evidence includes RCTs, nonrandomized studies, and systematic reviews of these studies. Relevant outcomes are functional improvement, quality of life, and adverse effects of treatment. One RCT included in a Cochrane review evaluated CO<sub>2</sub> fractional laser therapy monotherapy for keloids compared to no treatment. The review authors concluded that it is uncertain whether fractional CO<sub>2</sub> impacts on keloid scar severity compared to no treatment after up to 6 months, downgrading the evidence for very serious imprecision and serious risk of bias. Adverse events and function were not assessed. Scar pain and pruritus outcomes were not presented by the treatment arm. Another systematic review included 1 RCT of CO<sub>2</sub> fractional laser monotherapy compared to intralesional triamcinolone and found no



significant differences between keloid response but faster improvement in the intralesional triamcinolone group. Functional outcomes were not evaluated. Conclusions were limited by study heterogeneity and lack of functional outcome measures. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with hypertrophic scars who receive fractional CO2 ablative laser treatment as adjunctive therapy for functional improvement, the evidence includes an RCT, nonrandomized studies, and systematic reviews of these studies. Relevant outcomes are functional improvement, quality of life, and adverse effects of treatment. A systematic review included a 3-arm RCT that compared combination therapy with CO2 laser plus IPL laser, CO2 monotherapy, or no therapy in 23 individuals with hypertrophic scars. Statistically significant improvements were found on commonly used scar scales for both CO2 plus IPL laser and for CO2 alone. The reviewers determined the trial was at unclear risk of bias for unclear adequacy of allocation concealment and blinding. Functional outcomes were not evaluated, and adverse events were not reported. Conclusions were limited by study heterogeneity and lack of functional outcome measures. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with keloids who receive fractional CO2 ablative laser treatment as adjunctive therapy for functional improvement, the evidence includes an RCT, nonrandomized studies, and systematic reviews of these studies. Relevant outcomes are functional improvement, quality of life, and adverse effects of treatment. One RCT included in 2 systematic reviews compared CO2 laser plus intralesional triamcinolone to cryosurgery plus triamcinolone. Of 60 individuals enrolled, 23 were lost to follow-up and not assessed. Scar severity ratings favored the laser therapy group at 12 months, but certainty of the evidence was downgraded due to very serious imprecision and serious risk of bias. Pain not related to treatment favored the CO2 group, but there was no difference in pruritus score. There were more frequent early adverse effects in the CO2 laser group. At 12 months, there was a recurrence of 6 keloid scars (16.7%), all of which were in the CO2 laser group. Conclusions were limited by heterogeneity of subject characteristics and study outcomes measures, small sample sizes, and inconsistent study designs. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

## Ongoing and Unpublished Clinical Trials

Some currently ongoing trials that might influence this review are listed in [Table 1](#).



**Table 1. Summary of Key Trials**

<b>NCT No.</b>	<b>Trial Name</b>	<b>Planned Enrollment</b>	<b>Completion Date</b>
<b>Ongoing</b>			
<b>NCT04736251</b>	A Prospective Intra-patient Single-blinded Randomised Trial to Examine the Mechanistic Basis of fractional Ablative carbon Dioxide Laser Therapy in Treating Adult Burns and/or Trauma Patients With Hypertrophic Scarring (SMOOTH)	60	Aug 2023
<b>NCT04567537</b>	Ablative Fractional Laser Treatment for the Improvement of Hypertrophic Scars and Scleroderma: a Prospective Cohort Study	20	May 2024
<b>NCT03692273</b>	A Within-Scar, Randomized Control Trial Comparing Fractional Ablative Carbon Dioxide Laser to Non-Energy-Based, Mechanical Tissue Extraction and No Treatment	120	Dec 2024
<b>NCT04364217</b>	Evaluating the Mechanism of Pain and Itch Reduction in Burn Scars Following Fractional Ablative CO2 Laser Treatment	28	Jul 2025

NCT: national clinical trial.

## **Practice Guidelines and Position Statements**

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the policy conclusions.

Guidelines or position statements will be considered for inclusion if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

### ***International Advisory Panel on Scar Management***

In 2014, Gold et al published updated international clinical recommendations on scar management.<sup>4</sup> Although they were not informed by a systematic review and strength of evidence ratings were not provided, the recommendations are frequently cited and were accompanied by a narrative review of the literature.<sup>5</sup> The recommendation document notes that,



where clinical evidence was lacking, management recommendations were based on advisory panel member consensus.

Specific recommendations on laser therapy include the following, according to scar classification:

- Immature or Erythematous Hypertrophic Scars

"In the case of persistent erythema for more than a month despite preventative efforts, management should transition to that of a linear hypertrophic scar or alternatively, pulsed-dye laser therapy may be applied once monthly for 2 to 3 months. If the scar is unresponsive to the pulsed-dye laser, fractional laser therapy or treatment as a linear hypertrophic scar may be instituted."

- Linear Hypertrophic Scars Arising from Surgery or Trauma

"Pulsed-dye or fractional laser therapy are second-line and, often, first -line options for linear hypertrophic scars."

- Widespread Burn Hypertrophic Scars

"Positive data for fractional lasers support their use for burn scar treatment. Ablative fractional lasers offer the advantage of fewer treatment sessions compared with nonablative options."

- Minor Keloids

If improvement with conservative therapy is not observed within 8 to 12 weeks, 5-FU in combination with intralesional corticosteroids and, ultimately, laser therapy or surgical excision may be considered. Although data from published clinical trials are lacking, some advisory panel members suggest the use of ablative fractional lasers over other types of laser therapy for the treatment of refractory keloids."

- Major Keloids

"Secondary management options for refractory keloids include laser treatment and surgical excision with appropriate prophylactic therapy."



## *Consensus Recommendations*

In 2020, Seago et al published consensus recommendations on laser treatment of scars and contractures.<sup>6</sup> The recommendations were developed by a panel of 26 dermatologists and plastic and reconstructive surgeons from 13 countries between March 2018 and March 2019. The panel used a modified Delphi method consisting of 2 rounds of email questionnaires and supplementary face-to-face meetings. The threshold for consensus recommendations was at least 80% concurrence among the panel members. The recommendations were not informed by a systematic review and do not include strength of evidence ratings.

Specific recommendation statements on laser therapy include the following:

- "The panel members are unanimous in their view that lasers are a first-line therapy in the management of traumatic scars and contractures."
- "The potential indications for laser treatment are determined based on clinical findings (i.e., erythema, hypopigmentation, hyperpigmentation, atrophy, hypertrophy, degree of epithelialization, pliability, and restriction) as well as subjective symptoms including pain and pruritus."
- "The fractional lasers, especially AFL, have the most potential to treat the entire range of clinical issues as a single modality. The optimal treatment routinely includes multiple laser types in concurrent or alternating treatment sessions to suit varying clinical presentations and treatment goals in a particular location at a particular time. Effective comprehensive traumatic scar management frequently incorporates surgical evaluation, ongoing conservative measures (i.e., compression, massage, and silicone gels and sheets); physical and occupational therapy; medical management such as corticosteroids and antimetabolites; and mental health evaluation where appropriate."

The document also includes recommendations on device application and settings but notes, "Optimal wavelengths and settings for traumatic scar management have not yet been fully elaborated in the literature and settings will vary depending on the characteristics of the particular device chosen by the operator, the clinical findings on the day of the visit (e.g., degree of erythema, presence of a tan, etc.), and issues specific to the individual (e.g., pain tolerance, approximate downtime, etc.)."

In its recommendations on scar assessment, the panel noted, "Continuing research is vital to determining the optimal laser devices, timing, combinations, and settings in the management of traumatic scars," and "Given the greater range of scar response to current laser techniques such as AFL, future scar assessment should incorporate evaluation of function, symptom relief, and overall quality of life to a greater extent."





## Medicare National Coverage

There is no national coverage determination.

## Regulatory Status

Multiple fractional CO2 laser systems have been approved by FDA through the 510(k) program. These devices have broad indications for dermatological procedures requiring ablation, resurfacing, and coagulation of soft tissue.

FDA Product Codes GEX, ONG.

## References

1. Leszczynski R, da Silva CA, Pinto ACPN, et al. Laser therapy for treating hypertrophic and keloid scars. *Cochrane Database Syst Rev*. Sep 26 2022; 9(9): CD011642. PMID 36161591
2. Walsh LA, Wu E, Pontes D, et al. Keloid treatments: an evidence-based systematic review of recent advances. *Syst Rev*. Mar 14 2023; 12(1): 42. PMID 36918908
3. UpToDate. Laser Therapy for Hypertrophic Scars and Keloids. 2023. <https://www.uptodate.com/contents/search>. (requires sign in access) Accessed February 12, 2024.
4. Gold MH, McGuire M, Mustoe TA, et al. Updated international clinical recommendations on scar management: part 2-- algorithms for scar prevention and treatment. *Dermatol Surg*. Aug 2014; 40(8): 825-31. PMID 25068544
5. Gold MH, Berman B, Clementoni MT, et al. Updated international clinical recommendations on scar management: part 1-- evaluating the evidence. *Dermatol Surg*. Aug 2014; 40(8): 817-24. PMID 25068543
6. Seago M, Shumaker PR, Spring LK, et al. Laser Treatment of Traumatic Scars and Contractures: 2020 International Consensus Recommendations. *Lasers Surg Med*. Feb 2020; 52(2): 96-116. PMID 31820478
7. Buhalog B, Moustafa F, Arkin L, et al. Ablative fractional laser treatment of hypertrophic burn and traumatic scars: a systematic review of the literature. *Arch Dermatol Res*. Jul 2021; 313(5): 301-317. PMID 32926192
8. Annabathula A, Sekar CS, Srinivas CR. Fractional Carbon Dioxide, Long Pulse Nd:YAG and Pulsed Dye Laser in the Management of Keloids. *J Cutan Aesthet Surg*. 2017; 10(2): 76-80. PMID 28852292
9. Azzam OA, Bassiouny DA, El-Hawary MS, et al. Treatment of hypertrophic scars and keloids by fractional carbon dioxide laser: a clinical, histological, and immunohistochemical study. *Lasers Med Sci*. Jan 2016; 31(1): 9-18. PMID 26498451
10. Behera B, Kumari R, Thappa DM, et al. Therapeutic Efficacy of Intralesional Steroid With Carbon Dioxide Laser Versus With Cryotherapy in Treatment of Keloids: A Randomized Controlled Trial. *Dermatol Surg*. Oct 2016; 42(10): 1188-98. PMID 27661432
11. Blome-Eberwein S, Gogal C, Weiss MJ, et al. Prospective Evaluation of Fractional CO2 Laser Treatment of Mature Burn Scars. *J Burn Care Res*. 2016; 37(6): 379-387. PMID 27828835



12. Daoud AA, Gianatasio C, Rudnick A, et al. Efficacy of Combined Intense Pulsed Light (IPL) With Fractional CO<sub>2</sub>-Laser Ablation in the Treatment of Large Hypertrophic Scars: A Prospective, Randomized Control Trial. *Lasers Surg Med.* Oct 2019; 51(8): 678-685. PMID 31090087
13. Garg GA, Sao PP, Khopkar US. Effect of carbon dioxide laser ablation followed by intralesional steroids on keloids. *J Cutan Aesthet Surg.* Jan 2011; 4(1): 2-6. PMID 21572673
14. Maari C (2017). Randomized, controlled, within-patient, single-blinded pilot study to evaluate the efficacy of the ablatie fractional CO<sub>2</sub> laser in the treatment of hypertrophic scars in adult burn patients. *J Am Acad Dermatol* 76:AB212.
15. Majid I, Imran S. Fractional Carbon Dioxide Laser Resurfacing in Combination With Potent Topical Corticosteroids for Hypertrophic Burn Scars in the Pediatric Age Group: An Open Label Study. *Dermatol Surg.* Aug 2018; 44(8): 1102-1108. PMID 30045141
16. Makboul M, Makboul R, Abdelhafez AH, et al. Evaluation of the effect of fractional CO<sub>2</sub> laser on histopathological picture and TGF-β<sub>1</sub> expression in hypertrophic scar. *J Cosmet Dermatol.* Sep 2014; 13(3): 169-79. PMID 25196683
17. Srivastava S, Kumari H, Singh A. Comparison of Fractional CO<sub>2</sub> Laser, Verapamil, and Triamcinolone for the Treatment of Keloid. *Adv Wound Care (New Rochelle).* Jan 01 2019; 8(1): 7-13. PMID 30705785
18. Wang J, Wu J, Xu M, et al. Combination therapy of refractory keloid with ultrapulse fractional carbon dioxide (CO<sub>2</sub>) laser and topical triamcinolone in Asians-long-term prevention of keloid recurrence. *Dermatol Ther.* Nov 2020; 33(6): e14359. PMID 33002270
19. Zuccaro J, Muser I, Singh M, et al. Laser Therapy for Pediatric Burn Scars: Focusing on a Combined Treatment Approach. *J Burn Care Res.* Apr 20 2018; 39(3): 457-462. PMID 29897540

## History

Date	Comments
04/01/24	New policy, approved March 12, 2024, Policy created with literature review through November 30, 2023. CO <sub>2</sub> fractional laser ablation treatment for hypertrophic scars and keloids to improve function is considered investigational. Added CPT codes 0479T and 0480T.

**Disclaimer:** This medical policy is a guide in evaluating the medical necessity of a particular service or treatment. The Company adopts policies after careful review of published peer-reviewed scientific literature, national guidelines and local standards of practice. Since medical technology is constantly changing, the Company reserves the right to review and update policies as appropriate. Member contracts differ in their benefits. Always consult the member benefit booklet or contact a member service representative to determine coverage for a specific medical service or supply. CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). ©2024 Premera All Rights Reserved.

**Scope:** Medical policies are systematically developed guidelines that serve as a resource for Company staff when determining coverage for specific medical procedures, drugs or devices. Coverage for medical services is subject to the limits and conditions of the member benefit plan. Members and their providers should consult the member



benefit booklet or contact a customer service representative to determine whether there are any benefit limitations applicable to this service or supply. This medical policy does not apply to Medicare Advantage.



## Discrimination is Against the Law

Premera Blue Cross (Premera) complies with applicable Federal and Washington state civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, sex, gender identity, or sexual orientation. Premera does not exclude people or treat them differently because of race, color, national origin, age, disability, sex, gender identity, or sexual orientation. Premera provides free aids and services to people with disabilities to communicate effectively with us, such as qualified sign language interpreters and written information in other formats (large print, audio, accessible electronic formats, other formats). Premera provides free language services to people whose primary language is not English, such as qualified interpreters and information written in other languages. If you need these services, contact the Civil Rights Coordinator. If you believe that Premera has failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, sex, gender identity, or sexual orientation, you can file a grievance with: Civil Rights Coordinator — Complaints and Appeals, PO Box 91102, Seattle, WA 98111, Toll free: 855-332-4535, Fax: 425-918-5592, TTY: 711, Email [AppealsDepartmentInquiries@Premera.com](mailto:AppealsDepartmentInquiries@Premera.com). You can file a grievance in person or by mail, fax, or email. If you need help filing a grievance, the Civil Rights Coordinator is available to help you. You can also file a civil rights complaint with the U.S. Department of Health and Human Services, Office for Civil Rights, electronically through the Office for Civil Rights Complaint Portal, available at <https://ocrportal.hhs.gov/ocr/portal/lobby.jsf>, or by mail or phone at: U.S. Department of Health and Human Services, 200 Independence Ave SW, Room 509F, HHH Building, Washington, D.C. 20201, 1-800-368-1019, 800-537-7697 (TDD). Complaint forms are available at <http://www.hhs.gov/ocr/office/file/index.html>.

**Washington residents:** You can also file a civil rights complaint with the Washington State Office of the Insurance Commissioner, electronically through the Office of the Insurance Commissioner Complaint Portal available at <https://www.insurance.wa.gov/file-complaint-or-check-your-complaint-status>, or by phone at 800-562-6900, 360-586-0241 (TDD). Complaint forms are available at <https://fortress.wa.gov/oic/online-services/cc/pub/complaintinformation.aspx>.

**Alaska residents:** Contact the Alaska Division of Insurance via email at [insurance@alaska.gov](mailto:insurance@alaska.gov), or by phone at 907-269-7900 or 1-800-INSURAK (in-state, outside Anchorage).

## Language Assistance

**ATENCIÓN:** si habla español, tiene a su disposición servicios gratuitos de asistencia lingüística. Llame al 800-722-1471 (TTY: 711).

**PAUNAWA:** Kung nagsasalita ka ng Tagalog, maaari kang gumamit ng mga serbisyo ng tulong sa wika nang walang bayad. Tumawag sa 800-722-1471 (TTY: 711).

**注意:** 如果您使用繁體中文，您可以免費獲得語言援助服務。請致電 800-722-1471 (TTY: 711)。

**CHÚ Ý:** Nếu bạn nói Tiếng Việt, có các dịch vụ hỗ trợ ngôn ngữ miễn phí dành cho bạn. Gọi số 800-722-1471 (TTY: 711).

**주의:** 한국어를 사용하시는 경우, 언어 지원 서비스를 무료로 이용하실 수 있습니다. 800-722-1471 (TTY: 711) 번으로 전화해 주십시오.

**ВНИМАНИЕ:** Если вы говорите на русском языке, то вам доступны бесплатные услуги перевода. Звоните 800-722-1471 (телетайп: 711).

**LUS CEEV:** Yog tias koj hais lus Hmoob, cov kev pab txog lus, muaj kev pab dawb rau koj. Hu rau 800-722-1471 (TTY: 711).

**MO LOU SILAFIA:** Afai e te tautala Gagana fa'a Sāmoa, o loo iai auunaga fesoasoan, e fai fua e leai se totagi, mo oe, Telefoni mai: 800-722-1471 (TTY: 711).

**ໂປດອຸລາ:** ຖ້າວ່າ ທ່ານເວົ້າພາສາ ລາວ, ການບໍລິການຊ່ວຍເຫຼືອດ້ານພາສາ, ໂດຍບໍ່ຄ່າສ່ຽງຄ່າ, ຄມມນມີພ້ອມໃຫ້ທ່ານ. ໂທ 800-722-1471 (TTY: 711).

**注意事項:** 日本語を話される場合、無料の言語支援をご利用いただけます。800-722-1471 (TTY:711) まで、お電話にてご連絡ください。

**PAKDAAR:** Nu saritaem ti Ilocano, ti serbisyo para ti baddang ti lengguahe nga awanan bayadna, ket sidadaan para kenyan. Awagan ti 800-722-1471 (TTY: 711).

**УВАГА!** Якщо ви розмовляєте українською мовою, ви можете звернутися до безкоштовної служби мовної підтримки. Телефонуйте за номером 800-722-1471 (телетайп: 711).

**ប្រយ័ត្ន:** បើសិនជាអ្នកនិយាយ ភាសាខ្មែរ, សេវាជំនួយផ្នែកភាសា ដោយមិនគិតលុយ គឺអាចមានសំរាប់អ្នក។ ចូរ ទូរស័ព្ទ 800-722-1471 (TTY: 711)។

**ማስታወሻ:** የሚናገሩት ቋንቋ አማርኛ ከሆነ የትርጉም አርዳታ ድርጅቶች: በነጻ ሊያግኙዎት ተዘጋጅተዋል: ወደ ሚከተለው ቁጥር ይደውሉ 800-722-1471 (መስማት ለተሳናቸው: 711).

**XIYYEEFFANNAA:** Afaan dubbattu Oroomiffa, tajaajjila gargaarsa afaanii, kanfaltiidhaan ala, ni argama. Bilbilaa 800-722-1471 (TTY: 711).

**ملحوظة:** إذا كنت تتحدث اذكر اللغة، فإن خدمات المساعدة اللغوية تتوافر لك بالمجان. اتصل برقم 800-722-1471 (رقم هاتف الصم والبكم: 711).

**ਧਿਆਨ ਦਿਓ:** ਜੇ ਤੁਸੀਂ ਪੰਜਾਬੀ ਬੋਲਦੇ ਹੋ, ਤਾਂ ਭਾਸ਼ਾ ਵਿੱਚ ਸਹਾਇਤਾ ਸੇਵਾ ਤੁਹਾਡੇ ਲਈ ਮੁਫਤ ਉਪਲਬਧ ਹੈ। 800-722-1471 (TTY: 711) 'ਤੇ ਕਾਲ ਕਰੋ।

**ထိပ်စီး:** ถ้าคุณพูดภาษาไทยคุณสามารถใช้บริการช่วยเหลือทางภาษาได้ฟรี โทร 800-722-1471 (TTY: 711).

**ACHTUNG:** Wenn Sie Deutsch sprechen, stehen Ihnen kostenlos sprachliche Hilfsdienstleistungen zur Verfügung. Rufnummer: 800-722-1471 (TTY: 711).

**UWAGA:** Jeżeli mówisz po polsku, możesz skorzystać z bezpłatnej pomocy językowej. Zadzwoń pod numer 800-722-1471 (TTY: 711).

**ATANSYON:** Si w pale Kreyòl Ayisyen, gen sèvis èd pou lang ki disponib gratis pou ou. Rele 800-722-1471 (TTY: 711).

**ATTENTION:** Si vous parlez français, des services d'aide linguistique vous sont proposés gratuitement. Appelez le 800-722-1471 (ATS: 711).

**ATENÇÃO:** Se fala português, encontram-se disponíveis serviços linguísticos, grátis. Ligue para 800-722-1471 (TTY: 711).

**ATTENZIONE:** In caso la lingua parlata sia l'italiano, sono disponibili servizi di assistenza linguistica gratuiti. Chiamare il numero 800-722-1471 (TTY: 711).

**توجہ:** اگر بہ زبان فارسی گفتگو می کنید، تسهیلات زبانی بصورت رایگان برای شما فراهم می باشد. با 800-722-1471 (TTY: 711) تماس بگیرید.