

MEDICAL POLICY – 2.01.38

Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease

BCBSA Ref. Policy: 2.01.38

Effective Date: Feb. 1, 2020

Last Revised: Nov. 1, 2020

Replaces: N/A

RELATED MEDICAL POLICIES:


2.01.91 Peroral Endoscopic Myotomy for Treatment of Esophageal Achalasia

2.01.533 Sinus Surgery

7.01.137 Magnetic Esophageal Sphincter Augmentation to Treat Gastroesophageal Reflux Disease

Select a hyperlink below to be directed to that section.

[POLICY CRITERIA](#) | [CODING](#) | [RELATED INFORMATION](#)
[EVIDENCE REVIEW](#) | [REFERENCES](#) | [HISTORY](#)

 Clicking this icon returns you to the hyperlinks menu above.

Introduction

GERD — gastroesophageal reflux disease — is a long-term medical condition. It’s a digestive problem that affects the ring of muscles between the esophagus (the tube that carries swallowed food to the stomach) and the stomach. When food is swallowed, the muscles at the end of the esophagus open so food can pass into the stomach. The muscles then close to prevent acid and stomach contents from backing up into the esophagus. In GERD, however, the ring of muscles is too weak, and acid can leak back up into the esophagus. GERD is usually treated with changes to lifestyle and diet, or medications, or in some cases a surgery called fundoplication. A number of other treatments have been studied. These include a procedure that is done through the mouth that wraps the upper part of the stomach around the esophagus, the use of radiofrequency energy to try to improve the barrier between the stomach and the esophagus, and the placement of implants or fillers in the esophagus. These procedures are investigational (unproven). More studies are needed to determine if they are as effective as other standard treatments.

Note: The Introduction section is for your general knowledge and is not to be taken as policy coverage criteria. The rest of the policy uses specific words and concepts familiar to medical professionals. It is intended for providers. A provider can be a person, such as a doctor, nurse, psychologist, or dentist. A provider also can be a place where medical care is given, like a hospital, clinic, or lab. This policy informs them about when a service may be covered.

Policy Coverage Criteria

Service	Investigational
Transoral incisionless fundoplication (TIF)	Transoral incisionless fundoplication (TIF) (ie, Esophyx®) is considered investigational as a treatment of gastroesophageal reflux disease.
Transesophageal radiofrequency	Transesophageal radiofrequency to create submucosal thermal lesions of the gastroesophageal junction (ie, Stretta® procedure) is considered investigational as a treatment of gastroesophageal reflux disease.
Endoscopic submucosal implantation of a prosthesis or injection of a bulking agent	Endoscopic submucosal implantation of a prosthesis (ie, Gatekeeper™ Reflux Repair System) or injection of a bulking agent (eg, polymethylmethacrylate beads [PMMA], zirconium oxide spheres [ie, Durasphere®]) is considered investigational as a treatment of gastroesophageal reflux disease.

Coding

Code	Description
CPT	
43201	Esophagoscopy, flexible, transoral; with directed submucosal injection(s), any substance
43210	Esophagogastroduodenoscopy, flexible, transoral; with esophagogastric fundoplasty, partial or complete, includes duodenoscopy when performed
43236	Esophagogastroduodenoscopy, flexible, transoral; with directed submucosal injection(s), any substance
43257	Esophagogastroduodenoscopy, flexible, transoral; with delivery of thermal energy to the muscle of lower esophageal sphincter and/or gastric cardia, for treatment of gastroesophageal reflux disease
43266	Esophagogastroduodenoscopy, flexible, transoral; with placement of endoscopic stent (includes pre- and post-dilation and guide wire passage, when performed)
43499	Unlisted procedure, esophagus



Note: CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). HCPCS codes, descriptions and materials are copyrighted by Centers for Medicare Services (CMS).

Related Information

N/A

Evidence Review

Description

Transesophageal endoscopic therapies are being developed for the treatment of gastroesophageal reflux disease (GERD). A variety of procedures are being evaluated, including transesophageal (or transoral) incisionless fundoplication (TIF), application of radiofrequency energy, and injection/implantation of prosthetic devices or bulking agents.

Background

Gastroesophageal Reflux Disease

Gastroesophageal reflux disease (GERD) is a common disorder characterized by heartburn and other symptoms related to reflux of stomach acid into the esophagus. Nearly all individuals experience such symptoms at some point in their lives; a smaller number have chronic symptoms and are at risk for complications of GERD. The prevalence of GERD has been estimated to be 10% to 20% in the Western world, with a lower prevalence in Asia.¹

Pathophysiology

The pathophysiology of GERD involves excessive exposure of the esophagus to stomach acid, which occurs for several reasons. There can be an incompetent barrier between the esophagus and stomach, either due to dysfunction of the lower esophageal sphincter or incompetence of



the diaphragm. Another mechanism is abnormally slow clearance of stomach acid. In this situation, delayed clearance leads to an increased reservoir of stomach acid and a greater tendency to reflux.

In addition to troubling symptoms, some patients will have more serious disease, which results in complications such as erosive esophagitis, dysphagia, Barrett esophagus, and esophageal carcinoma. Pulmonary complications may result from aspiration of stomach acid into the lungs and can include asthma, pulmonary fibrosis and bronchitis, or symptoms of chronic hoarseness, cough, and sore throat.

Treatment

Guidelines on the management of GERD emphasize initial medical management. Weight loss, smoking cessation, elevating the head of the bed, and elimination of food triggers are all recommended in recent practice guidelines.¹ Proton pump inhibitors have been shown to be the most effective medical treatment. In a Cochrane systematic review, van Pinxteren et al (2010) reported that proton pump inhibitors demonstrated superiority to H₂-receptor agonists and prokinetics in both network meta-analyses and direct comparisons.²

Surgical Treatment

The most common surgical procedure used for GERD is laparoscopic Nissen fundoplication. Fundoplication involves wrapping a portion of the gastric fundus around the distal esophagus to increase lower esophageal sphincter pressure. If a hiatal hernia is present, the procedure also restores the position of the lower esophageal sphincter to the correct location. Laparoscopic fundoplication was introduced in 1991 and has been rapidly adopted because it avoids complications associated with an open procedure.

Although fundoplication results in a high proportion of patients reporting symptom relief, complications can occur and sometimes require conversion to an open procedure. Patients who have relief of symptoms of GERD after fundoplication may have dysphagia or gas-bloat syndrome (excessive gastrointestinal gas).



Other Treatment Options

Due in part to the high prevalence of GERD, there has been interest in creating a minimally invasive transesophageal therapeutic alternative to open or laparoscopic fundoplication or chronic medical therapy. This type of procedure may be considered natural orifice transluminal surgery. Three types of procedures have been investigated.

1. Transesophageal endoscopic gastroplasty (gastroplication, transoral incisionless fundoplication) can be performed as an outpatient procedure. During this procedure, the fundus of the stomach is folded and then held in place with staples or fasteners that are deployed by the device. The endoscopic procedure is designed to recreate a valve and barrier to reflux.
2. Radiofrequency energy has been used to produce submucosal thermal lesions at the gastroesophageal junction. (This technique has also been referred to as the Stretta procedure.) Specifically, radiofrequency energy is applied through 4 electrodes inserted into the esophageal wall at multiple sites both above and below the squamocolumnar junction. The mechanism of action of the thermal lesions is not precisely known but may be related to ablation of the nerve pathways responsible for sphincter relaxation or may induce a tissue-tightening effect related to heat-induced collagen contraction and fibrosis.
3. Submucosal injection or implantation of a prosthetic or bulking agent to enhance the volume of the lower esophageal sphincter has also been investigated.

One bulking agent, pyrolytic carbon-coated zirconium oxide spheres (Durasphere®), is being evaluated. The Gatekeeper™ Reflux Repair System (Medtronic) uses a soft, pliable, expandable prosthesis made of a polyacrylonitrile-based hydrogel. The prosthesis is implanted into the esophageal submucosa, and with time, the prosthesis absorbs water and expands, creating bulk in the region of implantation. U.S. Food and Drug Administration (FDA) product code: DQX. Endoscopic submucosal implantation of polymethylmethacrylate beads into the lower esophageal folds has also been investigated.

The Agency for Healthcare Research and Quality published a systematic review of management strategies for GERD in 2005, which was updated by Ip et al (2011).^{3,4} The 2005 comparative effectiveness review evaluated studies on the EndoCinch Suturing System, Stretta, Enteryx, and the NDO Plicator.³ The 2011 update excluded Enteryx and the NDO Plicator, because they were no longer available in the United States, and added the EsophyX procedure (endoscopic fundoplication), which was commercialized after the 2005 review.⁴ The 2011 report concluded that, for the 3 available endoscopic procedures (EndoCinch, Stretta, EsophyX), effectiveness remained substantially uncertain for the long-term management of GERD. All procedures have



been associated with complications, including dysphagia, infection/fever, and bloating, although bloating and dysphagia are also adverse events of laparoscopic fundoplication.⁵ A review of endoscopic treatment of GERD by Hummel and Richards (2015) noted that EndoCinch is no longer manufactured.⁶

Summary of Evidence

For individuals with GERD and hiatal hernia of 2 cm or less that is not controlled by PPIs who receive TIF (eg, EsophyX), the evidence includes two randomized controlled trials (RCTs) comparing TIF with PPI therapy, nonrandomized studies comparing TIF with fundoplication, and case series with longer term follow-up. The relevant outcomes are symptoms, change in disease status, quality of life, medication use, and treatment-related morbidity. The highest quality RCT (RESPECT) was sham-controlled that compared TIF with PPI therapy while the other RCT (TEMPO) compared TIF with maximum PPI therapy. Both trials found a significant benefit of TIF on the primary outcome measure in about 65% of patients. The sham-controlled trial reported improvement in 45% of the sham-controlled group and no benefit on secondary subjective outcome measures. The nonblinded RCT found significant improvements in subjective measures but no difference in objective outcome measures when compared with PPI therapy. Together, these trial results would suggest a strong placebo effect of the surgery and a modest benefit of TIF in patients whose symptoms were not controlled by PPIs. For these patients, the most appropriate comparator would be laparoscopic fundoplication. Studies comparing TIF with fundoplication have limitations that include earlier TIF procedures and unbalanced groups at baseline and are inadequate to determine relative efficacy. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals with GERD and a hiatal hernia of 2 cm or less that is controlled by PPIs who receive TIF (eg, EsophyX), the evidence includes two RCTs and observational studies with longer term follow-up. The relevant outcomes are symptoms, change in disease status, quality of life, medication use, and treatment-related morbidity. A sham-controlled trial found that the time to resume PPI therapy was longer following TIF and the remission rate was higher, indicating that TIF is more effective than no therapy. The nonblinded RCT found a benefit of TIF compared with continued PPI therapy for subjective measures, but not for the objective measures of pH normalization and esophagitis. These results raise questions about a possible placebo effect for the procedure. Also, observational studies have indicated a loss of treatment effectiveness over time. Adverse events associated with the procedure (eg, perforation) may be severe. At present, the available evidence does not support the use of this intervention in patients whose symptoms



are adequately controlled by medical therapy. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals with GERD who receive endoscopic radiofrequency energy (eg, Stretta), the evidence includes four small RCTs, a nonrandomized comparative study, and observational studies with longer term follow-up. The relevant outcomes are symptoms, change in disease status, quality of life, medication use, and treatment-related morbidity. The RCTs report improvements in symptoms and quality of life following treatment with RF energy compared with sham controls. However, objective measures of GERD and a meta-analysis of these studies found no significant improvements in outcomes, raising questions about the mechanism of the symptom relief. Symptom relief is reported to be lower than after fundoplication, and reoperations greater. Larger RCTs with longer follow-up, preferably compared with fundoplication, are needed to define the risks and benefits of this procedure better. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals with GERD who receive esophageal bulking agents, the evidence includes an RCT and case series. The relevant outcomes are symptoms, change in disease status, quality of life, medication use, and treatment-related morbidity. The RCT for a single product was terminated early due to lack of efficacy, while other products have only case series to support use. High-quality data from large RCTs are needed to compare bulking procedures with both sham controls and with the currently accepted treatments for GERD (ie, drug therapy, laparoscopic fundoplication). Well-designed trials should use standardized outcome measures to examine whether subjective improvement (eg, discontinuation of medication therapy, GERD–HRQL scores) is supported by objective improvement (eg, esophageal acid exposure). The evidence is insufficient to determine the effects of the technology on health outcomes.

Ongoing and Unpublished Clinical Trials

Some currently ongoing trials that might influence this review are listed in [Table 1](#).

Table 1. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
Ongoing			



NCT No.	Trial Name	Planned Enrollment	Completion Date
NCT02366169 ^a	A Worldwide Post-Market Surveillance Registry to Assess the Medigus Ultrasonic Surgical Endostapler (MUSE™) System for the Treatment of GERD	200	Dec 2019
Unpublished			
NCT01110811 ^a	A Randomized Controlled Trial Comparing Transoral Incisionless Fundoplication (TIF) Using EsophyX With Sham Procedure for the Treatment of PPI Dependent GERD: the TIF vs. Sham Study	60	Dec 2018 (completed)
NCT01682265	Stretta in Reflux Uncontrolled by Intake of Inhibitors of Protons Pump (IPP)-The SIRUP Trial-Multicentric, Randomized, Double Blind, Prospective Study	62	Nov 2018 (completed)
NCT01118585 ^a	Prospective Outcome Evaluation of Transoral Incisionless Fundoplication (TIF) for the Treatment of Gastroesophageal Reflux Disease (GERD): The TIF Registry Study	500	Dec 2018 (completed)

NCT: national clinical trial

^a Denotes industry-sponsored or cosponsored trial

Clinical Input Received from Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

2015 Input

In response to requests for clinical input on transesophageal radiofrequency (Stretta) as a treatment of gastroesophageal reflux disease (GERD), input was received from 1 physician specialty society (two reviewers) and three academic medical centers while this policy was under review for 2015. Input was mixed on the treatment of GERD with transesophageal radiofrequency to create submucosal thermal lesions of the gastroesophageal junction (ie, Stretta). Potential conflicts of interest were noted by two reviewers.



2011 Input

In response to requests for clinical input on transoral incisionless fundoplication (TIF) using EsophyX[®], input was received from two physician specialty societies and four academic medical centers while this policy was under review in 2011. Reviewers agreed that TIF differed sufficiently from laparoscopic Nissen fundoplication to warrant evaluation as a separate procedure. Reviewers considered TIF (ie, EsophyX[®]) to be investigational for the treatment of GERD.

Practice Guidelines and Position Statements

American Society for Gastrointestinal Endoscopy

The American Society for Gastrointestinal Endoscopy (2015) published guidelines on endoscopic procedures for GERD.³⁷ In its review of the EsophyX and Stretta procedures, the Society noted some positive findings but discrepancies between subjective and objective outcome measures or a lack of objective outcome measures in reported trials, concluding that these techniques represent “potentially new therapeutic indications for GI endoscopy,” but that prospective trials using objective measures of GERD as the primary end point could be useful in defining the clinical role of these procedures.

American College of Gastroenterology

Updated guidelines released by the American College of Gastroenterology (2013) indicated the use of current endoscopic therapy or SAGES; (2017) provided a clinical spotlight review on endoluminal treatments for GERD.³⁸ The SAGES gave a strong recommendation based on moderate-quality evidence that TIF using EsophyX can be performed with an acceptable safety risk in selected patients. The SAGES concluded that EsophyX results in better control of GERD symptoms than proton pump inhibitor (PPI) treatment in the short term (six months) and leads to similar improvements in objective GERD measures compared with PPIs. TIF appears to lose effectiveness during longer-term follow-up and is associated with moderate patient satisfaction scores. SAGES found no comparative, controlled trials between TIF and surgical fundoplication, but preliminary evidence suggested that surgical fundoplication can be used safely after TIF failure.



The SAGES gave a strong recommendation based on moderate-quality evidence that Stretta is safe for adults and significantly improves health-related quality of life score, heartburn scores, the incidence of esophagitis, and esophageal acid exposure in patients with GERD. Stretta was found to decrease PPI use by about 50%, and be more effective than PPIs, but less effective compared to fundoplication. The effectiveness of the procedure decreases over time.

American Society of General Surgeons

The American Society of General Surgeons (2011) issued a position statement on transoral fundoplication stating that “ASGS supports the use of transoral fundoplication by trained General Surgeons for the treatment of symptomatic chronic gastroesophageal reflux disease (GERD) in patients who fail to achieve satisfactory response to a standard dose of proton pump inhibitor (PPI) therapy or for those who wish to avoid the need for a lifetime of medication dependence.”³⁹

National Institute for Health and Care Excellence

The National Institute for Health and Care Excellence (NICE; 2013) updated its guidance on endoscopic radiofrequency treatment for GERD, concluding: “The evidence on the safety of endoscopic radiofrequency ablation for gastro-esophageal reflux disease is adequate in the short and medium term but there is uncertainty about longer term outcomes. With regard to efficacy, there is evidence of symptomatic relief but objective evidence on reduction of reflux is inconclusive. Therefore, this procedure should only be used with special arrangements...”⁴⁰ The NICE noted “concern on the part of some specialists about the possibility that symptoms may improve as a result of denervation caused by the procedure; if that were the case then failure to recognize and treat reflux might lead to complications in the long term.”

The NICE (2011) issued guidance on endoluminal gastroplication for GERD, concluding that “The evidence on endoluminal gastroplication for gastroesophageal reflux disease raises no major safety concerns. Evidence from a number of RCTs [randomized controlled trials] shows a degree of efficacy in terms of reduced medication requirement in the short term, but changes in other efficacy outcomes are inconsistent, and there is no good evidence of sustained improvement in esophageal pH measurements. ...”⁴¹

The NICE (2004) issued guidance on bulking agents for GERD found that “Current evidence on the safety and efficacy of endoscopic injection of bulking agents for gastro-esophageal reflux disease does not appear adequate for this procedure to be used without special



arrangements...."⁴² The NICE (2016) removed guidance on endoscopic bulking agents/hydrogel implants from guidelines on treatment for "dyspepsia and gastro-esophageal reflux" because the product had been withdrawn by the manufacturer.

Medicare National Coverage

There is no national coverage determination.

Regulatory Status

In 2007, Esophyx® (EndoGastric Solutions) was cleared for marketing by FDA through the 510(k) process for full-thickness plication. In 2016, EsophyX® Z Device with SerosaFuse Fasteners was cleared for marketing by the FDA through the 510(k) process (K160960) for use in transoral tissue approximation, full thickness plication, ligation in the gastrointestinal tract, narrowing the gastroesophageal junction, and reduction of hiatal hernia of 2 cm or less in patients with symptomatic chronic GERD.⁷ In June 2017, EsophyX2 HD and the third-generation EsophyX Z Devices with SerosaFuse fasteners and accessories were cleared for marketing by FDA through the 510(k) process (K171307) for expanded indications, including patients who require and respond to pharmacologic therapy and in patients with hiatal hernias larger than 2 cm when a laparoscopic hiatal hernia repair reduces a hernia to 2 cm or less.⁸ FDA code: ODE.

The Medigus SRS Endoscopic Stapling System (MUSE, Medigus) was cleared for marketing by FDA through the 510(k) process in 2012 (K120299) and 2014 (K132151). MUSE is intended for endoscopic placement of surgical staples in the soft tissue of the esophagus and stomach to create anterior partial fundoplication for treatment of symptomatic chronic GERD in patients who require and respond to pharmacologic therapy. FDA product code: ODE.

In 2000, the CSM Stretta® System was cleared for marketing by FDA through the 510(k) process for general use in the electrosurgical coagulation of tissue and is specifically intended for use in the treatment of GERD. Stretta® is currently manufactured by Mederi Therapeutics. FDA product code: GEI.

Durasphere® is a bulking agent approved for treatment of urinary and fecal incontinence (see [Related Policies](#)). Use of this product for esophageal reflux would be considered off-label use. The website of Carbon Medical Technologies states that the Durasphere® GR product is "intended to treat problems associated with GERD" but is considered an investigational device in the United States.



References

1. Katz PO, Gerson LB, Vela MF. Guidelines for the diagnosis and management of gastroesophageal reflux disease. *Am J Gastroenterol*. Mar 2013;108(3):308-328; quiz 329. PMID 23419381.
2. van Pinxteren B, Sigterman KE, Bonis P, et al. Short-term treatment with proton pump inhibitors, H2-receptor antagonists and prokinetics for gastro-oesophageal reflux disease-like symptoms and endoscopy negative reflux disease. *Cochrane Database Syst Rev*. Nov 10 2010(11):CD002095. PMID 21069670.
3. Ip S, Bonis P, Tatsoni A, et al. Comparative Effectiveness of Management Strategies for Gastroesophageal Reflux Disease (Evidence Report/Technology Assessment No. 1). Rockville, MD: Agency for Healthcare Research and Quality; 2005.
4. Ip S, Chung M, Moorthy D, et al. Management strategies for gastroesophageal reflux disease: An update (Comparative Effectiveness Review No. 29). Rockville, MD: Agency for Healthcare Research and Quality; 2011.
5. Humphries LA, Hernandez JM, Clark W, et al. Causes of dissatisfaction after laparoscopic fundoplication: the impact of new symptoms, recurrent symptoms, and the patient experience. *Surg Endosc*. May 2013;27(5):1537-1545. PMID 23508812.
6. Hummel K, Richards W. Endoscopic treatment of gastroesophageal reflux disease. *Surg Clin North Am*. Jun 2015;95(3):653-667. PMID 25965137.
7. Food and Drug Administration (FDA). 510(k) Summary: EsophyX (K106960). 2016; https://www.accessdata.fda.gov/cdrh_docs/pdf16/K160960.pdf Accessed January 2020.
8. Food and Drug Administration (FDA). EsophyX Summary K171307. 2017; https://www.accessdata.fda.gov/cdrh_docs/pdf17/K171307.pdf Accessed January 2020.
9. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Transesophageal Endoscopic Treatments for Gastroesophageal Reflux Disease. TEC Assessment. 2003;Volume 18:Tab 20.
10. McCarty TR, Itidiare M, Njei B, et al. Efficacy of transoral incisionless fundoplication for refractory gastroesophageal reflux disease: a systematic review and meta-analysis. *Endoscopy*. Jul 2018;50(7):708-725. PMID 29625507.
11. Richter JE, Kumar A, Lipka S, et al. Efficacy of laparoscopic Nissen fundoplication vs transoral incisionless fundoplication or proton pump inhibitors in patients with gastroesophageal reflux disease: a systematic review and network meta-analysis. *Gastroenterology*. Apr 2018;154(5):1298-1308 e1297. PMID 29305934.
12. Hunter JG, Kahrilas PJ, Bell RC, et al. Efficacy of transoral fundoplication vs omeprazole for treatment of regurgitation in a randomized controlled trial. *Gastroenterology*. Feb 2015;148(2):324-333 e325. PMID 25448925.
13. Trad KS, Barnes WE, Simoni G, et al. Transoral incisionless fundoplication effective in eliminating GERD symptoms in partial responders to proton pump inhibitor therapy at 6 months: the TEMPO Randomized Clinical Trial. *Surg Innov*. Feb 2015;22(1):26-40. PMID 24756976.
14. Trad KS, Fox MA, Simoni G, et al. Transoral fundoplication offers durable symptom control for chronic GERD: 3-year report from the TEMPO randomized trial with a crossover arm. *Surg Endosc*. Jun 2017;31(6):2498-2508. PMID 27655380.
15. Trad KS, Barnes WE, Prevou ER, et al. The TEMPO Trial at 5 years: transoral fundoplication (TIF 2.0) Is safe, durable, and cost-effective. *Surg Innov*. Apr 2018;25(2):149-157. PMID 29405886.
16. Hakansson B, Montgomery M, Cadiere GB, et al. Randomised clinical trial: transoral incisionless fundoplication vs. sham intervention to control chronic GERD. *Aliment Pharmacol Ther*. Dec 2015;42(11-12):1261-1270. PMID 26463242.



17. Witteman BP, Conchillo JM, Rinsma NF, et al. Randomized controlled trial of transoral incisionless fundoplication vs. proton pump inhibitors for treatment of gastroesophageal reflux disease. *Am J Gastroenterol*. Apr 2015;110(4):531-542. PMID 25823768.
18. Toomey P, Teta A, Patel K, et al. Transoral incisionless fundoplication: is it as safe and efficacious as a Nissen or Toupet fundoplication? *Am Surg*. Sep 2014;80(9):860-867. PMID 25197871.
19. Frazzoni M, Conigliaro R, Manta R, et al. Reflux parameters as modified by EsophyX or laparoscopic fundoplication in refractory GERD. *Aliment Pharmacol Ther*. Jul 2011;34(1):67-75. PMID 21539587.
20. Stefanidis G, Viazis N, Kotsikoros N, et al. Long-term benefit of transoral incisionless fundoplication using the esophyx device for the management of gastroesophageal reflux disease responsive to medical therapy. *Dis Esophagus*. Feb 01 2017;30(3):1-8. PMID 27868281.
21. Testoni PA, Testoni S, Mazzoleni G, et al. Long-term efficacy of transoral incisionless fundoplication with Esophyx (Tif 2.0) and factors affecting outcomes in GERD patients followed for up to 6 years: a prospective single-center study. *Surg Endosc*. Sep 2015;29(9):2770-2780. PMID 25480624.
22. Testoni PA, Testoni S, Distefano G et al. Transoral incisionless fundoplication with EsophyX for gastroesophageal reflux disease: clinical efficacy is maintained up to 10 years. *Endosc Int Open*, 2019 May 7;7(5). PMID 31058207.
23. Huang X, Chen S, Zhao H, et al. Efficacy of transoral incisionless fundoplication (TIF) for the treatment of GERD: a systematic review with meta-analysis. *Surg Endosc*. Mar 2017;31(3):1032-1044. PMID 27495332.
24. Lipka S, Kumar A, Richter JE. No evidence for efficacy of radiofrequency ablation for treatment of gastroesophageal reflux disease: a systematic review and meta-analysis. *Clin Gastroenterol Hepatol*. Jun 2015;13(6):1058-1067 e1051. PMID 25459556.
25. Corley DA, Katz P, Wo JM, et al. Improvement of gastroesophageal reflux symptoms after radiofrequency energy: a randomized, sham-controlled trial. *Gastroenterology*. Sep 2003;125(3):668-676. PMID 12949712.
26. Aziz AM, El-Khayat HR, Sadek A, et al. A prospective randomized trial of sham, single-dose Stretta, and double-dose Stretta for the treatment of gastroesophageal reflux disease. *Surg Endosc*. Apr 2010;24(4):818-825. PMID 19730952.
27. Arts J, Bisschops R, Blondeau K, et al. A double-blind sham-controlled study of the effect of radiofrequency energy on symptoms and distensibility of the gastro-esophageal junction in GERD. *Am J Gastroenterol*. Feb 2012;107(2):222-230. PMID 22108449.
28. Fass R, Cahn F, Scotti DJ, et al. Systematic review and meta-analysis of controlled and prospective cohort efficacy studies of endoscopic radiofrequency for treatment of gastroesophageal reflux disease. *Surg Endosc*. Dec 2017;31(12):4865-4882. PMID 28233093.
29. Coron E, Sebillé V, Cadiot G, et al. Clinical trial: Radiofrequency energy delivery in proton pump inhibitor-dependent gastro-oesophageal reflux disease patients. *Aliment Pharmacol Ther*. Nov 1 2008;28(9):1147-1158. PMID 18616516.
30. Kalapala R, Shah H, Nabi Z, et al. Treatment of gastroesophageal reflux disease using radiofrequency ablation (Stretta procedure): An interim analysis of a randomized trial. *Indian J Gastroenterol*. Sep 2017;36(5):337-342. PMID 29030802.
31. Liang WT, Yan C, Wang ZG, et al. Early and midterm outcome after laparoscopic fundoplication and a minimally invasive endoscopic procedure in patients with gastroesophageal reflux disease: a prospective observational study. *J Laparoendosc Adv Surg Tech A*. Aug 2015;25(8):657-661. PMID 26258269.
32. Liang WT, Wang ZG, Wang F, et al. Long-term outcomes of patients with refractory gastroesophageal reflux disease following a minimally invasive endoscopic procedure: a prospective observational study. *BMC Gastroenterol*. Oct 10 2014;14:178. PMID 25304252.
33. Noar M, Squires P, Noar E, et al. Long-term maintenance effect of radiofrequency energy delivery for refractory GERD: a decade later. *Surg Endosc*. Aug 2014;28(8):2323-2333. PMID 24562599.
34. Ganz RA, Fallon E, Wittchow T, et al. A new injectable agent for the treatment of GERD: results of the Durasphere pilot trial. *Gastrointest Endosc*. Feb 2009;69(2):318-323. PMID 19185691.



35. Fockens P, Cohen L, Edmundowicz SA, et al. Prospective randomized controlled trial of an injectable esophageal prosthesis versus a sham procedure for endoscopic treatment of gastroesophageal reflux disease. *Surg Endosc.* Jun 2010;24(6):1387-1397. PMID 20198491.
36. Feretis C, Benakis P, Dimopoulos C, et al. Endoscopic implantation of Plexiglas (PMMA) microspheres for the treatment of GERD. *Gastrointest Endosc.* Apr 2001;53(4):423-426. PMID 11275880.
37. ASGE Standards of Practice Committee, Muthusamy VR, Lightdale JR, et al. The role of endoscopy in the management of GERD. *Gastrointest Endosc.* Apr 2015;81(6):1305-1310. PMID 25863867.
38. Society of American Gastrointestinal and Endoscopic Surgeons. Clinical Spotlight Review: Endoluminal Treatments for Gastroesophageal Reflux Disease (GERD). 2017; <https://www.sages.org/publications/guidelines/endoluminal-treatments-for-gastroesophageal-reflux-disease-gerd/> Accessed January 2020.
39. American Society of General Surgeons (ASGS). Coverage of Transoral fundoplication. 2011; <https://theasgs.org/position-statements/coverage-of-transoral-fundoplication-2/> Accessed January 2020.
40. National Institute for Health and Care Excellence (NICE). Endoscopic radiofrequency ablation for gastro-oesophageal reflux disease [IPG461]. 2013; <https://www.nice.org.uk/guidance/ipg461> Accessed January 2020.
41. National Institute for Health and Care Excellence (NICE). Endoluminal gastroplication for gastro-oesophageal reflux disease [IPG404]. 2011; <https://www.nice.org.uk/guidance/ipg404> Accessed January 2020.
42. National Institute for Health and Care Excellence (NICE). Endoscopic injection of bulking agents for gastro-oesophageal reflux disease [IPG55]. 2004; <https://www.nice.org.uk/guidance/ipg55> Accessed January 2020.

History

Date	Comments
01/18/01	Add to Medicine Section - New Policy
01/08/02	Replace Policy - Updated with new references; policy statement expanded.
11/12/02	Replace Policy - Policy updated using 2002 TEC Assessment; policy statement unchanged.
05/13/03	Replace Policy - CPT codes updated.
01/01/04	Replace Policy - CPT code updates only.
05/11/04	Replace Policy - Policy updated with reference to 2003 Assessment; policy statement amended to include ENTERYX procedure as investigational. New CPT code added. Title changed.
07/13/04	Replace Policy - Policy updated with literature search; references added; policy statement unchanged.
02/08/05	Replace Policy - Policy updated with new CPT category I code for Stretta added and category III code deletion; policy statement unchanged.
03/22/06	Code update - HCPCs code removed only, no other changes.



Date	Comments
05/09/06	Replace Policy - Policy updated with literature and research; references added; policy statement enhanced
06/16/06	Update Scope and Disclaimer - No other changes.
09/12/06	Replace Policy - Policy updated with information on Plicator procedure, which was incorporated into the benefit statement as an additional investigational treatment; Rationale updated; references added; no actual change in policy statement.
10/9/07	Replace Policy - Policy updated with literature search through April 2007; policy statement unchanged. Another FDA-cleared device (StomaphyX) added to description. References added.
08/12/08	Replace Policy - Policy updated with literature search; no change to the policy statement. References and code added.
12/16/08	Code Update - 0133T deleted no other changes.
09/15/09	Replace Policy - Policy updated with literature search, no change to policy statement. References added.
01/12/10	Cross Reference Update - No other changes.
04/13/10	Minor update - No other changes.
10/12/10	Replace Policy - Policy updated with literature review, reference numbers 41-52 added. No change in policy statements.
10/11/11	Replace Policy – Policy updated with literature review through May 2011; Rationale section revised; policy statements on biocompatible polymer and PMMA beads combined as bulking agents; remains investigational.
02/14/12	Replace Policy – Clinical input reviewed; reference 3 added and references reordered; policy statements unchanged.
08/27/12	Update Coding Section – ICD-10 codes are now effective 10/01/2014.
10/18/12	Update Related Policies – Add 7.01.137.
01/29/13	Replace policy. Policy updated with literature review through September 2012; references 15, 19, 26 and 29 added and references reordered; policy statements unchanged. Add Related Policy 2.01.58.
08/15/13	Update Related Policies. Remove deleted policy 2.01.520 and add 2.01.20.
01/21/14	Replace policy. Policy updated with literature review through October 16, 2013; references added and reordered; policy statements unchanged. CPT coding updated; 43212 added to the policy and descriptors updated on others. ICD-9 diagnosis and ICD-10-CM codes removed; policy not adjudicated by diagnoses.
08/18/14	Update Related Policies. Remove 2.01.20 and 2.01.81 as they were archived.
12/03/14	Update Related Policies. Add 2.01.91.



Date	Comments
04/24/15	Annual Review. Policy updated with literature review through October 8, 2014; clinical input reviewed; Rationale revised; references 8, 11, and 17 added and some references removed; NDO Plicator, EndoCinch, and Enteryx removed from policy because they are no longer available in the US. Remove ICD-9 and ICD-10 procedure codes; these are not utilized in policy adjudication.
12/16/15	Update Related Policies. Remove 2.01.58 as it is archived.
12/01/16	Annual Review, approved November 8, 2016. Policy updated with literature review through August 2016. References 5-11, 14,17,19,21,28,30 added. No change to policy statements.
01/01/17	Coding update. Added CPT code 43210, removed 43200, 43212, 43232, and 43266.
12/01/17	Annual Review, approved November 9, 2017. Policy updated with literature review through November 2017. Policy statement unchanged. Updated the Practice Guidelines and Position Statements section.
05/01/18	Annual Review, approved April 18, 2018. Policy updated with literature review through October 2017; new references added. Policy statements unchanged.
02/01/19	Annual Review, approved January 4, 2019. Policy updated with literature review through September 2018; references 10-11, 15, 23, and 30 added; references 42-43 updated. Policy statements unchanged.
02/01/20	Annual Review, approved January 9, 2020. Policy updated with literature review through October 2019; references added. Policy statements unchanged.
07/02/20	Coding update. CPT code 43266 removed.
07/14/20	Coding update. CPT code 43236 removed from policy. This applies to other criteria set.
11/01/20	Coding update. Added CPT 43266 and 43236. Related Policy 2.01.55 Sinus Surgery added.

Disclaimer: This medical policy is a guide in evaluating the medical necessity of a particular service or treatment. The Company adopts policies after careful review of published peer-reviewed scientific literature, national guidelines and local standards of practice. Since medical technology is constantly changing, the Company reserves the right to review and update policies as appropriate. Member contracts differ in their benefits. Always consult the member benefit booklet or contact a member service representative to determine coverage for a specific medical service or supply. CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). ©2020 Premera All Rights Reserved.

Scope: Medical policies are systematically developed guidelines that serve as a resource for Company staff when determining coverage for specific medical procedures, drugs or devices. Coverage for medical services is subject to the limits and conditions of the member benefit plan. Members and their providers should consult the member benefit booklet or contact a customer service representative to determine whether there are any benefit limitations applicable to this service or supply. This medical policy does not apply to Medicare Advantage.



Discrimination is Against the Law

Premera Blue Cross complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex. Premera does not exclude people or treat them differently because of race, color, national origin, age, disability or sex.

Premera:

- Provides free aids and services to people with disabilities to communicate effectively with us, such as:
 - Qualified sign language interpreters
 - Written information in other formats (large print, audio, accessible electronic formats, other formats)
- Provides free language services to people whose primary language is not English, such as:
 - Qualified interpreters
 - Information written in other languages

If you need these services, contact the Civil Rights Coordinator.

If you believe that Premera has failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, or sex, you can file a grievance with:

Civil Rights Coordinator - Complaints and Appeals
PO Box 91102, Seattle, WA 98111
Toll free 855-332-4535, Fax 425-918-5592, TTY 800-842-5357
Email AppealsDepartmentInquiries@Premera.com

You can file a grievance in person or by mail, fax, or email. If you need help filing a grievance, the Civil Rights Coordinator is available to help you.

You can also file a civil rights complaint with the U.S. Department of Health and Human Services, Office for Civil Rights, electronically through the Office for Civil Rights Complaint Portal, available at <https://ocrportal.hhs.gov/ocr/portal/lobby.jsf>, or by mail or phone at: U.S. Department of Health and Human Services
200 Independence Avenue SW, Room 509F, HHH Building
Washington, D.C. 20201, 1-800-368-1019, 800-537-7697 (TDD)
Complaint forms are available at <http://www.hhs.gov/ocr/office/file/index.html>.

Getting Help in Other Languages

This Notice has Important Information. This notice may have important information about your application or coverage through Premera Blue Cross. There may be key dates in this notice. You may need to take action by certain deadlines to keep your health coverage or help with costs. You have the right to get this information and help in your language at no cost. Call 800-722-1471 (TTY: 800-842-5357).

አማርኛ (Amharic):

ይህ ማስታወቂያ አስፈላጊ መረጃ ይዟል። ይህ ማስታወቂያ ስለ ማመልከቻዎ ወይም የ Premera Blue Cross ሽፋን አስፈላጊ መረጃ ሊኖረው ይችላል። በዚህ ማስታወቂያ ውስጥ ቁልፍ ቀናት ሊኖሩ ይችላሉ። የጤና ሽፋንዎን ለመጠበቅና በአስፈላጊ እርዳታ ለማግኘት በተውሰኑ የጊዜ ገደቦች እርምጃ መውሰድ ይገባዎት ይሆናል። ይህን መረጃ እንዲያገኙ እና የለምንም ክፍያ በቋንቋዎ እርዳታ እንዲያገኙ መሰታወቅ አለዎት። በስልክ ቁጥር 800-722-1471 (TTY: 800-842-5357) ይደውሉ።

العربية (Arabic):

يحتوي هذا الإشعار على معلومات هامة. قد يحتوي هذا الإشعار على معلومات مهمة بخصوص طلبك أو التخطيط التي تزيد الحصول عليها من خلال Premera Blue Cross. قد تكون هناك تواريخ مهمة في هذا الإشعار. وقد تحتاج لاتخاذ إجراء في تاريخ معينة للحفاظ على تغطيتك الصحية أو للمساعدة في دفع التكاليف. يحق لك الحصول على هذه المعلومات والمساعدة بلغتك دون تكبد أية تكلفة. اتصل بـ 800-722-1471 (TTY: 800-842-5357)

中文 (Chinese):

本通知有重要的訊息。本通知可能有關於您透過 Premera Blue Cross 提交的申請或保險的重要訊息。本通知內可能有重要日期。您可能需要在截止日期之前採取行動，以保留您的健康保險或者費用補貼。您有權利免費以您的母語得到本訊息和幫助。請撥電話 800-722-1471 (TTY: 800-842-5357)。

Oromoo (Cushite):

Beeksisni kun odeeffannoo barbaachisaa qaba. Beeksisni kun sagantaa yookan karaa Premera Blue Cross tiin tajaajila keessan ilaalchisee odeeffannoo barbaachisaa qabaachuu danda'a. Guyyaawwan murteessaa ta'an beeksisa kana keessatti ilaalaa. Tarii kaffaltiidhaan deeggaramuuf yookan tajaajila fayyaa keessaniif guyyaa dhumaa irratti wanti raawwattan jiraachuu danda'a. Kaffaltii irraa bilisa haala ta'een afaan keessaniin odeeffannoo argachuu fi deeggarsa argachuuf mirga ni qabaattu. Lakkoofsa bilbilaa 800-722-1471 (TTY: 800-842-5357) tii bilbilaa.

Français (French):

Cet avis a d'importantes informations. Cet avis peut avoir d'importantes informations sur votre demande ou la couverture par l'intermédiaire de Premera Blue Cross. Le présent avis peut contenir des dates clés. Vous devez peut-être prendre des mesures par certains délais pour maintenir votre couverture de santé ou d'aide avec les coûts. Vous avez le droit d'obtenir cette information et de l'aide dans votre langue à aucun coût. Appelez le 800-722-1471 (TTY: 800-842-5357).

Kreyòl ayisyen (Creole):

Avi sila a gen Enfòmasyon Enpòtan ladann. Avi sila a kapab genyen enfòmasyon enpòtan konsènan aplikasyon w lan oswa konsènan kouvèti asirans lan atravè Premera Blue Cross. Kapab genyen dat ki enpòtan nan avi sila a. Ou ka gen pou pran kèk aksyon avan sèten dat limit pou ka kenbe kouvèti asirans sante w la oswa pou yo ka ede w avèk depans yo. Se dwa w pou resewva enfòmasyon sa a ak asistans nan lang ou pale a, san ou pa gen pou peye pou sa. Rele nan 800-722-1471 (TTY: 800-842-5357).

Deutsche (German):

Diese Benachrichtigung enthält wichtige Informationen. Diese Benachrichtigung enthält unter Umständen wichtige Informationen bezüglich Ihres Antrags auf Krankenversicherungsschutz durch Premera Blue Cross. Suchen Sie nach eventuellen wichtigen Terminen in dieser Benachrichtigung. Sie könnten bis zu bestimmten Stichtagen handeln müssen, um Ihren Krankenversicherungsschutz oder Hilfe mit den Kosten zu behalten. Sie haben das Recht, kostenlose Hilfe und Informationen in Ihrer Sprache zu erhalten. Rufen Sie an unter 800-722-1471 (TTY: 800-842-5357).

Hmoob (Hmong):

Tsab ntawv tshaj xo no muaj cov ntshiab lus tseem ceeb. Tej zaum tsab ntawv tshaj xo no muaj cov ntshiab lus tseem ceeb txog koj daim ntawv thov kev pab los yog koj qhov kev pab cuam hnu ntawm Premera Blue Cross. Tej zaum muaj cov hnuv tseem ceeb uas sau rau hauv daim ntawv no. Tej zaum koj kuj yuav tau ua qee yam uas peb kom koj ua tsis pub dhau cov caij nyoog uas teev tseg rau hauv daim ntawv no mas koj thiaj yuav tau txais kev pab cuam kho mob los yog kev pab them tej nqi kho mob ntawd. Koj muaj cai kom lawv muab cov ntshiab lus no uas tau muab sau ua koj hom lus pub dawb rau koj. Hu rau 800-722-1471 (TTY: 800-842-5357).

Iloko (Ilocano):

Daytoy a Pakdaar ket naglaon iti Napateg nga Impormasion. Daytoy a pakdaar mabalin nga adda ket naglaon iti napateg nga impormasion maipanggep iti aplikasyonyo wenno coverage babaen iti Premera Blue Cross. Daytoy ket mabalin dagiti importante a petsa iti daytoy a pakdaar. Mabalin nga adda rumbeng nga aramidenyo nga addang sakbay dagiti partikular a naituding nga aldaw tapno mapagtalinaedyo ti coverage ti salun-atyto wenno tulong kadagiti gastos. Adda karbenganyo a mangala iti daytoy nga impormasion ken tulong iti bukodyo a pagsasao nga awan ti bayadanyo. Tumawag iti numero nga 800-722-1471 (TTY: 800-842-5357).

Italiano (Italian):

Questo avviso contiene informazioni importanti. Questo avviso può contenere informazioni importanti sulla tua domanda o copertura attraverso Premera Blue Cross. Potrebbero esserci date chiave in questo avviso. Potrebbe essere necessario un tuo intervento entro una scadenza determinata per consentirti di mantenere la tua copertura o sovvenzione. Hai il diritto di ottenere queste informazioni e assistenza nella tua lingua gratuitamente. Chiama 800-722-1471 (TTY: 800-842-5357).

日本語 (Japanese):

この通知には重要な情報が含まれています。この通知には、Premera Blue Cross の申請または補償範囲に関する重要な情報が含まれている場合があります。この通知に記載されている可能性がある重要な日付をご確認ください。健康保険や有料サポートを維持するには、特定の期日までに行動を取らなければならない場合があります。ご希望の言語による情報とサポートが無料で提供されます。800-722-1471 (TTY: 800-842-5357)までお電話ください。

한국어 (Korean):

본 통지서에는 중요한 정보가 들어 있습니다. 즉 이 통지서는 귀하의 신청에 관하여 그리고 Premera Blue Cross 를 통한 커버리지에 관한 정보를 포함하고 있을 수 있습니다. 본 통지서에는 핵심이 되는 날짜들이 있을 수 있습니다. 귀하의 귀하의 건강 커버리지를 계속 유지하거나 비용을 절감하기 위해서 일정한 마감일까지 조치를 취해야 할 필요가 있을 수 있습니다. 귀하의 이러한 정보와 도움을 귀하의 언어로 비용 부담없이 얻을 수 있는 권리가 있습니다. 800-722-1471 (TTY: 800-842-5357) 로 전화하십시오.

ລາວ (Lao):

ແຈ້ງການນີ້ມີຂໍ້ມູນສໍາຄັນ. ແຈ້ງການນີ້ອາດຈະມີຂໍ້ມູນສໍາຄັນກ່ຽວກັບຄໍາຮ້ອງສະໝັກ ຫຼື ຄວາມຄົມຄອງປະກັນໄພຂອງທ່ານຜ່ານ Premera Blue Cross. ອາດຈະມີວັນທີ່ສໍາຄັນໃນແຈ້ງການນີ້. ທ່ານອາດຈະຈໍາເປັນຕ້ອງດໍາເນີນການຕາມກຳນົດ ເວລາສະເພາະເພື່ອຮັກສາຄວາມຄົມຄອງປະກັນສະພາບ ຫຼື ຄວາມຊ່ວຍເຫຼືອເວັ້ນເວີ້ ຄ່າໃຊ້ຈ່າຍຂອງທ່ານໄດ້. ທ່ານມີສິດໄດ້ຮັບຂໍ້ມູນນີ້ ແລະ ຄວາມຊ່ວຍເຫຼືອເປັນພາສາຂອງທ່ານໂດຍບໍ່ເສຍຄ່າ. ໃຫ້ໃບທາ 800-722-1471 (TTY: 800-842-5357).

ភាសាខ្មែរ (Khmer):

សេចក្តីជូនដំណឹងនេះមានព័ត៌មានយ៉ាងសំខាន់។ សេចក្តីជូនដំណឹងនេះប្រហែលជាមានព័ត៌មានយ៉ាងសំខាន់អំពីទម្រង់បែបបទ ឬការរៀបចំរបស់អ្នកកាមរយ: Premera Blue Cross ។ ប្រហែលជាមាន កាលបរិច្ឆេទសំខាន់នៅក្នុងសេចក្តីជូនដំណឹងនេះ។ អ្នកប្រហែលជាត្រូវការបញ្ជាក់សមត្ថភាព ដល់កិច្ចការផ្ទៃក្នុងនានា នៃស្ត្រីនិងក្មេងៗក្នុងគ្រួសាររបស់អ្នក ឬប្រាក់ចំណូលចេញថ្លៃ។ អ្នកមានសិទ្ធិទទួលបានព័ត៌មាននេះ និងដំណោះស្រាយនៅក្នុងភាសារបស់អ្នកដោយមិនអស់លុយឡើយ។ សូមទូរស័ព្ទ 800-722-1471 (TTY: 800-842-5357)។

ਪੰਜਾਬੀ (Punjabi):

ਇਸ ਨੋਟਿਸ ਵਿਚ ਖਾਸ ਜਾਣਕਾਰੀ ਹੈ. ਇਸ ਨੋਟਿਸ ਵਿਚ Premera Blue Cross ਵਲੋਂ ਤੁਹਾਡੀ ਕਵਰੇਜ ਅਤੇ ਅਰਜੀ ਬਾਰੇ ਮਹੱਤਵਪੂਰਨ ਜਾਣਕਾਰੀ ਹੋ ਸਕਦੀ ਹੈ . ਇਸ ਨੋਟਿਸ ਨਵ ਖਾਸ ਤਾਰੀਖਾਂ ਹੋ ਸਕਦੀਆਂ ਹਨ. ਜੇਕਰ ਤੁਸੀਂ ਜਸਰਤ ਕਵਰੇਜ ਰਿੱਖਣੀ ਹੋਵੇ ਜਾਂ ਓਸ ਦੀ ਲਾਗਤ ਜਵਿੱਚ ਮਦਦ ਦੇ ਇਕੱਠ ਹੋ ਤਾਂ ਤੁਹਾਨੂੰ ਅੰਤਮ ਤਾਰੀਖ ਤੋਂ ਪਹਿਲਾਂ ਢੁੱਝ ਖਾਸ ਕਰਮ ਚੁੱਕਣ ਦੀ ਲੋੜ ਹੋ ਸਕਦੀ ਹੈ ,ਤੁਹਾਨੂੰ ਮੁਫਤ ਵਿੱਚ ਤੋਂ ਅਪਣੀ ਭਾਸ਼ਾ ਵਿੱਚ ਜਾਣਕਾਰੀ ਅਤੇ ਮਦਦ ਪ੍ਰਾਪਤ ਕਰਨ ਦਾ ਅਧਿਕਾਰ ਹੈ ,ਕਾਲ 800-722-1471 (TTY: 800-842-5357).

فارسی (Farsi):

این اعلامیه حاوی اطلاعات مهم میباشد. این اعلامیه ممکن است حاوی اطلاعات مهم درباره فرم تقاضا و یا پوشش بیمه ای شما از طریق Premera Blue Cross باشد. به تاریخ های مهم در این اعلامیه توجه نمایید. شما ممکن است برای حفظ پوشش بیمه تان یا کمک در پرداخت هزینه های درمانی تان، به تاریخ های مشخصی برای انجام کارهای خاصی احتیاج داشته باشید. شما حق این را دارید که این اطلاعات و کمک را به زبان خود به طور رایگان دریافت نمایید. برای کسب اطلاعات با شماره 800-722-1471 (کلیران TTY تماس باشماره 800-842-5357) تماس برقرار نمایید.

Polskie (Polish):

To ogłoszenie może zawierać ważne informacje. To ogłoszenie może zawierać ważne informacje odnośnie Państwa wniosku lub zakresu świadczeń poprzez Premera Blue Cross. Prosimy zwrócić uwagę na kluczowe daty, które mogą być zawarte w tym ogłoszeniu aby nie przekroczyć terminów w przypadku utrzymania polisy ubezpieczeniowej lub pomocy związanej z kosztami. Macie Państwo prawo do bezpłatnej informacji we własnym języku. Zadzwońcie pod 800-722-1471 (TTY: 800-842-5357).

Português (Portuguese):

Este aviso contém informações importantes. Este aviso poderá conter informações importantes a respeito de sua aplicação ou cobertura por meio do Premera Blue Cross. Poderão existir datas importantes neste aviso. Talvez seja necessário que você tome providências dentro de determinados prazos para manter sua cobertura de saúde ou ajuda de custos. Você tem o direito de obter esta informação e ajuda em seu idioma e sem custos. Ligue para 800-722-1471 (TTY: 800-842-5357).

Română (Romanian):

Prezenta notificare conține informații importante privind cererea sau acoperirea asigurării dumneavoastră de sănătate prin Premera Blue Cross. Pot exista date cheie în această notificare. Este posibil să fie nevoie să acționați până la anumite termene limită pentru a vă menține acoperirea asigurării de sănătate sau asistența provizorie la costuri. Aveți dreptul de a obține gratuit aceste informații și ajutor în limba dumneavoastră. Sunați la 800-722-1471 (TTY: 800-842-5357).

Русский (Russian):

Настоящее уведомление содержит важную информацию. Это уведомление может содержать важную информацию о вашем заявлении или страховом покрытии через Premera Blue Cross. В настоящем уведомлении могут быть указаны ключевые даты. Вам, возможно, потребуется принять меры к определенным предельным срокам для сохранения страхового покрытия или помощи с расходами. Вы имеете право на бесплатное получение этой информации и помощь на вашем языке. Звоните по телефону 800-722-1471 (TTY: 800-842-5357).

Fa'asamoa (Samoan):

Atonu ua iai i lenei fa'asilasilaga ni fa'amatalaga e sili ona taua e tatau ona e malamalama i ai. O lenei fa'asilasilaga o se fesoasoani e fa'amatala atili i ai i le tulaga o le polokalame, Premera Blue Cross, ua e tau fia maua atu i ai. Fa'amolemole, ia e iloilo fa'alelei i aso fa'apitoa olo'o iai i lenei fa'asilasilaga taua. Masalo o le'a iai ni feau e tatau ona e faia ao le'i aulia le aso ua ta'ua i lenei fa'asilasilaga ina ia e iai pea ma maua fesoasoani mai ai i le polokalame a le Malo olo'o e iai i ai. Olo'o iai iate oe le aia tatau e maua atu i lenei fa'asilasilaga ma lenei fa'matalaga i legagana e te malamalama i ai aunoa ma se togiga tupe. Vili atu i le telefoni 800-722-1471 (TTY: 800-842-5357).

Español (Spanish):

Este Aviso contiene información importante. Es posible que este aviso contenga información importante acerca de su solicitud o cobertura a través de Premera Blue Cross. Es posible que haya fechas clave en este aviso. Es posible que deba tomar alguna medida antes de determinadas fechas para mantener su cobertura médica o ayuda con los costos. Usted tiene derecho a recibir esta información y ayuda en su idioma sin costo alguno. Llame al 800-722-1471 (TTY: 800-842-5357).

Tagalog (Tagalog):

Ang Paunawa na ito ay naglalaman ng mahalagang impormasyon tungkol sa iyong aplikasyon o pagsakop sa pamamagitan ng Premera Blue Cross. Maaaring may mga mahalagang petsa dito sa paunawa. Maaring mangailangan ka na magsagawa ng hakbang sa ilang mga itinakdang panahon upang mapanatili ang iyong pagsakop sa kalusugan o tulong na walang gastos. May karapatan ka na makakuha ng ganiitong impormasyon at tulong sa iyong wika ng walang gastos. Tumawag sa 800-722-1471 (TTY: 800-842-5357).

ไทย (Thai):

ประกาศนี้มีข้อมูลสำคัญ ประกาศนี้อาจมีข้อมูลที่สำคัญเกี่ยวกับกาการสมัครหรือขอบเขตประกันสุขภาพของคุณผ่าน Premera Blue Cross และอาจมีกำหนดการในประกาศนี้ คุณอาจจะต้องดำเนินการภายในกำหนดระยะเวลาที่แน่นอนเพื่อจะรักษาการประกันสุขภาพของคุณหรือการช่วยเหลือที่มีค่าใช้จ่าย คุณมีสิทธิที่จะได้รับข้อมูลและความช่วยเหลือนี้ในภาษาของคุณโดยไม่มีค่าใช้จ่าย โทร 800-722-1471 (TTY: 800-842-5357)

Український (Ukrainian):

Це повідомлення містить важливу інформацію. Це повідомлення може містити важливу інформацію про Ваше звернення щодо страховального покриття через Premera Blue Cross. Зверніть увагу на ключові дати, які можуть бути вказані у цьому повідомленні. Існує імовірність того, що Вам треба буде здійснити певні кроки у конкретні кінцеві строки для того, щоб зберегти Ваше медичне страхування або отримати фінансову допомогу. У Вас є право на отримання цієї інформації та допомоги безкоштовно на Вашій рідній мові. Дзвоніть за номером телефону 800-722-1471 (TTY: 800-842-5357).

Tiếng Việt (Vietnamese):

Thông báo này cung cấp thông tin quan trọng. Thông báo này có thông tin quan trọng về đơn xin tham gia hoặc hợp đồng bảo hiểm của quý vị qua chương trình Premera Blue Cross. Xin xem ngày quan trọng trong thông báo này. Quý vị có thể phải thực hiện theo thông báo đúng trong thời hạn để duy trì bảo hiểm sức khỏe hoặc được trợ giúp thêm về chi phí. Quý vị có quyền được biết thông tin này và được trợ giúp bằng ngôn ngữ của mình miễn phí. Xin gọi số 800-722-1471 (TTY: 800-842-5357).