

## MEDICAL POLICY – 2.01.91

## Peroral Endoscopic Myotomy for Treatment of Esophageal Achalasia

BCBSA Ref. Policy: 2.01.91

Effective Date: Feb. 1, 2019

Last Revised: Jan. 4, 2019

Replaces: N/A

## RELATED MEDICAL POLICIES:


2.01.38 Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease

7.01.137 Magnetic Esophageal Sphincter Augmentation to Treat Gastroesophageal Reflux Disease

8.01.17 Hematopoietic Cell Transplantation for Plasma Cell Dyscrasias, Including Multiple Myeloma and POEMS Syndrome

Select a hyperlink below to be directed to that section.

[POLICY CRITERIA](#) | [CODING](#) | [RELATED INFORMATION](#)  
[EVIDENCE REVIEW](#) | [REFERENCES](#) | [HISTORY](#)

 Clicking this icon returns you to the hyperlinks menu above.

## Introduction

Esophageal achalasia is a rare problem with the esophagus (the swallowing tube). It affects the ability to pass food through the esophagus and into the stomach. The muscles of the esophagus don't move food down, and the ring of muscles at the end of the esophagus don't relax to easily allow food into the stomach. This makes swallowing very difficult. A new surgery, POEM (peroral endoscopic myotomy), is being tried. A viewing scope with a special cutting blade is passed through the mouth and into the esophagus. Part of the muscle layer of the lower part of the esophagus, the sphincter, and the upper part of the stomach is removed. POEM is investigational. More and larger studies are needed to compare POEM with standard surgery to treat esophageal achalasia.

**Note:** The Introduction section is for your general knowledge and is not to be taken as policy coverage criteria. The rest of the policy uses specific words and concepts familiar to medical professionals. It is intended for providers. A provider can be a person, such as a doctor, nurse, psychologist, or dentist. A provider also can be a place where medical care is given, like a hospital, clinic, or lab. This policy informs them about when a service may be covered.

## Policy Coverage Criteria

Service	Investigational
<b>Peroral endoscopic myotomy</b>	<b>Peroral endoscopic myotomy is considered investigational as a treatment for esophageal achalasia.</b>  <b>Note:</b> This policy addresses POEM. A similar acronym, POEMS syndrome, describes a different condition and is addressed in a separate medical policy. Please see <a href="#">Related Policies</a> .

## Coding

Code	Description
<b>CPT</b>	
43499	Unlisted procedure, esophagus

**Note:** CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). HCPCS codes, descriptions and materials are copyrighted by Centers for Medicare Services (CMS).

## Related Information

N/A

## Evidence Review

### Description

Esophageal achalasia is characterized by reduced numbers of neurons in the esophageal myenteric plexuses and reduced peristaltic activity, making it difficult for patients to swallow food and possibly leading to complications such as regurgitation, coughing, choking, aspiration



pneumonia, esophagitis, ulceration, and weight loss. Peroral endoscopic myotomy (POEM) is a novel endoscopic procedure that uses the oral cavity as a natural orifice entry point to perform myotomy of the lower esophageal sphincter. This procedure is intended to reduce the total number of incisions needed and thus the overall invasiveness of surgery.

## Background

### *Esophageal Achalasia*

Esophageal achalasia is characterized by reduced numbers of neurons in the esophageal myenteric plexuses and reduced peristaltic activity, making it difficult for patients to swallow food and possibly leading to complications such as regurgitation, coughing, choking, aspiration pneumonia, esophagitis, ulceration, and weight loss. The estimated U.S. prevalence of achalasia is 10 cases per 100,000, and the estimated incidence is 0.6 cases per 100,000 per year.<sup>1</sup>

### Treatment

Treatment options for achalasia have included pharmacotherapy such as injections with botulinum toxin, pneumatic dilation, and laparoscopic Heller myotomy.<sup>1,2</sup> Although the latter two are considered the standard treatments because of higher success rates and relatively long-term efficacy compared with pharmacotherapy, both are associated with a perforation risk of about 1%. Heller myotomy is the most invasive of the procedures, requiring laparoscopy and surgical dissection of the esophagogastric junction.<sup>2</sup> One-year response rates of 86% and major mucosal tear rates requiring subsequent intervention of 0.6% have been reported.<sup>3</sup>

Peroral endoscopic myotomy (POEM) is a novel endoscopic procedure developed in Japan.<sup>2,4</sup> POEM is performed with the patient under general anesthesia.<sup>5</sup> After tunneling an endoscope down the esophagus toward the esophageal-gastric junction, a surgeon performs the myotomy by cutting only the inner, circular lower esophageal sphincter muscles through a submucosal tunnel created in the proximal esophageal mucosa. POEM differs from laparoscopic surgery, which involves complete division of both circular and longitudinal lower esophageal sphincter muscle layers. Cutting the dysfunctional muscle fibers that prevent the lower esophageal sphincter from opening allows food to enter the stomach more easily.<sup>2,5</sup>

**NOTE:** The acronym POEM in this policy refers to peroral endoscopic myotomy. POEMS syndrome, which uses a similar acronym, is discussed in a separate medical policy (see [Related Policies](#)).



## Summary of Evidence

For individuals with achalasia who receive POEM, the evidence includes systematic reviews of observational studies, nonrandomized comparative studies, and case series. Relevant outcomes are symptoms, functional outcomes, health status measures, resource utilization, and treatment-related morbidity. The comparative studies have primarily reported similar outcomes with POEM and for Heller myotomy in symptom relief, as assessed by the Eckardt score. Some studies have shown shorter length of stay and less postoperative pain with POEM. However, potential imbalances in patient characteristics in these nonrandomized studies might have biased the treatment comparisons. In the case series, treatment success at short follow-up periods was reported for a high proportion of patients treated with POEM. However, incidence of adverse events was relatively high, with POEM-specific complications, including subcutaneous emphysema, pneumothorax, and thoracic effusion, reported across studies. Additionally, a substantial proportion of patients undergoing POEM developed gastroesophageal reflux disease and esophagitis and required treatment. Case series do not permit conclusions about the efficacy of POEM relative to established treatment, and long-term outcomes of the procedure are not well described in the literature. The evidence is insufficient to determine the effects of the technology on health outcomes.

## Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in [Table 1](#).

**Table 1. Summary of Key Trials**

NCT No.	Trial Name	Planned Enrollment	Completion Date
<b>Ongoing</b>			
<a href="#">NCT02138643</a>	Laparoscopy Heller Myotomy With Fundoplication Associated Versus Peroral Endoscopic Myotomy (POEM)	30	Dec 2017 (ongoing)
<a href="#">NCT03228758</a>	Efficacy of Anterior Versus Posterior Myotomy Approach in Peroral Endoscopic Myotomy (POEM) for the Treatment of Achalasia – a Single Operator Analysis	290	Nov 2019



NCT No.	Trial Name	Planned Enrollment	Completion Date
<a href="#">NCT01402518</a>	Observational Study of the Peroral Endoscopic Myotomy (POEM) Procedure	100	Nov 2019
<a href="#">NCT01601678</a>	Endoscopic Versus Laparoscopic Myotomy for Treatment of Idiopathic Achalasia: A Randomized, Controlled Trial	240	Dec 2019
<a href="#">NCT01832779</a>	Prospective Evaluation of the Clinical Utility of Peroral Endoscopic Myotomy (POEM)	600	Dec 2022
<a href="#">NCT01793922</a>	A Prospective Randomized Multi-center Study Comparing Endoscopic Pneumodilation and Per Oral Endoscopic Myotomy (POEM) as Treatment of Idiopathic Achalasia	150	Jan 2023

NCT: national clinical trial

## Practice Guidelines and Position Statements

### *American Gastroenterological Association Institute*

In 2017, the American Gastroenterological Association Institute published a clinical practice update on the use of peroral endoscopic myotomy (POEM) for the treatment of achalasia.<sup>30</sup>

Based on expert review, the Institute made the following recommendations:

- POEM should be performed by experienced physicians in high-volume centers (competence achieved after estimated 20 to 40 procedures).
- If expertise is available, POEM should be considered primary therapy for type III achalasia.
- If expertise is available, POEM should be considered comparable to Heller myotomy for any achalasia syndromes.
- Patients receiving POEM should be considered high risk to develop reflux esophagitis and be advised of management considerations (eg, proton pump inhibitor therapy and/or surveillance endoscopy) prior to undergoing POEM.



## ***American Society of American Gastrointestinal and Endoscopic Surgeons***

In 2014, the American Society of Gastrointestinal and Endoscopic Surgeons issued evidence-based, consensus guidelines on the use of endoscopy in the evaluation and management of dysphagia, including esophageal achalasia.<sup>31</sup> The Society recommended that:

... Endoscopic and surgical treatment options for achalasia should be discussed with the patient. In patients who opt for endoscopic management and are good surgical candidates, pneumatic dilation with large-caliber balloon dilators for the endoscopic treatment of achalasia was recommended... Long-term data and randomized trials comparing peroral endoscopic myotomy to conventional modalities of management are necessary before it can be adopted into clinical practice, but the procedure is becoming more widely used in expert centers.

## ***American College of Gastroenterology***

In 2013, the American College of Gastroenterology issued clinical guidelines on the diagnosis and management of achalasia.<sup>32</sup> POEM was discussed as an emerging therapy and stated to have promise as an alternative to the laparoscopic approach. The guidelines further stated that randomized prospective comparison trials are needed, and the procedure should be performed in the context of clinical trials.

## ***Society of American Gastrointestinal and Endoscopic Surgeons***

In 2012, the Society of American Gastrointestinal and Endoscopic Surgeons issued evidence-based, consensus guidelines on the surgical management of esophageal achalasia. The guidelines stated that the POEM technique "is in its infancy and further experience is needed before providing recommendations."<sup>33</sup>

## ***International Society for Diseases of the Esophagus***

In 2018, the International Society for Diseases of the Esophagus published guidelines on the diagnosis and management of achalasia.<sup>34</sup> The Society convened 51 experts from 11 countries, including several from the United States, to systematically review evidence, assess recommendations using the GRADE system, and vote to integrate the recommendations into



the guidelines (>80% approval required for inclusion). **Table 2** summarizes POEM recommendations.

**Table 2. Recommendations for the Treatment of Achalasia**

Recommendation	LOR	GOR
POEM is an effective therapy for achalasia both in short-term and medium-term follow-up with results comparable to Heller myotomy.	Conditional	Very low
POEM is an effective therapy for achalasia both in short-term and medium-term follow-up with results comparable to pneumatic dilations.	Conditional	Low
Pretreatment information on GERD, nonsurgical options (pneumatic dilation), and surgical options with lower GERD risk (Heller myotomy) should be provided to patient.	Good practice	NA
POEM is feasible and effective for symptom relief in patients previously treated with endoscopic therapies.	Conditional	Very low
POEM may be considered an option for treating recurrent symptoms after laparoscopic Heller myotomy.	Conditional	Low
Appropriate training (in vivo/in vitro animal model) and proctorship should be considered prior to a clinical program of POEM.	Good practice	NA

GERD: gastroesophageal reflux disease; GOR: grade of recommendation; LOR: level of recommendation; NA: not applicable; POEM: peroral endoscopic myotomy

## Medicare National Coverage

There is no national coverage determination.

## Regulatory Status

POEM uses available laparoscopic instrumentation and, as a surgical procedure, is not subject to regulation by the U.S. Food and Drug Administration.

## References



1. Cheatham JG, Wong RK. Current approach to the treatment of achalasia. *Curr Gastroenterol Rep.* Jun 2011;13(3):219-225. PMID 21424734
2. Pandolfino JE, Kahrilas PJ. Presentation, diagnosis, and management of achalasia. *Clin Gastroenterol Hepatol.* Aug 2013;11(8):887-897. PMID 23395699
3. Yaghoobi M, Mayrand S, Martel M, et al. Laparoscopic Heller's myotomy versus pneumatic dilation in the treatment of idiopathic achalasia: a meta-analysis of randomized, controlled trials. *Gastrointest Endosc.* Sep 2013;78(3):468-475. PMID 23684149
4. Inoue H, Minami H, Kobayashi Y, et al. Peroral endoscopic myotomy (POEM) for esophageal achalasia. *Endoscopy.* Apr 2010;42(4):265-271. PMID 20354937
5. Hungness ES, Teitelbaum EN, Santos BF, et al. Comparison of perioperative outcomes between peroral esophageal myotomy (POEM) and laparoscopic Heller myotomy. *J Gastrointest Surg.* Feb 2013;17(2):228-235. PMID 23054897
6. Eckardt AJ, Eckardt VF. Treatment and surveillance strategies in achalasia: an update. *Nat Rev Gastroenterol Hepatol.* Jun 2011;8(6):311-319. PMID 21522116
7. Schlottmann F, Lockett DJ, Fine J, et al. Laparoscopic Heller myotomy versus peroral endoscopic myotomy (POEM) for achalasia: a systematic review and meta-analysis. *Ann Surg.* Mar 2018;267(3):451-460. PMID 28549006
8. Crespin OM, Liu LWC, Parmar A, et al. Safety and efficacy of POEM for treatment of achalasia: a systematic review of the literature. *Surg Endosc.* May 2017;31(5):2187-2201. PMID 27633440
9. Akintoye E, Kumar N, Obaitan I, et al. Peroral endoscopic myotomy: a meta-analysis. *Endoscopy.* Dec 2016;48(12):1059-1068. PMID 27617421
10. Patel K, Abbassi-Ghadi N, Markar S, et al. Peroral endoscopic myotomy for the treatment of esophageal achalasia: systematic review and pooled analysis. *Dis Esophagus.* Oct 2016;29(7):807-819. PMID 26175119
11. Marano L, Pallabazzer G, Solito B, et al. Surgery or peroral esophageal myotomy for achalasia: a systematic review and meta-analysis. *Medicine (Baltimore).* Mar 2016;95(10):e3001. PMID 26962813
12. Zhang Y, Wang H, Chen X, et al. Per-oral endoscopic myotomy versus laparoscopic Heller myotomy for achalasia: a meta-analysis of nonrandomized comparative studies. *Medicine (Baltimore).* Feb 2016;95(6):e2736. PMID 26871816
13. Sanaka MR, Hayat U, Thota PN, et al. Efficacy of peroral endoscopic myotomy vs other achalasia treatments in improving esophageal function. *World J Gastroenterol.* May 28 2016;22(20):4918-4925. PMID 27239118
14. Wang X, Tan Y, Lv L, et al. Peroral endoscopic myotomy versus pneumatic dilation for achalasia in patients aged  $\geq$  65 years. *Rev Esp Enferm Dig.* Oct 2016;108(10):637-641. PMID 27649684
15. Kumbhari V, Tieu AH, Onimaru M, et al. Peroral endoscopic myotomy (POEM) vs laparoscopic Heller myotomy (LHM) for the treatment of Type III achalasia in 75 patients: a multicenter comparative study. *Endosc Int Open.* Jun 2015;3(3):E195-201. PMID 26171430
16. Bhayani NH, Kurian AA, Dunst CM, et al. A comparative study on comprehensive, objective outcomes of laparoscopic Heller myotomy with per-oral endoscopic myotomy (POEM) for achalasia. *Ann Surg.* Jun 2014;259(6):1098-1103. PMID 24169175
17. Patti MG, Fisichella PM. Controversies in management of achalasia. *J Gastrointest Surg.* Sep 2014;18(9):1705-1709. PMID 24972973
18. Ujiki MB, Yetasook AK, Zapf M, et al. Peroral endoscopic myotomy: A short-term comparison with the standard laparoscopic approach. *Surgery.* Oct 2013;154(4):893-897; discussion 897-900. PMID 24074429
19. Hungness ES, Sternbach JM, Teitelbaum EN, et al. Per-oral endoscopic myotomy (POEM) after the learning curve: durable long-term results with a low complication rate. *Ann Surg.* Sep 2016;264(3):508-517. PMID 27513156





20. Ramchandani M, Nageshwar Reddy D, Darisetty S, et al. Peroral endoscopic myotomy for achalasia cardia: Treatment analysis and follow up of over 200 consecutive patients at a single center. *Dig Endosc.* Jan 2016;28(1):19-26. PMID 26018637
21. Inoue H, Sato H, Ikeda H, et al. Per-oral endoscopic myotomy: a series of 500 patients. *J Am Coll Surg.* Aug 2015;221(2):256-264. PMID 26206634
22. Teitelbaum EN, Soper NJ, Santos BF, et al. Symptomatic and physiologic outcomes one year after peroral esophageal myotomy (POEM) for treatment of achalasia. *Surg Endosc.* Dec 2014;28(12):3359-3365. PMID 24939164
23. Ling T, Guo H, Zou X. Effect of peroral endoscopic myotomy in achalasia patients with failure of prior pneumatic dilation: A prospective case-control study. *J Gastroenterol Hepatol.* Aug 2014;29(8):1609-1613. PMID 24628480
24. Ling TS, Guo HM, Yang T, et al. Effectiveness of peroral endoscopic myotomy in the treatment of achalasia: A pilot trial in Chinese Han population with a minimum of one-year follow-up. *J Dig Dis.* Jul 2014;15(7):352-358. PMID 24739072
25. Von Renteln D, Fuchs KH, Fockens P, et al. Peroral endoscopic myotomy for the treatment of achalasia: an international prospective multicenter study. *Gastroenterology.* Aug 2013;145(2):309-311 e303. PMID 23665071
26. Ren Z, Zhong Y, Zhou P, et al. Perioperative management and treatment for complications during and after peroral endoscopic myotomy (POEM) for esophageal achalasia (EA) (data from 119 cases). *Surg Endosc.* Nov 2012;26(11):3267-3272. PMID 22609984
27. Onimaru M, Inoue H, Ikeda H, et al. Peroral endoscopic myotomy is a viable option for failed surgical esophagocardiomyotomy instead of redo surgical Heller myotomy: a single center prospective study. *J Am Coll Surg.* Oct 2013;217(4):598-605. PMID 23891071
28. Zhou PH, Li QL, Yao LQ, et al. Peroral endoscopic remyotomy for failed Heller myotomy: a prospective single-center study. *Endoscopy.* Mar 2013;45(3):161-166. PMID 23389963
29. Li QL, Chen WF, Zhou PH, et al. Peroral endoscopic myotomy for the treatment of achalasia: a clinical comparative study of endoscopic full-thickness and circular muscle myotomy. *J Am Coll Surg.* Jul 25 2013;217(3):442-451. PMID 23891074
30. PJ, Katzka D, Richter JE. Clinical Practice Update: the use of per-oral endoscopic myotomy in achalasia: expert review and best practice advice from the AGA Institute. *Gastroenterology.* Nov 2017;153(5):1205-1211. PMID 28989059
31. Pasha SF, Acosta RD, Chandrasekhara V, et al. The role of endoscopy in the evaluation and management of dysphagia. *Gastrointest Endosc.* Feb 2014;79(2):191-201. PMID 24332405
32. Vaezi MF, Pandolfino JE, Vela MF. ACG clinical guideline: diagnosis and management of achalasia. *Am J Gastroenterol.* Aug 2013;108(8):1238-1249; quiz 1250. PMID 23877351
33. Stefanidis D, Richardson W, Farrell TM, et al. SAGES guidelines for the surgical treatment of esophageal achalasia. *Surg Endosc.* Feb 2012;26(2):296-311. PMID 22044977
34. Zaninotto G, Bennett C, Boeckstaens G, et al. The 2018 ISDE achalasia guidelines. *Dis Esophagus.* Aug 30 2018. PMID 30169645

## History

Date	Comments
11/11/13	New Policy. Policy created with literature search through August 1, 2013; considered investigational.



Date	Comments
11/20/14	Annual Review. Policy updated with literature review through August 18, 2014; references 3, 6-7, 9-12, and 18 added; no change to policy statement. ICD-9 and ICD-10 diagnosis codes removed; these do not relate to adjudication of this policy.
12/08/15	Annual Review. Policy updated with literature review through October 15, 2015; references 8-11 and 23 added. Policy statement unchanged.
12/01/16	Annual Review, approved November 8, 2016. Policy reviewed with literature search through September 2016; No change to policy statement
02/01/17	Annual Review, approved January 10, 2017. Policy updated with literature review through October 10, 2016; references 6-8, 10-11, and 15-16 added. Policy statement unchanged.
11/10/17	Policy moved to new format, no changes to policy statement.
02/01/18	Annual Review, approved January 9, 2018. Policy updated with literature review through September 14, 2017; reference 28 added. Policy statement unchanged.
02/01/19	Annual Review, approved January 4, 2019. Policy updated with literature review through September 2018; reference 9, 19, 30, and 34 added. Policy statement unchanged.

**Disclaimer:** This medical policy is a guide in evaluating the medical necessity of a particular service or treatment. The Company adopts policies after careful review of published peer-reviewed scientific literature, national guidelines and local standards of practice. Since medical technology is constantly changing, the Company reserves the right to review and update policies as appropriate. Member contracts differ in their benefits. Always consult the member benefit booklet or contact a member service representative to determine coverage for a specific medical service or supply. CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). ©2019 Premera All Rights Reserved.

**Scope:** Medical policies are systematically developed guidelines that serve as a resource for Company staff when determining coverage for specific medical procedures, drugs or devices. Coverage for medical services is subject to the limits and conditions of the member benefit plan. Members and their providers should consult the member benefit booklet or contact a customer service representative to determine whether there are any benefit limitations applicable to this service or supply. This medical policy does not apply to Medicare Advantage.



**Discrimination is Against the Law**

Premera Blue Cross complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex. Premera does not exclude people or treat them differently because of race, color, national origin, age, disability or sex.

Premera:

- Provides free aids and services to people with disabilities to communicate effectively with us, such as:
  - Qualified sign language interpreters
  - Written information in other formats (large print, audio, accessible electronic formats, other formats)
- Provides free language services to people whose primary language is not English, such as:
  - Qualified interpreters
  - Information written in other languages

If you need these services, contact the Civil Rights Coordinator.

If you believe that Premera has failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, or sex, you can file a grievance with:

Civil Rights Coordinator - Complaints and Appeals  
PO Box 91102, Seattle, WA 98111  
Toll free 855-332-4535, Fax 425-918-5592, TTY 800-842-5357  
Email AppealsDepartmentInquiries@Premera.com

You can file a grievance in person or by mail, fax, or email. If you need help filing a grievance, the Civil Rights Coordinator is available to help you.

You can also file a civil rights complaint with the U.S. Department of Health and Human Services, Office for Civil Rights, electronically through the Office for Civil Rights Complaint Portal, available at <https://ocrportal.hhs.gov/ocr/portal/lobby.jsf>, or by mail or phone at: U.S. Department of Health and Human Services  
200 Independence Avenue SW, Room 509F, HHH Building  
Washington, D.C. 20201, 1-800-368-1019, 800-537-7697 (TDD)  
Complaint forms are available at <http://www.hhs.gov/ocr/office/file/index.html>.

**Getting Help in Other Languages**

**This Notice has Important Information.** This notice may have important information about your application or coverage through Premera Blue Cross. There may be key dates in this notice. You may need to take action by certain deadlines to keep your health coverage or help with costs. You have the right to get this information and help in your language at no cost. Call 800-722-1471 (TTY: 800-842-5357).

**አማርኛ (Amharic):**

ይህ ማስታወቂያ አስፈላጊ መረጃ ይዟል። ይህ ማስታወቂያ ስለ ማመልከቻዎ ወይም የ Premera Blue Cross ሽፋን አስፈላጊ መረጃ ሊኖረው ይችላል። በዚህ ማስታወቂያ ውስጥ ቁልፍ ቀናት ሊኖሩ ይችላሉ። የጤና ሽፋንዎን ለመጠበቅና በአስፈላጊ እርዳታ ለማግኘት በተውሰኑ የጊዜ ገደቦች እርምጃ መውሰድ ይገባዎት ይሆናል። ይህን መረጃ እንዲያገኙ እና የለምንም ክፍያ በቋንቋዎ እርዳታ እንዲያገኙ መሰታወቅ አለዎት። በስልክ ቁጥር 800-722-1471 (TTY: 800-842-5357) ይደውሉ።

**العربية (Arabic):**

يحتوي هذا الإشعار على معلومات هامة. قد يحتوي هذا الإشعار على معلومات مهمة بخصوص طلبك أو التغطية التي تزيد الحصول عليها من خلال Premera Blue Cross. قد تكون هناك تواريخ مهمة في هذا الإشعار. وقد تحتاج لاتخاذ إجراء في تاريخ معينه للحفاظ على تغطيتك الصحية أو للمساعدة في دفع التكاليف. يحق لك الحصول على هذه المعلومات والمساعدة بلغتك دون تكبد أية تكلفة. اتصل بـ 800-722-1471 (TTY: 800-842-5357)

**中文 (Chinese):**

**本通知有重要的訊息。**本通知可能有關於您透過 Premera Blue Cross 提交的申請或保險的重要訊息。本通知內可能有重要日期。您可能需要在截止日期之前採取行動，以保留您的健康保險或者費用補貼。您有權利免費以您的母語得到本訊息和幫助。請撥電話 800-722-1471 (TTY: 800-842-5357)。

**Oromoo (Cushite):**

**Beeksisni kun odeeffannoo barbaachisaa qaba.** Beeksisni kun sagantaa yookan karaa Premera Blue Cross tiin tajaajila keessan ilaalchisee odeeffannoo barbaachisaa qabaachuu danda'a. Guyyaawwan murteessaa ta'an beeksisa kana keessatti ilaalaa. Tarii kaffaltiidhaan deeggaramuuf yookan tajaajila fayyaa keessaniif guyyaa dhumaa irratti wanti raawwattan jiraachuu danda'a. Kaffaltii irraa bilisa haala ta'een afaan keessaniin odeeffannoo argachuu fi deeggarsa argachuuf mirga ni qabaattu. Lakkoofsa bilbilaa 800-722-1471 (TTY: 800-842-5357) tii bilbilaa.

**Français (French):**

**Cet avis a d'importantes informations.** Cet avis peut avoir d'importantes informations sur votre demande ou la couverture par l'intermédiaire de Premera Blue Cross. Le présent avis peut contenir des dates clés. Vous devez peut-être prendre des mesures par certains délais pour maintenir votre couverture de santé ou d'aide avec les coûts. Vous avez le droit d'obtenir cette information et de l'aide dans votre langue à aucun coût. Appelez le 800-722-1471 (TTY: 800-842-5357).

**Kreyòl ayisyen (Creole):**

**Avi sila a gen Enfòmasyon Enpòtan ladann.** Avi sila a kapab genyen enfòmasyon enpòtan konsènan aplikasyon w lan oswa konsènan kouvèti asirans lan atravè Premera Blue Cross. Kapab genyen dat ki enpòtan nan avi sila a. Ou ka gen pou pran kèk aksyon avan sèten dat limit pou ka kenbe kouvèti asirans sante w la oswa pou yo ka ede w avèk depans yo. Se dwa w pou resewva enfòmasyon sa a ak asistans nan lang ou pale a, san ou pa gen pou peye pou sa. Rele nan 800-722-1471 (TTY: 800-842-5357).

**Deutsche (German):**

**Diese Benachrichtigung enthält wichtige Informationen.** Diese Benachrichtigung enthält unter Umständen wichtige Informationen bezüglich Ihres Antrags auf Krankenversicherungsschutz durch Premera Blue Cross. Suchen Sie nach eventuellen wichtigen Terminen in dieser Benachrichtigung. Sie könnten bis zu bestimmten Stichtagen handeln müssen, um Ihren Krankenversicherungsschutz oder Hilfe mit den Kosten zu behalten. Sie haben das Recht, kostenlose Hilfe und Informationen in Ihrer Sprache zu erhalten. Rufen Sie an unter 800-722-1471 (TTY: 800-842-5357).

**Hmoob (Hmong):**

**Tsab ntawv tshaj xo no muaj cov ntshiab lus tseem ceeb.** Tej zaum tsab ntawv tshaj xo no muaj cov ntshiab lus tseem ceeb txog koj daim ntawv thov kev pab los yog koj qhov kev pab cuam hns ntawm Premera Blue Cross. Tej zaum muaj cov hnuv tseem ceeb uas sau rau hauv daim ntawv no. Tej zaum koj kuj yuav tau ua qee yam uas peb kom koj ua tsis pub dhau cov caij nyuog uas teev tseg rau hauv daim ntawv no mas koj thiaj yuav tau txais kev pab cuam kho mob los yog kev pab them tej nqi kho mob ntawd. Koj muaj cai kom lawv muab cov ntshiab lus no uas tau muab sau ua koj hom lus pub dawb rau koj. Hu rau 800-722-1471 (TTY: 800-842-5357).

**Iloko (Ilocano):**

**Daytoy a Pakdaar ket naglaon iti Napateg nga Impormasion.** Daytoy a pakdaar mabalin nga adda ket naglaon iti napateg nga impormasion maipanggep iti aplikasyonyo wenna coverage babaen iti Premera Blue Cross. Daytoy ket mabalin dagiti importante a petsa iti daytoy a pakdaar. Mabalin nga adda rumbeng nga aramidenyo nga addang sakbay dagiti partikular a naituding nga aldaw tapno mapagtalinaedyo ti coverage ti salun-atyto wenna tulong kadagiti gastos. Adda karbenganyo a mangala iti daytoy nga impormasion ken tulong iti bukodyo a pagsasao nga awan ti bayadanyo. Tumawag iti numero nga 800-722-1471 (TTY: 800-842-5357).

**Italiano (Italian):**

**Questo avviso contiene informazioni importanti.** Questo avviso può contenere informazioni importanti sulla tua domanda o copertura attraverso Premera Blue Cross. Potrebbero esserci date chiave in questo avviso. Potrebbe essere necessario un tuo intervento entro una scadenza determinata per consentirti di mantenere la tua copertura o sovvenzione. Hai il diritto di ottenere queste informazioni e assistenza nella tua lingua gratuitamente. Chiama 800-722-1471 (TTY: 800-842-5357).

**日本語 (Japanese):**

この通知には重要な情報が含まれています。この通知には、Premera Blue Cross の申請または補償範囲に関する重要な情報が含まれている場合があります。この通知に記載されている可能性がある重要な日付をご確認ください。健康保険や有料サポートを維持するには、特定の期日までに行動を取らなければならない場合があります。ご希望の言語による情報とサポートが無料で提供されます。800-722-1471 (TTY: 800-842-5357)までお電話ください。

**한국어 (Korean):**

본 통지서에는 중요한 정보가 들어 있습니다. 즉 이 통지서는 귀하의 신청에 관하여 그리고 Premera Blue Cross 를 통한 커버리지에 관한 정보를 포함하고 있을 수 있습니다. 본 통지서에는 핵심이 되는 날짜들이 있을 수 있습니다. 귀하의 귀하의 건강 커버리지를 계속 유지하거나 비용을 절감하기 위해서 일정한 마감일까지 조치를 취해야 할 필요가 있을 수 있습니다. 귀하의 이러한 정보와 도움을 귀하의 언어로 비용 부담없이 얻을 수 있는 권리가 있습니다. 800-722-1471 (TTY: 800-842-5357) 로 전화하십시오.

**ລາວ (Lao):**

ແຈງການນີ້ມີຂໍ້ມູນສໍາຄັນ. ແຈງການນີ້ອາດຈະມີຂໍ້ມູນສໍາຄັນກ່ຽວກັບຄໍາຮ້ອງສະໝັກ ຫຼື ຄວາມຄົມຄອງປະກັນໄພຂອງທ່ານຜ່ານ Premera Blue Cross. ອາດຈະມີວັນທີ່ສໍາຄັນໃນແຈງການນີ້. ທ່ານອາດຈະຈໍາເປັນຕ້ອງດໍາເນີນການຕາມກຳນົດ ເວລາສະເພາະເພື່ອຮັກສາຄວາມຄົມຄອງປະກັນສະພາບ ຫຼື ຄວາມຊ່ວຍເຫຼືອເວັ້ນເວົ້ອງຄ່າໃຊ້ຈ່າຍຂອງທ່ານໄວ້. ທ່ານມີສິດໄດ້ຮັບຂໍ້ມູນນີ້ ແລະ ຄວາມຊ່ວຍເຫຼືອເປັນພາສາຂອງທ່ານໂດຍບໍ່ເສຍຄ່າ. ໃຫ້ໃບທາ 800-722-1471 (TTY: 800-842-5357).

**ភាសាខ្មែរ (Khmer):**

សេចក្តីជូនដំណឹងនេះមានព័ត៌មានយ៉ាងសំខាន់។ សេចក្តីជូនដំណឹងនេះប្រហែលជាមានព័ត៌មានយ៉ាងសំខាន់អំពីទម្រង់បែបបទ ឬការរៀបចំរបស់អ្នកកាមរយ: Premera Blue Cross ។ ប្រហែលជាមាន កាលបរិច្ឆេទសំខាន់នៅក្នុងសេចក្តីជូនដំណឹងនេះ។ អ្នកប្រហែលជាត្រូវការបញ្ជាក់សមត្ថភាព ដល់កិច្ចការផ្ទៃក្នុងរបស់នានា ដើម្បីនឹងរក្សាទុកការធានារ៉ាប់រងអនាគតរបស់អ្នក ឬប្រាក់ដុល្លារចេញផ្លូវ។ អ្នកមានសិទ្ធិទទួលបានព័ត៌មាននេះ និងដុល្លារនៅក្នុងភាសារបស់អ្នកដោយមិនអស់លុយឡើយ។ សូមទូរស័ព្ទ 800-722-1471 (TTY: 800-842-5357)។

**ਪੰਜਾਬੀ (Punjabi):**

ਇਸ ਨੋਟਿਸ ਵਿਚ ਖਾਸ ਜਾਣਕਾਰੀ ਹੈ. ਇਸ ਨੋਟਿਸ ਵਿਚ Premera Blue Cross ਵਲੋਂ ਤੁਹਾਡੀ ਕਵਰੇਜ ਅਤੇ ਅਰਜੀ ਬਾਰੇ ਮਹੱਤਵਪੂਰਨ ਜਾਣਕਾਰੀ ਹੋ ਸਕਦੀ ਹੈ . ਇਸ ਨੋਟਿਸ ਨਵ ਖਾਸ ਤਾਰੀਖਾਂ ਹੋ ਸਕਦੀਆਂ ਹਨ. ਜੇਕਰ ਤੁਸੀਂ ਜਸਰਤ ਕਵਰੇਜ ਰਿੱਖਣੀ ਹੋਵੇ ਜਾਂ ਓਸ ਦੀ ਲਾਗਤ ਜਵਿੱਚ ਮਦਦ ਦੇ ਇਛੁੱਕ ਹੋ ਤਾਂ ਤੁਹਾਨੂੰ ਅੰਤਮ ਤਾਰੀਖ ਤੋਂ ਪਹਿਲਾਂ ਢੁੱਝ ਖਾਸ ਕਦਮ ਚੁੱਕਣ ਦੀ ਲੋੜ ਹੋ ਸਕਦੀ ਹੈ ,ਤੁਹਾਨੂੰ ਮੁਫਤ ਵਿੱਚ ਤੋਂ ਅਪਣੀ ਭਾਸ਼ਾ ਵਿੱਚ ਜਾਣਕਾਰੀ ਅਤੇ ਮਦਦ ਪ੍ਰਾਪਤ ਕਰਨ ਦਾ ਅਧਿਕਾਰ ਹੈ ,ਕਾਲ 800-722-1471 (TTY: 800-842-5357).

**فارسی (Farsi):**

این اعلامیه حاوی اطلاعات مهم میباشد. این اعلامیه ممکن است حاوی اطلاعات مهم درباره فرم تقاضا و یا پوشش بیمه ای شما از طریق Premera Blue Cross باشد. به تاریخ های مهم در این اعلامیه توجه نمایید. شما ممکن است برای حفظ پوشش بیمه تان یا کمک در پرداخت هزینه های درمانی تان، به تاریخ های مشخصی برای انجام کارهای خاصی احتیاج داشته باشید. شما حق این را دارید که این اطلاعات و کمک را به زبان خود به طور رایگان دریافت نمایید. برای کسب اطلاعات با شماره 800-722-1471 (کلیربران TTY تماس باشماره 800-842-5357) تماس برقرار نمایید.

**Polskie (Polish):**

To ogłoszenie może zawierać ważne informacje. To ogłoszenie może zawierać ważne informacje odnośnie Państwa wniosku lub zakresu świadczeń poprzez Premera Blue Cross. Prosimy zwrócić uwagę na kluczowe daty, które mogą być zawarte w tym ogłoszeniu aby nie przekroczyć terminów w przypadku utrzymania polisy ubezpieczeniowej lub pomocy związanej z kosztami. Macie Państwo prawo do bezpłatnej informacji we własnym języku. Zadzwońcie pod 800-722-1471 (TTY: 800-842-5357).

**Português (Portuguese):**

Este aviso contém informações importantes. Este aviso poderá conter informações importantes a respeito de sua aplicação ou cobertura por meio do Premera Blue Cross. Poderão existir datas importantes neste aviso. Talvez seja necessário que você tome providências dentro de determinados prazos para manter sua cobertura de saúde ou ajuda de custos. Você tem o direito de obter esta informação e ajuda em seu idioma e sem custos. Ligue para 800-722-1471 (TTY: 800-842-5357).

**Română (Romanian):**

Prezenta notificare conține informații importante privind cererea sau acoperirea asigurării dumneavoastră de sănătate prin Premera Blue Cross. Pot exista date cheie în această notificare. Este posibil să fie nevoie să acționați până la anumite termene limită pentru a vă menține acoperirea asigurării de sănătate sau asistența provizorie la costuri. Aveți dreptul de a obține gratuit aceste informații și ajutor în limba dumneavoastră. Sunați la 800-722-1471 (TTY: 800-842-5357).

**Русский (Russian):**

Настоящее уведомление содержит важную информацию. Это уведомление может содержать важную информацию о вашем заявлении или страховом покрытии через Premera Blue Cross. В настоящем уведомлении могут быть указаны ключевые даты. Вам, возможно, потребуется принять меры к определенным предельным срокам для сохранения страхового покрытия или помощи с расходами. Вы имеете право на бесплатное получение этой информации и помощь на вашем языке. Звоните по телефону 800-722-1471 (TTY: 800-842-5357).

**Fa'asamoa (Samoan):**

Atonu ua iai i lenei fa'asilasilaga ni fa'amatalaga e sili ona taua e tatau ona e malamalama i ai. O lenei fa'asilasilaga o se fesoasoani e fa'amatala atili i ai i le tulaga o le polokalame, Premera Blue Cross, ua e tau fia maua atu i ai. Fa'amolemole, ia e iloilo fa'alelei i aso fa'apitoa olo'o iai i lenei fa'asilasilaga taua. Masalo o le'a iai ni feau e tatau ona e faia ao le'i aulia le aso ua ta'ua i lenei fa'asilasilaga ina ia e iai pea ma maua fesoasoani mai ai i le polokalame a le Malo olo'o e iai i ai. Olo'o iai iate oe le aia tatau e maua atu i lenei fa'asilasilaga ma lenei fa'matalaga i legagana e te malamalama i ai aunoa ma se togiga tupe. Vili atu i le telefoni 800-722-1471 (TTY: 800-842-5357).

**Español (Spanish):**

Este Aviso contiene información importante. Es posible que este aviso contenga información importante acerca de su solicitud o cobertura a través de Premera Blue Cross. Es posible que haya fechas clave en este aviso. Es posible que deba tomar alguna medida antes de determinadas fechas para mantener su cobertura médica o ayuda con los costos. Usted tiene derecho a recibir esta información y ayuda en su idioma sin costo alguno. Llame al 800-722-1471 (TTY: 800-842-5357).

**Tagalog (Tagalog):**

Ang Paunawa na ito ay naglalaman ng mahalagang impormasyon tungkol sa iyong aplikasyon o pagsakop sa pamamagitan ng Premera Blue Cross. Maaaring may mga mahalagang petsa dito sa paunawa. Maaring mangailangan ka na magsagawa ng hakbang sa ilang mga itinakdang panahon upang mapanatili ang iyong pagsakop sa kalusugan o tulong na walang gastos. May karapatan ka na makakuha ng ganiitong impormasyon at tulong sa iyong wika ng walang gastos. Tumawag sa 800-722-1471 (TTY: 800-842-5357).

**ไทย (Thai):**

ประกาศนี้มีข้อมูลสำคัญ ประกาศนี้อาจมีข้อมูลที่สำคัญเกี่ยวกับกาการสมัครหรือขอบเขตประกันสุขภาพของคุณผ่าน Premera Blue Cross และอาจมีกำหนดการในประกาศนี้ คุณอาจจะต้องดำเนินการภายในกำหนดระยะเวลาที่แน่นอนเพื่อจะรักษาการประกันสุขภาพของคุณหรือการช่วยเหลือที่มีค่าใช้จ่าย คุณมีสิทธิที่จะได้รับข้อมูลและความช่วยเหลือนี้ในภาษาของคุณโดยไม่มีค่าใช้จ่าย โทร 800-722-1471 (TTY: 800-842-5357)

**Український (Ukrainian):**

Це повідомлення містить важливу інформацію. Це повідомлення може містити важливу інформацію про Ваше звернення щодо страховального покриття через Premera Blue Cross. Зверніть увагу на ключові дати, які можуть бути вказані у цьому повідомленні. Існує імовірність того, що Вам треба буде здійснити певні кроки у конкретні кінцеві строки для того, щоб зберегти Ваше медичне страхування або отримати фінансову допомогу. У Вас є право на отримання цієї інформації та допомоги безкоштовно на Вашій рідній мові. Дзвоніть за номером телефону 800-722-1471 (TTY: 800-842-5357).

**Tiếng Việt (Vietnamese):**

Thông báo này cung cấp thông tin quan trọng. Thông báo này có thông tin quan trọng về đơn xin tham gia hoặc hợp đồng bảo hiểm của quý vị qua chương trình Premera Blue Cross. Xin xem ngày quan trọng trong thông báo này. Quý vị có thể phải thực hiện theo thông báo đúng trong thời hạn để duy trì bảo hiểm sức khỏe hoặc được trợ giúp thêm về chi phí. Quý vị có quyền được biết thông tin này và được trợ giúp bằng ngôn ngữ của mình miễn phí. Xin gọi số 800-722-1471 (TTY: 800-842-5357).