

MEDICAL POLICY – 2.02.09

Closure Devices for Patent Foramen Ovale and Atrial Septal Defects

BCBSA Ref. Policy: 2.02.09

Effective Date: Aug. 1, 2020

Last Revised: Jul. 2, 2020


Replaces: N/A

RELATED MEDICAL POLICIES:

None

Select a hyperlink below to be directed to that section.

[POLICY CRITERIA](#) | [DOCUMENTATION REQUIREMENTS](#) | [CODING](#)
[RELATED INFORMATION](#) | [EVIDENCE REVIEW](#) | [REFERENCES](#) | [HISTORY](#)

 Clicking this icon returns you to the hyperlinks menu above.

Introduction

The heart's two upper chambers are called atria. The septum is the thin wall of tissue that separates the atria. Sometimes there may be a congenital heart defect in which there is a hole in the septum between the two atria. This is called an atrial septal defect. Atrial septal defects may not cause any problems and might not even be diagnosed until adulthood. Larger atrial septal defects may cause problems and may need to be closed.

One way to treat an atrial septal defect is to use a catheter. A catheter is a long, thin tube which is threaded through a blood vessel in the groin to the heart. Once the catheter is in the correct location, a small device is put in place to seal the opening between the atria. This policy discusses when a catheter may be considered medically necessary to treat atrial septal defects.

The foramen ovale is an opening in the septum between the two atria that is normally found in a baby before it is born. This opening usually closes soon after birth. If a foramen ovale doesn't automatically close after the baby is born, it is called a patent foramen ovale (PFO). For most people a patent foramen ovale does not cause problems. In the small subset of people who have had a stroke where the cause is uncertain, using a new device to close the PFO may decrease the risk of a second stroke.

Note: The Introduction section is for your general knowledge and is not to be taken as policy coverage criteria. The rest of the policy uses specific words and concepts familiar to medical professionals. It is intended for providers. A provider can be a person, such as a doctor, nurse, psychologist, or dentist. A provider also can be a place where medical care is given, like a hospital, clinic, or lab. This policy informs them about when a service may be covered.

Policy Coverage Criteria

FDA approved devices	Medical Necessity
<p>Atrial septal defects (ASD) closure devices:</p> <ul style="list-style-type: none"> • Amplatzer™ Septal Occluder • GORE CARDIOFORM Septal Occluder 	<p>Transcatheter closure of secundum atrial septal defects may be considered medically necessary when using a device that has been approved by the U.S. Food and Drug Administration for that purpose and used according to the label indications including:</p> <ul style="list-style-type: none"> • Patients with echocardiographic evidence of ostium secundum atrial septal defect; <p>AND either ONE of the following:</p> <ul style="list-style-type: none"> • Clinical evidence of right ventricular volume overload (ie, 1.5:1 degree of left-to-right shunt or right ventricular enlargement); <p>OR</p> <ul style="list-style-type: none"> • Clinical evidence of paradoxical embolism is present <p>Transcatheter closure of secundum atrial septal defects is considered investigational for all other indications not meeting criteria outlined above.</p>
<p>Patent foramen ovale (PFO) closure devices:</p> <ul style="list-style-type: none"> • Amplatzer™ PFO Occluder • GORE CARDIOFORM Septal Occluder 	<p>The percutaneous transcatheter closure of a patent foramen ovale using a device that has been approved by the U.S. Food and Drug Administration for that purpose may be considered medically necessary to reduce the risk of recurrent ischemic stroke if patient meets all of the following:</p> <ul style="list-style-type: none"> • The patient is between the ages of 18 and 60 years of age • Diagnosed with patent foramen ovale with a right-to-left interatrial shunt confirmed by echocardiography with at least ONE of the following characteristics: <ul style="list-style-type: none"> ○ PFO with large shunt (defined as >30 microbubbles in the left atrium within 3 cardiac cycles, after opacification of the



FDA approved devices	Medical Necessity
	<p>right atrium); OR</p> <ul style="list-style-type: none"> ○ PFO associated with atrial septal aneurysm on transesophageal examination (septum primum excursion >10 mm) • Documented history of a cryptogenic stroke due to a presumed paradoxical embolism, as determined by: <ul style="list-style-type: none"> ○ A neurologist and cardiologist following an evaluation to exclude any other identifiable cause of stroke, including large vessel atherosclerotic disease and small vessel occlusive disease <p>AND</p> <ul style="list-style-type: none"> • None of the following are present: <ul style="list-style-type: none"> ○ Uncontrolled vascular risk factors, including uncontrolled diabetes or uncontrolled hypertension ○ Other sources of right-to-left shunts, including an atrial septal defect and/or fenestrated septum ○ Active endocarditis or other untreated infections ○ Inferior vena cava filter

Documentation Requirements
<p>The patient's medical records submitted for review should document that medical necessity criteria are met. The record should include clinical documentation of:</p> <ul style="list-style-type: none"> • Diagnosis/condition • History and physical examination documenting the severity of the condition • Results and/or reports from prior imaging or testing completed • Any prior procedures • Name of device to be used for closure

Coding

Code	Description
CPT	
93580	Percutaneous transcatheter closure of congenital interatrial communication (ie, Fontan



Code	Description
CPT	
	fenestration, atrial septal defect) with implant

Note: CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). HCPCS codes, descriptions and materials are copyrighted by Centers for Medicare Services (CMS).

Related Information

Definition of Terms

Atrial septal aneurysm: Is a rare, but well-recognized cardiac abnormality in which there is an overabundant or weakened septal tissue allowing the septum to become very mobile and protrude or bulge into the right or left atrium of at least 10 mm to 15 mm. This can be visualized at echocardiography and the degree of deviation can be measured. Atrial septal aneurysms are often associated with larger PFOs.

Cryptogenic stroke: A stroke that happens for an unknown reason after other causes such as cardiac, pulmonary, vascular or neurologic sources have been ruled out.

Ischemic stroke: A stroke that happens when a blood vessel that carries blood to the brain is blocked either due to arteries narrowed by atherosclerosis or a blood clot (thrombus).

Paradoxical embolism (PDE): This happens when a clot (thrombus) passes through the patent foramen ovale (PFO) in the heart, bypassing the lungs that act as a clot filter.

Septum primum: A septum in the embryonic heart, dividing the primitive atrium into right and left chambers from Latin, meaning 'first septum'.

Evidence Review

Description

Patent foramen ovale (PFO) and atrial septal defects (ASDs) are relatively common congenital heart defects that can be associated with a range of symptoms. PFOs may be asymptomatic but have been associated with higher rates of cryptogenic stroke. PFOs have also been investigated



in association with a variety of other conditions, such as a migraine. Depending on their size, ASDs may lead to left-to-right shunting and signs and symptoms of pulmonary overload. Repair of ASDs is indicated for patients with a significant degree of left-to-right shunting. Transcatheter closure devices have been developed to repair PFO and ASDs. These devices are alternatives to open surgical repair for ASDs or treatment with antiplatelet and/or anticoagulant medications in patients with cryptogenic stroke and PFO.

Background

Patent Foramen Ovale

The foramen ovale, a component of fetal cardiovascular circulation, consists of a communication between the right and left atrium that functions as a vascular bypass of the uninflated lungs. The ductus arteriosus is another feature of the fetal cardiovascular circulation, consisting of a connection between the pulmonary artery and the distal aorta. Before birth, the foramen ovale is held open by the large flow of blood into the left atrium from the inferior vena cava. Over the course of months after birth, an increase in left atrial pressure and a decrease in right atrial pressure result in permanent closure of the foramen ovale in most individuals. However, a patent foramen ovale (PFO) is a common finding in 25% of asymptomatic adults.¹ In some epidemiologic studies, PFO has been associated with cryptogenic stroke, defined as an ischemic stroke occurring in the absence of potential cardiac, pulmonary, vascular, or neurologic sources. Studies have also shown an association between PFO and migraine headache.

Atrial Septal Defects

Unlike PFO, which represents the postnatal persistence of normal fetal cardiovascular physiology, atrial septal defects (ASDs) represent an abnormality in the development of the heart that results in free communication between the atria. ASDs are categorized by their anatomy. Ostium secundum describes defects located midseptally and are typically near the fossa ovalis. Ostium primum defects lie immediately adjacent to the atrioventricular valves and are within the spectrum of atrioventricular septal defects. Primum defects occur commonly in patients with Down syndrome. Sinus venous defects occur high in the atrial septum and are frequently associated with anomalies of the pulmonary veins.

Ostium secundum ASDs are the third most common form of congenital heart disorder and among the most common congenital cardiac malformations in adults, accounting for 30% to



40% of these patients older than age 40 years. The ASD often goes unnoticed for decades because the physical signs are subtle and the clinical sequelae are mild. However, virtually all patients who survive into their sixth decade are symptomatic; fewer than 50% of patients survive beyond age 40 to 50 years due to heart failure or pulmonary hypertension related to the left-to-right shunt. Symptoms related to ASD depend on the size of the defect and the relative diastolic filling properties of the left and right ventricles. Reduced left ventricular compliance, and mitral stenosis will increase left-to-right shunting across the defect. Conditions that reduce right ventricular compliance and tricuspid stenosis will reduce left-to-right shunting or cause a right-to-left shunt. Symptoms of an ASD include exercise intolerance and dyspnea, atrial fibrillation, and less commonly, signs of right heart failure. Patients with ASDs are also at risk for paradoxical emboli.

Treatment

Repair of ASDs is recommended for those with a pulmonary-to-systemic flow ratio ($Q_p: Q_s$) exceeding 1.5:1.0. Despite the success of surgical repair, there has been interest in developing a transcatheter-based approach to ASD repair to avoid the risks and morbidity of open heart surgery. A variety of devices have been researched. Technical challenges include minimizing the size of the device so that smaller catheters can be used, developing techniques to center the device properly across the ASD, and ensuring that the device can be easily retrieved or repositioned, if necessary.

Individuals with ASDs and a history of cryptogenic stroke are typically treated with antiplatelet agents, given an absence of evidence that systemic anticoagulation is associated with outcome improvements.

Transcatheter Closure Devices

Transcatheter PFO and ASD occluders typically consist of a single or paired wire mesh discs covered or filled with polyester or polymer fabric that are placed over the septal defect. Over time, the occlusion system is epithelialized. ASD occluder devices consist of flexible mesh discs delivered via catheter to cover the ASD.



Summary of Evidence

For individuals who have PFO and cryptogenic stroke who receive PFO closure with a transcatheter device, the evidence includes multiple randomized controlled trials (RCTs) comparing device-based PFO closure with medical therapy, systematic reviews, meta-analyses, and observational studies. Relevant outcomes are symptoms, change in disease status, overall survival, morbid events, and treatment-related morbidity and mortality. The RCTs comparing PFO closure with medical management have suggested that PFO closure is more effective than medical therapy in reducing event rates. Although these results were not statistically significant by intention-to-treat analyses in the first 3 trials [ie, Evaluation of the STARFlex Septal Closure System in Patients with a Stroke and/or Transient Ischemic Attack due to Presumed Paradoxical Embolism through a Patent Foramen Ovale (CLOSURE), Amplatzer PFO Occluder with Medical Treatment in Patients with Cryptogenic Embolism (PC-Trial), and Randomized Evaluation of Recurrent Stroke Comparing PFO Closure to Established Current Standard of Care Treatment (RESPECT; initial study)], they were statistically significant in later trials [ie, RESPECT (extended follow-up), Reduction in the Use of Corticosteroids in Exacerbated COPD (REDUCE), and Patent Foramen Ovale Closure or Anticoagulants versus Antiplatelet Therapy to Prevent Stroke Recurrence (CLOSE)]. Use of appropriate patient selection criteria to eliminate other causes of cryptogenic stroke in RESPECT, REDUCE, and CLOSE trials contributed to findings of the superiority of PFO closure compared with medical management. Of note, higher rates of atrial fibrillation were reported in a few of the individual trials and in the meta-analysis that incorporated evidence from RESPECT, REDUCE, and CLOSE trials. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome, and observational studies.

For individuals who have PFO and migraines who receive PFO closure with a transcatheter device, the evidence includes two RCTs of PFO closure and multiple observational studies reporting on the association between PFO and migraine. Relevant outcomes are symptoms, quality of life, medication use, and treatment-related morbidity and mortality. The available sham-controlled randomized trial did not demonstrate significant improvements in migraine symptoms after PFO closure. A second RCT with blinded end point evaluation did not demonstrate reductions in migraine days after PFO closure but likely was underpowered. Nonrandomized studies have shown highly variable rates of migraine reduction after PFO closure. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have PFO and conditions associated with PFO other than cryptogenic stroke or migraine (eg, platypnea-orthodeoxia syndrome, myocardial infarction with normal coronary arteries, decompression illness, high-altitude pulmonary edema, obstructive sleep apnea) who



receive PFO closure with a transcatheter device, the evidence includes small case series and case reports. Relevant outcomes are symptoms, change in disease status, morbid events, and treatment-related morbidity and mortality. Comparative studies are needed to evaluate outcomes in similar patient groups treated with and without PFO closure. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have ASD and evidence of left-to-right shunt or right ventricular overload who receive ASD closure with a transcatheter device, the evidence includes nonrandomized comparative studies and single-arm studies. Relevant outcomes are symptoms, change in disease status, and treatment-related morbidity and mortality. The available nonrandomized comparative studies and single-arm case series have shown rates of closure using transcatheter-based devices approaching the high success rates of surgery, which are supported by meta-analyses of these studies. The percutaneous approach has a low complication rate and avoids the morbidity and complications of open surgery. If the percutaneous approach is unsuccessful, ASD closure can be achieved using surgery. Because of the benefits of percutaneous closure over open surgery, it can be determined that transcatheter ASD closure improves outcomes in patients with an indication for ASD closure. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in [Table 1](#).

Table 1. Summary of Key Clinical Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
Ongoing			
NCT00738894^a	GORE® HELEX® Septal Occluder / GORE® Septal Occluder and Antiplatelet Medical Management for Reduction of Recurrent Stroke or Imaging-Confirmed TIA in Patients With Patent Foramen Ovale (PFO)	664	Feb 2020
NCT01550588	Device Closure Versus Medical Therapy for Secondary Prevention in Cryptogenic Stroke Patients With High-Risk Patent Foramen Ovale : DEFENSE-PFO	210	Feb 2020
NCT03867708	Outcomes of Transcatheter Closure of Secundum Atrial Septal Defect Guided by Three-dimensional	80	June 2021



NCT No.	Trial Name	Planned Enrollment	Completion Date
	Transesophageal Echocardiography		
NCT03309332	OBS Lead-AMPLATZER PFO Occluder New Enrollment Study	1214	Dec 2027
Unpublished			
NCT01960491	Prospective Single Center Pilot Clinical Study to Evaluate the Safety and Effectiveness of an Intracardiac Septal Closure Device With Biodegradable Framework in Patients With Clinically Significant Atrial Septum Defect (ASD) or Patent Foramen Ovale (PFO)	15	June 2018 <i>(updated 10/11/18)</i>

NCT: national clinical trial

^a Denotes industry-sponsored or cosponsored trial

Clinical Input Received from Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

In response to requests, input was received from 2 academic medical centers (1 of which provided 2 responses) while this policy was under review in 2016. Input was mixed about the medical necessity of closure devices for patent foramen ovale (PFO) in patients with cryptogenic stroke or transient ischemic attack due to presumed paradoxical embolism through the PFO. There was consensus that use of closure devices for PFO in patients with other conditions (eg, migraine, platypnea-orthodeoxia syndrome) is not medically necessary.

Practice Guidelines and Position Statements

The American College of Chest Physicians

In 2012, the American College of Chest Physicians updated its guidelines on antithrombotic therapy and the prevention of thrombosis, which made the following recommendations related to patent foramen ovale (PFO) and cryptogenic stroke⁴⁶:



We suggest that patients with stroke and PFO are treated with antiplatelet therapy following the recommendations for patients with noncardioembolic stroke.... In patients with a history of noncardioembolic ischemic stroke or TIA, we recommend long-term treatment with aspirin (75-100 mg once daily), clopidogrel (75 mg once daily), aspirin/extended release dipyridamole (25 mg/200 mg bid), or cilostazol (100 mg bid) over no antiplatelet therapy (Grade 1A), oral anticoagulants (Grade 1B), the combination of clopidogrel plus aspirin (Grade 1B), or triflusal (Grade 2B).

The American Academy of Neurology

In 2016, the American Academy of Neurology updated its evidence-based guidelines on the management of patients with stroke and PFO to address whether percutaneous closure of PFO is superior to medical therapy alone.⁴⁷ Following a systematic review of the literature and structured formulation of recommendations, the Academy developed conclusions for the Amplatzer PFO Occluder devices. For patients with cryptogenic stroke and PFO, percutaneous PFO closure with the Amplatzer PFO Occluder:

- "Possibly decreases the risk of recurrent stroke—RD [risk difference] -1.68%, 95% CI [confidence interval]-3.18% to -0.19%;"
- "Possibly increases the risk of new-onset AF [atrial fibrillation] —RD 1.64%, 95% CI 0.07%–3.2% (2 Class I studies; confidence downgraded to low for risk of bias relative to magnitude of effect and imprecision);"
- "Is highly likely to be associated with a procedural complication risk of 3.4%, 95% CI 2.3%–5% (2 Class I studies)."

The guidelines concluded:

Clinicians should not routinely offer percutaneous PFO closure to patients with cryptogenic ischemic stroke outside of a research setting (Level R). In rare circumstances, such as recurrent strokes despite adequate medical therapy with no other mechanism identified, clinicians may offer the AMPLATZER PFO Occluder if it is available (Level C).



American Heart Association and American Stroke Association

In 2014, the American Heart Association and American Stroke Association updated its guidelines on the prevention of stroke in patients with ischemic stroke or transient ischemic attack. The guidelines made the following recommendations for device-based closure for PFO⁴⁸:

- For patients with a cryptogenic ischemic stroke or TIA [transient ischemic attack] and a PFO without evidence for DVT [deep vein thrombosis], available data do not support a benefit for PFO closure (Class III; Level of Evidence A).
- In the setting of PFO and DVT, PFO closure by a transcatheter device might be considered, depending on the risk of recurrent DVT (Class IIb; Level of Evidence C).

American College of Cardiology and American Heart Association

In 2018, the American College of Cardiology and American Heart Association updated guidelines on the management of adults with congenital heart disease.⁴⁹ The treatment recommendations are summarized in Table 10., coronary sinus, or primum atrial septal defect, however, surgery rather than percutaneous closure was recommended.

Table 2. American College of Cardiology and American Heart Association Recommendations for Treating Atrial Septal Defect

Condition	Recommendation	COR ^a /LOE ^b
Symptomatic isolated secundum ASD, right atrial and/or RV enlargement, and net left-to-right shunt sufficiency large enough to cause physiological sequelae, without cyanosis at rest or during exercise	Transcatheter or surgical closure	I1/B-NR2
Symptomatic primum ASD, sinus venosus defect, or coronary sinus defect, right atrial and/or RV enlargement, and net left-to-right shunt sufficiency large enough to cause physiological sequelae, without cyanosis at rest or during exercise	Surgical closure unless precluded by comorbidities	I1/B-NR2
Asymptomatic isolated secundum ASD, right atrial and RV enlargement, and net left-to-right shunt sufficiency large enough to cause physiological sequelae, without cyanosis at rest or during	Transcatheter or surgical closure	I1a1/C-LD2



exercise		
Secundum ASD when a concomitant surgical procedure is being performed and there is a net left-to-right shunt sufficiently large enough to cause physiological sequelae, and right atrial and RV enlargement without cyanosis at rest or during exercise	Surgical closure	IIa1/C-LD2
ASD when net left-to-right shunt is $\geq 1.5:1$, PA systolic pressure and/or pulmonary vascular resistance is greater than of one-third of systemic resistance	Percutaneous or surgical closure	IIb1/B-NR2
ASD with PA systolic pressure greater than two-thirds systemic, pulmonary vascular resistance greater than two-thirds systemic, and/or a net left-to-right shunt	ASD closure should not be performed	III-Harm1/C-LD2

Adapted from Stout et al (2019)⁴⁹.

ASD: atrial septal defect; COR: class (strength) of recommendation; LOE: level (quality) of evidence; PA: pulmonary artery; RCT: randomized controlled trial; RV: right ventricular.

^a COR key: I=strong; IIa=moderate; IIb=weak; III: No Benefit=weak; III: Harm=strong.⁴⁹

^b LOE key: A=high quality from >1 RCT, meta-analyses of high-quality RCTs, ≥ 1 RCT corroborated by high-quality registry studies; B-R=randomized, moderate-quality evidence from ≥ 1 RCT or meta-analysis of moderate-quality RCTs; B-NR=nonrandomized, moderate-quality evidence from ≥ 1 well-designed, well-executed nonrandomized study, observational study, or registry study, or meta-analyses of such studies; C-LD: limited data, randomized or nonrandomized observational or registry studies with limitations of design or execution, meta-analyses of such studies, or physiological or mechanistic studies in human subjects; C-EO: expert opinion.⁴⁹

Medicare National Coverage

There is no national coverage determination.

Regulatory Status

Patent Foramen Ovale (PFO) Closure Devices

The U.S. Food and Drug Administration (FDA) has approved three devices for ASD closure through the premarket approval process or a premarket approval supplement: the Amplatzer Septal Occluder, the GORE HELEX Septal Occluder (discontinued), and the GORE CARDIOFORM Septal Occluder (see [Table 3](#)) (FDA product code: MLV).

In 2002, two transcatheter devices were cleared for marketing by the U.S. Food and Drug Administration FDA through a humanitarian device exemption as treatment for patients with cryptogenic stroke and PFO: the CardioSEAL® Septal Occlusion System (NMT Medical; device



no longer commercially available) and the Amplatzer® PFO Occluder (Amplatzer, now St. Jude Medical). Following the limited FDA approval, use of PFO closure devices increased by more than 50-fold, well in excess of the 4000 per year threshold intended under the humanitarian device exemption,² prompting the FDA to withdraw the humanitarian device exemption approval for these devices in 2007. The Amplatzer PFO Occluder was approved through the premarket approval process in 2016.

In March 2018, the FDA granted an expanded indication to the Gore Cardioform Septal Occluder to include closure of PFO to reduce the risk of recurrent stroke (see [Table 3](#)). The new indication was based on results of the REDUCE pivotal clinical trial.³

Table 3. Patent Foramen Ovale Closure Devices Approved by the Food and Drug Administration

Device	Manufacturer	PMA Approval Date	Indications
Amplatzer™ PFO Occluder	St. Jude Medical	Nov 2016	For percutaneous transcatheter closure of a patent foramen ovale (PFO) to reduce the risk of recurrent ischemic stroke in patients, predominantly between the ages of 18 and 60 years, who have had a cryptogenic stroke due to a presumed paradoxical embolism, as determined by a neurologist and cardiologist following an evaluation to exclude known causes of ischemic stroke. ⁴
GORE HELEXSeptal Occluder	W.L. Gore & Associates	Aug 2006 (discontinued)	Percutaneous, transcatheter closure of ostium secundum ASDs
GORE CARDIOFORM Septal Occluder	W.L. Gore & Associates	Mar 2018 (supplement)	PFO closure to reduce the risk of recurrent ischemic stroke in patients, predominantly between the ages of 18 and 60 years, who have had a cryptogenic stroke due to a presumed paradoxical embolism, as determined by a neurologist and cardiologist following an evaluation to exclude known causes of ischemic stroke

PFO: patent foramen ovale; PMA: premarket approval. FDA product code: MLV.



Atrial Septal Defect (ASD) Closure Devices

The FDA has approved three devices for ASD closure through the premarket approval process or a premarket approval supplement: the Amplatzer Septal Occluder, the GORE HELEX Septal Occluder (discontinued), and the GORE CARDIOFORM Septal Occluder (see [Table 4](#)) (FDA product code: MLV).

Table 4. Atrial Septal Defect Closure Devices Approved by the Food and Drug Administration

Device	Manufacturer	PMA Approval Date	Indications
Amplatzer™ Septal Occluder	St. Jude Medical (Abbot Medical)	Dec 2001	Occlusion of ASDs in the secundum position Use in patients who have had a fenestrated Fontan procedure who require closure of the fenestration Patients indicated for ASD closure have echocardiographic evidence of ostium secundum ASD and clinical evidence of right ventricular volume overload
GORE HELEX Septal Occluder	W.L. Gore & Associates	Aug 2006 (discontinued)	Percutaneous, transcatheter closure of ostium secundum ASDs
GORE CARDIOFORM ASD Occluder (formerly GORE CARDIOFORM Septal Occluder)	W.L. Gore & Associates	May 2019 (supplement; name change) Oct 2016 (supplement)	Percutaneous, transcatheter closure of ostium secundum ASDs

ASD: atrial septal defect; PMA: premarket approval.

References

1. Messe SR, Kasner SE. Is closure recommended for patent foramen ovale and cryptogenic stroke? Patent foramen ovale in cryptogenic stroke: not to close. *Circulation*. Nov 04 2008; 118(19): 1999-2004. PMID 18981314



2. Slottow TL, Steinberg DH, Waksman R. Overview of the 2007 Food and Drug Administration Circulatory System Devices Panel meeting on patent foramen ovale closure devices. *Circulation*. Aug 07 2007; 116(6): 677-82. PMID 17679629
3. U.S. Food and Drug Administration. Summary of safety and effectiveness data. Gore Cardioform Septal Occluder. March 30, 2018. https://www.accessdata.fda.gov/cdrh_docs/pdf5/P050006s060b.pdf. Accessed June, 2020.
4. Food and Drug Administration (FDA). Summary of Safety and Effectiveness Data (SSED): Patent Foramen Ovale (PFO) Occluder (PMA P120021). 2016; https://www.accessdata.fda.gov/cdrh_docs/pdf12/P120021B.pdf. Accessed June, 2020.
5. Meier B, Kalesan B, Mattle HP, et al. Percutaneous closure of patent foramen ovale in cryptogenic embolism. *N Engl J Med*. Mar 21 2013; 368(12): 1083-91. PMID 23514285
6. Carroll JD, Saver JL, Thaler DE, et al. Closure of patent foramen ovale versus medical therapy after cryptogenic stroke. *N Engl J Med*. Mar 21 2013; 368(12): 1092-100. PMID 23514286
7. Saver JL, Carroll JD, Thaler DE, et al. Long-Term Outcomes of Patent Foramen Ovale Closure or Medical Therapy after Stroke. *N Engl J Med*. Sep 14 2017; 377(11): 1022-1032. PMID 28902590
8. Rogers T, Slack M, Waksman R. Overview of the 2016 US Food and Drug Administration Circulatory System Devices Panel Meeting on the Amplatzer Patent Foramen Ovale Occluder. *Am J Cardiol*. Jan 01 2017; 119(1): 153-155. PMID 27810099
9. Lee PH, Song JK, Kim JS, et al. Cryptogenic Stroke and High-Risk Patent Foramen Ovale: The DEFENSE-PFO Trial. *J Am Coll Cardiol*. May 22 2018; 71(20): 2335-2342. PMID 29544871
10. Sondergaard L, Kasner SE, Rhodes JF, et al. Patent Foramen Ovale Closure or Antiplatelet Therapy for Cryptogenic Stroke. *N Engl J Med*. Sep 14 2017; 377(11): 1033-1042. PMID 28902580
11. Mas JL, Derumeaux G, Guillon B, et al. Patent Foramen Ovale Closure or Anticoagulation vs. Antiplatelets after Stroke. *N Engl J Med*. Sep 14 2017; 377(11): 1011-1021. PMID 28902593
12. Kent DM, Dahabreh IJ, Ruthazer R, et al. Device Closure of Patent Foramen Ovale After Stroke: Pooled Analysis of Completed Randomized Trials. *J Am Coll Cardiol*. Mar 01 2016; 67(8): 907-17. PMID 26916479
13. Li J, Liu J, Liu M, et al. Closure versus medical therapy for preventing recurrent stroke in patients with patent foramen ovale and a history of cryptogenic stroke or transient ischemic attack. *Cochrane Database Syst Rev*. Sep 08 2015; (9): CD009938. PMID 26346232
14. Shah R, Nayyar M, Jovin IS, et al. Device Closure Versus Medical Therapy Alone for Patent Foramen Ovale in Patients With Cryptogenic Stroke: A Systematic Review and Meta-analysis. *Ann Intern Med*. Mar 06 2018; 168(5): 335-342. PMID 29310136
15. De Rosa S, Sievert H, Sabatino J, et al. Percutaneous Closure Versus Medical Treatment in Stroke Patients With Patent Foramen Ovale: A Systematic Review and Meta-analysis. *Ann Intern Med*. Mar 06 2018; 168(5): 343-350. PMID 29310133
16. Alushi B, Lauten A, Cassese S, et al. Patent foramen ovale closure versus medical therapy for prevention of recurrent cryptogenic embolism: updated meta-analysis of randomized clinical trials. *Clin Res Cardiol*. Sep 2018; 107(9): 788-798. PMID 29644412
17. Rigatelli G, Pedon L, Zecchel R, et al. Long-Term Outcomes and Complications of Intracardiac Echocardiography-Assisted Patent Foramen Ovale Closure in 1,000 Consecutive Patients. *J Interv Cardiol*. Oct 2016; 29(5): 530-538. PMID 27500752
18. Wintzer-Wehekind J, Alperi A, Houde C, et al. Long-Term Follow-Up After Closure of Patent Foramen Ovale in Patients With Cryptogenic Embolism. *J Am Coll Cardiol*. Jan 29 2019; 73(3): 278-287. PMID 30678757
19. Dowson A, Mullen MJ, Peatfield R, et al. Migraine Intervention With STARFlex Technology (MIST) trial: a prospective, multicenter, double-blind, sham-controlled trial to evaluate the effectiveness of patent foramen ovale closure with STARFlex septal repair implant to resolve refractory migraine headache. *Circulation*. Mar 18 2008; 117(11): 1397-404. PMID 18316488
20. Mattle HP, Evers S, Hildick-Smith D, et al. Percutaneous closure of patent foramen ovale in migraine with aura, a randomized controlled trial. *Eur Heart J*. Jul 07 2016; 37(26): 2029-36. PMID 26908949
21. Tobis JM, Charles A, Silberstein SD, et al. Percutaneous Closure of Patent Foramen Ovale in Patients With Migraine: The PREMIUM Trial. *J Am Coll Cardiol*. Dec 05 2017; 70(22): 2766-2774. PMID 29191325



22. Lip PZ, Lip GY. Patent foramen ovale and migraine attacks: a systematic review. *Am J Med.* May 2014; 127(5): 411-20. PMID 24355354
23. Biasco L, Infantino V, Orzan F, et al. Impact of transcatheter closure of patent foramen ovale in the evolution of migraine and role of residual shunt. *J Cardiol.* Nov 2014; 64(5): 390-4. PMID 24713153
24. Snijder RJ, Luermans JG, de Heij AH, et al. Patent Foramen Ovale With Atrial Septal Aneurysm Is Strongly Associated With Migraine With Aura: A Large Observational Study. *J Am Heart Assoc.* Dec 01 2016; 5(12). PMID 27930349
25. Tobis J, Shenoda M. Percutaneous treatment of patent foramen ovale and atrial septal defects. *J Am Coll Cardiol.* Oct 30 2012; 60(18): 1722-32. PMID 23040567
26. Mojadidi MK, Gevorgyan R, Noureddin N, et al. The effect of patent foramen ovale closure in patients with platypnea-orthodeoxia syndrome. *Catheter Cardiovasc Interv.* Oct 2015; 86(4): 701-7. PMID 26063336
27. Du ZD, Hijazi ZM, Kleinman CS, et al. Comparison between transcatheter and surgical closure of secundum atrial septal defect in children and adults: results of a multicenter nonrandomized trial. *J Am Coll Cardiol.* Jun 05 2002; 39(11): 1836-44. PMID 12039500
28. Butera G, Biondi-Zoccai G, Sangiorgi G, et al. Percutaneous versus surgical closure of secundum atrial septal defects: a systematic review and meta-analysis of currently available clinical evidence. *EuroIntervention.* Jul 2011; 7(3): 377-85. PMID 21729841
29. Abaci A, Unlu S, Alsancak Y, et al. Short and long term complications of device closure of atrial septal defect and patent foramen ovale: meta-analysis of 28,142 patients from 203 studies. *Catheter Cardiovasc Interv.* Dec 01 2013; 82(7): 1123-38. PMID 23412921
30. Suchon E, Pieculewicz M, Tracz W, et al. Transcatheter closure as an alternative and equivalent method to the surgical treatment of atrial septal defect in adults: comparison of early and late results. *Med Sci Monit.* Dec 2009; 15(12): CR612-7. PMID 19946231
31. Berger F, Vogel M, Alexi-Meskishvili V, et al. Comparison of results and complications of surgical and Amplatzer device closure of atrial septal defects. *J Thorac Cardiovasc Surg.* Oct 1999; 118(4): 674-8; discussion 678-80. PMID 10504632
32. Kotowycz MA, Therrien J, Ionescu-Iltu R, et al. Long-term outcomes after surgical versus transcatheter closure of atrial septal defects in adults. *JACC Cardiovasc Interv.* May 2013; 6(5): 497-503. PMID 23602461
33. Chen TH, Hsiao YC, Cheng CC, et al. In-Hospital and 4-Year Clinical Outcomes Following Transcatheter Versus Surgical Closure for Secundum Atrial Septal Defect in Adults: A National Cohort Propensity Score Analysis. *Medicine (Baltimore).* Sep 2015; 94(38): e1524. PMID 26402807
34. Xu XD, Liu SX, Zhao XX, et al. Comparison of medium-term results of transcatheter correction versus surgical treatment for secundum type atrial septal defect combined with pulmonary valve stenosis. *Int Heart J.* 2014; 55(4): 326-30. PMID 24898601
35. Fischer G, Stieh J, Uebing A, et al. Experience with transcatheter closure of secundum atrial septal defects using the Amplatzer septal occluder: a single centre study in 236 consecutive patients. *Heart.* Feb 2003; 89(2): 199-204. PMID 12527678
36. Javois AJ, Rome JJ, Jones TK, et al. Results of the U.S. Food and Drug Administration continued access clinical trial of the GORE HELEX septal occluder for secundum atrial septal defect. *JACC Cardiovasc Interv.* Aug 2014; 7(8): 905-12. PMID 25147036
37. Baruteau AE, Petit J, Lambert V, et al. Transcatheter closure of large atrial septal defects: feasibility and safety in a large adult and pediatric population. *Circ Cardiovasc Interv.* Dec 2014; 7(6): 837-43. PMID 25423959
38. Du ZD, Koenig P, Cao QL, et al. Comparison of transcatheter closure of secundum atrial septal defect using the Amplatzer septal occluder associated with deficient versus sufficient rims. *Am J Cardiol.* Oct 15 2002; 90(8): 865-9. PMID 12372575
39. Oho S, Ishizawa A, Akagi T, et al. Transcatheter closure of atrial septal defects with the Amplatzer septal occluder--a Japanese clinical trial. *Circ J.* Sep 2002; 66(9): 791-4. PMID 12224813
40. Brochu MC, Baril JF, Dore A, et al. Improvement in exercise capacity in asymptomatic and mildly symptomatic adults after atrial septal defect percutaneous closure. *Circulation.* Oct 01 2002; 106(14): 1821-6. PMID 12356636



41. Furlan AJ, Reisman M, Massaro J, et al. Closure or medical therapy for cryptogenic stroke with patent foramen ovale. *N Engl J Med.* Mar 15 2012; 366(11): 991-9. PMID 22417252
42. Grohmann J, Hohn R, Fleck T, et al. Transcatheter closure of atrial septal defects in children and adolescents: single-center experience with the GORE(R) septal occluder. *Catheter Cardiovasc Interv.* Nov 15 2014; 84(6): E51-7. PMID 24664494
43. Nyboe C, Hjortdal VE, Nielsen-Kudsk JE. First experiences with the GORE((R)) Septal Occluder in children and adults with atrial septal defects. *Catheter Cardiovasc Interv.* Nov 15 2013; 82(6): 929-34. PMID 23404677
44. Yilmazer MM, Guven B, Vupa-Cilengiroglu O, et al. Improvement in cardiac structure and functions early after transcatheter closure of secundum atrial septal defect in children and adolescents. *Turk J Pediatr.* Jul-Aug 2013; 55(4): 401-10. PMID 24292034
45. Jalal Z, Hascoet S, Gronier C, et al. Long-Term Outcomes After Percutaneous Closure of Ostium Secundum Atrial Septal Defect in the Young: A Nationwide Cohort Study. *JACC Cardiovasc Interv.* Apr 23 2018; 11(8): 795-804. PMID 29673513
46. Lansberg MG, O'Donnell MJ, Khatri P, et al. Antithrombotic and thrombolytic therapy for ischemic stroke: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest.* Feb 2012; 141(2 Suppl): e601S-e636S. PMID 22315273
47. Messe SR, Gronseth G, Kent DM, et al. Practice advisory: Recurrent stroke with patent foramen ovale (update of practice parameter): Report of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology. *Neurology.* Aug 23 2016; 87(8): 815-21. PMID 27466464
48. Kernan WN, Ovbiagele B, Black HR, et al. Guidelines for the prevention of stroke in patients with stroke and transient ischemic attack: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke.* Jul 2014; 45(7): 2160-236. PMID 24788967
49. Stout KK, Daniels CJ, Aboulhosn JA, et al. 2018 AHA/ACC Guideline for the Management of Adults With Congenital Heart Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol.* Apr 02 2019; 73(12): e81-e192. PMID 30121239

History

Date	Comments
09/07/99	Add to Medicine Section - New Policy
01/18/01	Replace policy - New information on patent foramen ovale; rest unchanged.
03/12/02	Replace policy - Revised; added requirements to policy statement patients with PFO must fail trial of oral anticoagulants. Noted FDA approval of Amplatzer device. Policy replaces 2.02.09.
08/13/02	Replace policy - Policy statement revised to indicate transcatheter treatment of ASD may be considered medically necessary. Replaces P2.02.100.
05/13/03	Replace policy - Policy reviewed; no change to policy statement; CPT codes updated.
05/26/06	Update Scope and Disclaimer - No other changes.
03/11/08	Cross Reference Update - No other changes.
10/14/08	Cross Reference Update - No other changes.



Date	Comments
03/10/09	Replace policy - Policy updated with literature search; policy rationale extensively revised. Policy statement for PFO changed to investigational due to the FDA's withdrawal of the humanitarian device exemption approval. References added.
06/08/10	Replace policy - Policy updated with literature search; no change to the policy statement. References added.
10/11/11	Replace policy – Policy updated with literature search. Policy statements unchanged. References 5, 8, 15 and 25 added. ICD-10 codes added to policy.
11/27/12	Replace policy - Policy updated with literature search. References 3, 6, 7, and 30 added. No change to policy statement.
12/04/13	Replace Policy. Policy guidelines reformatted for usability. Updated Regulatory Status with information about the FDA medical device alert issued 10/17/13 for the Amplatzer ASO. Rationale updated with literature search through July 31, 2013. References 4-7, 25 added; others renumbered/removed. Policy statement unchanged.
11/20/14	Annual Review. Policy updated with literature review through August 1, 2014. References 8-17, 21, 26, 33-37, 40, 48, 53-55, 58-59 added. Policy statement unchanged. ICD-9 and ICD-10 diagnosis codes removed; these are not utilized in adjudication of the policy.
06/09/15	Coding update: Correct ICD-10-PCS codes to support remediation efforts.
12/08/15	Annual Review. Policy updated with literature review through July, 2015; references 16-19, 48, 55-56, and 67 added. Policy statements unchanged.
07/01/16	Annual Review, approved June 14, 2016. Policy statements unchanged. Clinical input received from physician specialty societies and academic medical centers added. No new literature added.
01/01/17	Interim review, approved December 13, 2016. Changed policy statement from investigational to medically necessary for closure of PFO in the presence of cryptogenic stroke due to paradoxical embolism using a PFO occluder device (Amplatzer PFO) when criteria are met. Updated Regulatory Status with information about the Amplatzer device for PFO. Policy updated with literature review through October 2016.
07/01/17	Annual Review, approved June 22, 2017. Policy moved into new format. Policy updated with literature review through March 23, 2017; references 3, 6-7, 9-10, 48-49, 51-52, 64, and 78 added. Statement, "There are currently no transcatheter devices with the U.S. Food and Drug Administration [FDA] approval or clearance for this indication," removed from investigational statement for PFO closure devices; policy statements otherwise unchanged.
08/01/18	Annual Review, approved July 10, 2018. Policy updated with literature review through March 2018; references 9-11, 14-15, and 17 added. Policy statement changed to: The percutaneous transcatheter closure of a patent foramen ovale using AMPLATZER PFO Occluder may be considered medically necessary to reduce the risk of recurrent



Date	Comments
	ischemic stroke if patient meets all of the specified criteria.
08/10/18	Corrected errors in the description of the heart anatomy discussed under Background, Patent Foramen Ovale on page 5.
04/01/19	Minor update, added Documentation Requirements section.
08/01/19	Annual Review, approved July 25, 2019. Policy updated with literature review through March 2019; references added. Added new FDA approved patent foramen ovale closure device: Gore Cardioform Septal Occluder. An investigational statement was added for situations not meeting criteria.
04/01/20	Delete policy, approved March 10, 2020. This policy will be deleted effective July 2, 2020, and replaced with InterQual criteria for dates of service on or after July 2, 2020.
05/06/20	Interim Review, approved May 5, 2020. This policy is reinstated immediately and will no longer be deleted or replaced with InterQual criteria on July 2, 2020.
08/01/20	Annual Review, approved July 2, 2020. Policy updated with literature review through March 9, 2020; references added. Policy statements unchanged.

Disclaimer: This medical policy is a guide in evaluating the medical necessity of a particular service or treatment. The Company adopts policies after careful review of published peer-reviewed scientific literature, national guidelines and local standards of practice. Since medical technology is constantly changing, the Company reserves the right to review and update policies as appropriate. Member contracts differ in their benefits. Always consult the member benefit booklet or contact a member service representative to determine coverage for a specific medical service or supply. CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). ©2020 Premera All Rights Reserved.

Scope: Medical policies are systematically developed guidelines that serve as a resource for Company staff when determining coverage for specific medical procedures, drugs or devices. Coverage for medical services is subject to the limits and conditions of the member benefit plan. Members and their providers should consult the member benefit booklet or contact a customer service representative to determine whether there are any benefit limitations applicable to this service or supply. This medical policy does not apply to Medicare Advantage.



Discrimination is Against the Law

Premera Blue Cross complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex. Premera does not exclude people or treat them differently because of race, color, national origin, age, disability or sex.

Premera:

- Provides free aids and services to people with disabilities to communicate effectively with us, such as:
 - Qualified sign language interpreters
 - Written information in other formats (large print, audio, accessible electronic formats, other formats)
- Provides free language services to people whose primary language is not English, such as:
 - Qualified interpreters
 - Information written in other languages

If you need these services, contact the Civil Rights Coordinator.

If you believe that Premera has failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, or sex, you can file a grievance with:

Civil Rights Coordinator - Complaints and Appeals
PO Box 91102, Seattle, WA 98111
Toll free 855-332-4535, Fax 425-918-5592, TTY 800-842-5357
Email AppealsDepartmentInquiries@Premera.com

You can file a grievance in person or by mail, fax, or email. If you need help filing a grievance, the Civil Rights Coordinator is available to help you.

You can also file a civil rights complaint with the U.S. Department of Health and Human Services, Office for Civil Rights, electronically through the Office for Civil Rights Complaint Portal, available at <https://ocrportal.hhs.gov/ocr/portal/lobby.jsf>, or by mail or phone at: U.S. Department of Health and Human Services
200 Independence Avenue SW, Room 509F, HHH Building
Washington, D.C. 20201, 1-800-368-1019, 800-537-7697 (TDD)
Complaint forms are available at <http://www.hhs.gov/ocr/office/file/index.html>.

Getting Help in Other Languages

This Notice has Important Information. This notice may have important information about your application or coverage through Premera Blue Cross. There may be key dates in this notice. You may need to take action by certain deadlines to keep your health coverage or help with costs. You have the right to get this information and help in your language at no cost. Call 800-722-1471 (TTY: 800-842-5357).

አማርኛ (Amharic):

ይህ ማስታወቂያ አስፈላጊ መረጃ ይዟል። ይህ ማስታወቂያ ስለ ማመልከቻዎ ወይም የ Premera Blue Cross ሽፋን አስፈላጊ መረጃ ሊኖረው ይችላል። በዚህ ማስታወቂያ ውስጥ ቁልፍ ቀናት ሊኖሩ ይችላሉ። የጤና ሽፋንዎን ለመጠበቅና በአስፈላጊ እርዳታ ለማግኘት በተውሰኑ የጊዜ ገደቦች እርምጃ መውሰድ ይገባዎት ይሆናል። ይህን መረጃ እንዲያገኙ እና የለምንም ክፍያ በቋንቋዎ እርዳታ እንዲያገኙ መሰታወቅ አለዎት። በስልክ ቁጥር 800-722-1471 (TTY: 800-842-5357) ይደውሉ።

العربية (Arabic):

يحتوي هذا الإشعار على معلومات هامة. قد يحتوي هذا الإشعار على معلومات مهمة بخصوص طلبك أو التغطية التي تزيد الحصول عليها من خلال Premera Blue Cross. قد تكون هناك تواريخ مهمة في هذا الإشعار. وقد تحتاج لاتخاذ إجراء في تاريخ معينه للحفاظ على تغطيتك الصحية أو للمساعدة في دفع التكاليف. يحق لك الحصول على هذه المعلومات والمساعدة بلغتك دون تكبد أية تكلفة. اتصل بـ 800-722-1471 (TTY: 800-842-5357)

中文 (Chinese):

本通知有重要的訊息。本通知可能有關於您透過 Premera Blue Cross 提交的申請或保險的重要訊息。本通知內可能有重要日期。您可能需要在截止日期之前採取行動，以保留您的健康保險或者費用補貼。您有權利免費以您的母語得到本訊息和幫助。請撥電話 800-722-1471 (TTY: 800-842-5357)。

Oromoo (Cushite):

Beeksisni kun odeeffannoo barbaachisaa qaba. Beeksisni kun sagantaa yookan karaa Premera Blue Cross tiin tajaajila keessan ilaalchisee odeeffannoo barbaachisaa qabaachuu danda'a. Guyyaawwan murteessaa ta'an beeksisa kana keessatti ilaalaa. Tarii kaffaltiidhaan deeggaramuuf yookan tajaajila fayyaa keessaniif guyyaa dhumaa irratti wanti raawwattan jiraachuu danda'a. Kaffaltii irraa bilisa haala ta'een afaan keessaniin odeeffannoo argachuu fi deeggarsa argachuuf mirga ni qabaattu. Lakkoofsa bilbilaa 800-722-1471 (TTY: 800-842-5357) tii bilbilaa.

Français (French):

Cet avis a d'importantes informations. Cet avis peut avoir d'importantes informations sur votre demande ou la couverture par l'intermédiaire de Premera Blue Cross. Le présent avis peut contenir des dates clés. Vous devez peut-être prendre des mesures par certains délais pour maintenir votre couverture de santé ou d'aide avec les coûts. Vous avez le droit d'obtenir cette information et de l'aide dans votre langue à aucun coût. Appelez le 800-722-1471 (TTY: 800-842-5357).

Kreyòl ayisyen (Creole):

Avi sila a gen Enfòmasyon Enpòtan ladann. Avi sila a kapab genyen enfòmasyon enpòtan konsènan aplikasyon w lan oswa konsènan kouvèti asirans lan atravè Premera Blue Cross. Kapab genyen dat ki enpòtan nan avi sila a. Ou ka gen pou pran kèk aksyon avan sèten dat limit pou ka kenbe kouvèti asirans sante w la oswa pou yo ka ede w avèk depans yo. Se dwa w pou resewva enfòmasyon sa a ak asistans nan lang ou pale a, san ou pa gen pou peye pou sa. Rele nan 800-722-1471 (TTY: 800-842-5357).

Deutsche (German):

Diese Benachrichtigung enthält wichtige Informationen. Diese Benachrichtigung enthält unter Umständen wichtige Informationen bezüglich Ihres Antrags auf Krankenversicherungsschutz durch Premera Blue Cross. Suchen Sie nach eventuellen wichtigen Terminen in dieser Benachrichtigung. Sie könnten bis zu bestimmten Stichtagen handeln müssen, um Ihren Krankenversicherungsschutz oder Hilfe mit den Kosten zu behalten. Sie haben das Recht, kostenlose Hilfe und Informationen in Ihrer Sprache zu erhalten. Rufen Sie an unter 800-722-1471 (TTY: 800-842-5357).

Hmoob (Hmong):

Tsab ntawv tshaj xo no muaj cov ntshiab lus tseem ceeb. Tej zaum tsab ntawv tshaj xo no muaj cov ntshiab lus tseem ceeb txog koj daim ntawv thov kev pab los yog koj qhov kev pab cuam hnu ntawm Premera Blue Cross. Tej zaum muaj cov hnuv tseem ceeb uas sau rau hauv daim ntawv no. Tej zaum koj kuj yuav tau ua qee yam uas pab kom koj ua tsis pub dhau cov caij nyoog uas teev tseg rau hauv daim ntawv no mas koj thiaj yuav tau txais kev pab cuam kho mob los yog kev pab them tej nqi kho mob ntawd. Koj muaj cai kom lawv muab cov ntshiab lus no uas tau muab sau ua koj hom lus pub dawb rau koj. Hu rau 800-722-1471 (TTY: 800-842-5357).

Iloko (Ilocano):

Daytoy a Pakdaar ket naglaon iti Napateg nga Impormasion. Daytoy a pakdaar mabalin nga adda ket naglaon iti napateg nga impormasion maipanggep iti aplikasyonyo wenna coverage babaen iti Premera Blue Cross. Daytoy ket mabalin dagiti importante a petsa iti daytoy a pakdaar. Mabalin nga adda rumbeng nga aramidenyo nga addang sakbay dagiti partikular a naituding nga aldaw tapno mapagtalinaedyo ti coverage ti salun-atyto wenna tulong kadagiti gastos. Adda karbenganyo a mangala iti daytoy nga impormasion ken tulong iti bukodyo a pagsasao nga awan ti bayadanyo. Tumawag iti numero nga 800-722-1471 (TTY: 800-842-5357).

Italiano (Italian):

Questo avviso contiene informazioni importanti. Questo avviso può contenere informazioni importanti sulla tua domanda o copertura attraverso Premera Blue Cross. Potrebbero esserci date chiave in questo avviso. Potrebbe essere necessario un tuo intervento entro una scadenza determinata per consentirti di mantenere la tua copertura o sovvenzione. Hai il diritto di ottenere queste informazioni e assistenza nella tua lingua gratuitamente. Chiama 800-722-1471 (TTY: 800-842-5357).

日本語 (Japanese):

この通知には重要な情報が含まれています。この通知には、Premera Blue Cross の申請または補償範囲に関する重要な情報が含まれている場合があります。この通知に記載されている可能性がある重要な日付をご確認ください。健康保険や有料サポートを維持するには、特定の期日までに行動を取らなければならない場合があります。ご希望の言語による情報とサポートが無料で提供されます。800-722-1471 (TTY: 800-842-5357)までお電話ください。

한국어 (Korean):

본 통지서에는 중요한 정보가 들어 있습니다. 즉 이 통지서는 귀하의 신청에 관하여 그리고 Premera Blue Cross 를 통한 커버리지에 관한 정보를 포함하고 있을 수 있습니다. 본 통지서에는 핵심이 되는 날짜들이 있을 수 있습니다. 귀하의 귀하의 건강 커버리지를 계속 유지하거나 비용을 절감하기 위해서 일정한 마감일까지 조치를 취해야 할 필요가 있을 수 있습니다. 귀하의 이러한 정보와 도움을 귀하의 언어로 비용 부담없이 얻을 수 있는 권리가 있습니다. 800-722-1471 (TTY: 800-842-5357) 로 전화하십시오.

ລາວ (Lao):

ແຈ້ງການນີ້ມີຂໍ້ມູນສໍາຄັນ. ແຈ້ງການນີ້ອາດຈະມີຂໍ້ມູນສໍາຄັນກ່ຽວກັບຄໍາຮ້ອງສະໝັກ ຫຼື ຄວາມຄົມຄອງປະກັນໄພຂອງທ່ານຜ່ານ Premera Blue Cross. ອາດຈະມີວັນທີ່ສໍາຄັນໃນແຈ້ງການນີ້. ທ່ານອາດຈະຈໍາເປັນຕ້ອງດໍາເນີນການຕາມກຳນົດ ເວລາສະເພາະເພື່ອຮັກສາຄວາມຄົມຄອງປະກັນສະພາບ ຫຼື ຄວາມຊ່ວຍເຫຼືອເວັ້ນເວົ້ອງຄ່າໃຊ້ຈ່າຍຂອງທ່ານໄດ້. ທ່ານມີສິດໄດ້ຮັບຂໍ້ມູນນີ້ ແລະ ຄວາມຊ່ວຍເຫຼືອເປັນພາສາຂອງທ່ານໂດຍບໍ່ເສຍຄ່າ. ໃຫ້ໃບທາ 800-722-1471 (TTY: 800-842-5357).

ភាសាខ្មែរ (Khmer):

សេចក្តីជូនដំណឹងនេះមានព័ត៌មានយ៉ាងសំខាន់។ សេចក្តីជូនដំណឹងនេះប្រហែលជាមានព័ត៌មានយ៉ាងសំខាន់អំពីទម្រង់បែបបទ ឬការរៀបចំរបស់អ្នកកាមរយ: Premera Blue Cross ។ ប្រហែលជាមាន កាលបរិច្ឆេទសំខាន់នៅក្នុងសេចក្តីជូនដំណឹងនេះ។ អ្នកប្រហែលជាត្រូវការបញ្ជាក់សមត្ថភាព ដល់កំណត់ថ្លៃជាតំបន់នានា ដើម្បីនឹងរក្សាទុកការធានារ៉ាប់រងអន្តរជាតិរបស់អ្នក ឬប្រាក់ដុល្លារចេញថ្លៃ។ អ្នកមានសិទ្ធិទទួលព័ត៌មាននេះ និងដុល្លារនៅក្នុងភាសារបស់អ្នកដោយមិនអស់លុយឡើយ។ សូមទូរស័ព្ទ 800-722-1471 (TTY: 800-842-5357)។

ਪੰਜਾਬੀ (Punjabi):

ਇਸ ਨੋਟਿਸ ਵਿਚ ਖਾਸ ਜਾਣਕਾਰੀ ਹੈ. ਇਸ ਨੋਟਿਸ ਵਿਚ Premera Blue Cross ਵਲੋਂ ਤੁਹਾਡੀ ਕਵਰੇਜ ਅਤੇ ਅਰਜੀ ਬਾਰੇ ਮਹੱਤਵਪੂਰਨ ਜਾਣਕਾਰੀ ਹੋ ਸਕਦੀ ਹੈ . ਇਸ ਨੋਟਿਸ ਨਵ ਖਾਸ ਤਾਰੀਖਾਂ ਹੋ ਸਕਦੀਆਂ ਹਨ. ਜੇਕਰ ਤੁਸੀਂ ਜਸਰਤ ਕਵਰੇਜ ਰਿੱਖਣੀ ਹੋਵੇ ਜਾਂ ਓਸ ਦੀ ਲਾਗਤ ਜਵਿੱਚ ਮਦਦ ਦੇ ਇਛੁੱਕ ਹੋ ਤਾਂ ਤੁਹਾਨੂੰ ਅੰਤਮ ਤਾਰੀਖ ਤੋਂ ਪਹਿਲਾਂ ਢੁੱਝ ਖਾਸ ਕਦਮ ਚੁੱਕਣ ਦੀ ਲੋੜ ਹੋ ਸਕਦੀ ਹੈ ,ਤੁਹਾਨੂੰ ਮੁਫਤ ਵਿੱਚ ਤੋਂ ਅਪਣੀ ਭਾਸ਼ਾ ਵਿੱਚ ਜਾਣਕਾਰੀ ਅਤੇ ਮਦਦ ਪ੍ਰਾਪਤ ਕਰਨ ਦਾ ਅਧਿਕਾਰ ਹੈ ,ਕਾਲ 800-722-1471 (TTY: 800-842-5357).

فارسی (Farsi):

این اعلامیه حاوی اطلاعات مهم میباشد. این اعلامیه ممکن است حاوی اطلاعات مهم درباره فرم تقاضا و یا پوشش بیمه ای شما از طریق Premera Blue Cross باشد. به تاریخ های مهم در این اعلامیه توجه نمایید. شما ممکن است برای حفظ پوشش بیمه تان یا کمک در پرداخت هزینه های درمانی تان، به تاریخ های مشخصی برای انجام کارهای خاصی احتیاج داشته باشید. شما حق این را دارید که این اطلاعات و کمک را به زبان خود به طور رایگان دریافت نمایید. برای کسب اطلاعات با شماره 800-722-1471 (کلیر بران TTY تماس باشماره 800-842-5357) تماس برقرار نمایید.

Polskie (Polish):

To ogłoszenie może zawierać ważne informacje. To ogłoszenie może zawierać ważne informacje odnośnie Państwa wniosku lub zakresu świadczeń poprzez Premera Blue Cross. Prosimy zwrócić uwagę na kluczowe daty, które mogą być zawarte w tym ogłoszeniu aby nie przekroczyć terminów w przypadku utrzymania polisy ubezpieczeniowej lub pomocy związanej z kosztami. Macie Państwo prawo do bezpłatnej informacji we własnym języku. Zadzwońcie pod 800-722-1471 (TTY: 800-842-5357).

Português (Portuguese):

Este aviso contém informações importantes. Este aviso poderá conter informações importantes a respeito de sua aplicação ou cobertura por meio do Premera Blue Cross. Poderão existir datas importantes neste aviso. Talvez seja necessário que você tome providências dentro de determinados prazos para manter sua cobertura de saúde ou ajuda de custos. Você tem o direito de obter esta informação e ajuda em seu idioma e sem custos. Ligue para 800-722-1471 (TTY: 800-842-5357).

Română (Romanian):

Prezenta notificare conține informații importante privind cererea sau acoperirea asigurării dumneavoastră de sănătate prin Premera Blue Cross. Pot exista date cheie în această notificare. Este posibil să fie nevoie să acționați până la anumite termene limită pentru a vă menține acoperirea asigurării de sănătate sau asistența provizorie la costuri. Aveți dreptul de a obține gratuit aceste informații și ajutor în limba dumneavoastră. Sunați la 800-722-1471 (TTY: 800-842-5357).

Русский (Russian):

Настоящее уведомление содержит важную информацию. Это уведомление может содержать важную информацию о вашем заявлении или страховом покрытии через Premera Blue Cross. В настоящем уведомлении могут быть указаны ключевые даты. Вам, возможно, потребуется принять меры к определенным предельным срокам для сохранения страхового покрытия или помощи с расходами. Вы имеете право на бесплатное получение этой информации и помощь на вашем языке. Звоните по телефону 800-722-1471 (TTY: 800-842-5357).

Fa'asamoa (Samoan):

Atonu ua iai i lenei fa'asilasilaga ni fa'amatalaga e sili ona taua e tatau ona e malamalama i ai. O lenei fa'asilasilaga o se fesoasoani e fa'amatala atili i ai i le tulaga o le polokalame, Premera Blue Cross, ua e tau fia maua atu i ai. Fa'amolemole, ia e iloilo fa'alelei i aso fa'apitoa olo'o iai i lenei fa'asilasilaga taua. Masalo o le'a iai ni feau e tatau ona e faia ao le'i aulia le aso ua ta'ua i lenei fa'asilasilaga ina ia e iai pea ma maua fesoasoani mai ai i le polokalame a le Malo olo'o e iai i ai. Olo'o iai iate oe le aia tatau e maua atu i lenei fa'asilasilaga ma lenei fa'matalaga i legagana e te malamalama i ai aunoa ma se togiga tupe. Vili atu i le telefoni 800-722-1471 (TTY: 800-842-5357).

Español (Spanish):

Este Aviso contiene información importante. Es posible que este aviso contenga información importante acerca de su solicitud o cobertura a través de Premera Blue Cross. Es posible que haya fechas clave en este aviso. Es posible que deba tomar alguna medida antes de determinadas fechas para mantener su cobertura médica o ayuda con los costos. Usted tiene derecho a recibir esta información y ayuda en su idioma sin costo alguno. Llame al 800-722-1471 (TTY: 800-842-5357).

Tagalog (Tagalog):

Ang Paunawa na ito ay naglalaman ng mahalagang impormasyon tungkol sa iyong aplikasyon o pagsakop sa pamamagitan ng Premera Blue Cross. Maaaring may mga mahalagang petsa dito sa paunawa. Maaring mangailangan ka na magsagawa ng hakbang sa ilang mga itinakdang panahon upang mapanatili ang iyong pagsakop sa kalusugan o tulong na walang gastos. May karapatan ka na makakuha ng ganiitong impormasyon at tulong sa iyong wika ng walang gastos. Tumawag sa 800-722-1471 (TTY: 800-842-5357).

ไทย (Thai):

ประกาศนี้มีข้อมูลสำคัญ ประกาศนี้อาจมีข้อมูลที่สำคัญเกี่ยวกับกาการสมัครหรือขอบเขตประกันสุขภาพของคุณผ่าน Premera Blue Cross และอาจมีกำหนดการในประกาศนี้ คุณอาจจะต้องดำเนินการภายในกำหนดระยะเวลาที่แน่นอนเพื่อจะรักษาการประกันสุขภาพของคุณหรือการช่วยเหลือที่มีค่าใช้จ่าย คุณมีสิทธิที่จะได้รับข้อมูลและความช่วยเหลือนี้ในภาษาของคุณโดยไม่มีค่าใช้จ่าย โทร 800-722-1471 (TTY: 800-842-5357)

Український (Ukrainian):

Це повідомлення містить важливу інформацію. Це повідомлення може містити важливу інформацію про Ваше звернення щодо страховального покриття через Premera Blue Cross. Зверніть увагу на ключові дати, які можуть бути вказані у цьому повідомленні. Існує імовірність того, що Вам треба буде здійснити певні кроки у конкретні кінцеві строки для того, щоб зберегти Ваше медичне страхування або отримати фінансову допомогу. У Вас є право на отримання цієї інформації та допомоги безкоштовно на Вашій рідній мові. Дзвоніть за номером телефону 800-722-1471 (TTY: 800-842-5357).

Tiếng Việt (Vietnamese):

Thông báo này cung cấp thông tin quan trọng. Thông báo này có thông tin quan trọng về đơn xin tham gia hoặc hợp đồng bảo hiểm của quý vị qua chương trình Premera Blue Cross. Xin xem ngày quan trọng trong thông báo này. Quý vị có thể phải thực hiện theo thông báo đúng trong thời hạn để duy trì bảo hiểm sức khỏe hoặc được trợ giúp thêm về chi phí. Quý vị có quyền được biết thông tin này và được trợ giúp bằng ngôn ngữ của mình miễn phí. Xin gọi số 800-722-1471 (TTY: 800-842-5357).