

MEDICAL POLICY – 2.02.26

Percutaneous Left Atrial Appendage Closure Devices for Stroke Prevention in Atrial Fibrillation

BCBSA Ref. Policy: 2.02.26

Effective Date: Aug. 1, 2018

Last Revised: July 13, 2018


Replaces: N/A

RELATED MEDICAL POLICIES:

None

Select a hyperlink below to be directed to that section.

[POLICY CRITERIA](#) | [DOCUMENTATION REQUIREMENTS](#) | [CODING](#)
[RELATED INFORMATION EVIDENCE REVIEW](#) | [REFERENCES](#) | [HISTORY](#)

 Clicking this icon returns you to the hyperlinks menu above.

Introduction

The heart is divided into two upper and two lower chambers. Atrial fibrillation, also called a-fib, occurs when the heart's upper chambers beat irregularly—and often rapidly. Because blood isn't pumped out the way that it should be, blood tends to pool in these two upper chambers. The pooling blood increases the risk of blood clots in the area of the heart called the left atrial appendage. If a blood clot comes loose, it may travel to the brain and cause a stroke. Blood thinners are the usual method of preventing blood clots in people with a-fib. If taking a blood thinner poses too much risk or a person can't tolerate this medication, placing a device in the heart is a different way of helping to prevent stroke. This device seals off the left atrial appendage. Should a clot develop, the device blocks it from entering the bloodstream. This policy describes when a left atrial appendage closure device is considered medically necessary.

Note: The Introduction section is for your general knowledge and is not to be taken as policy coverage criteria. The rest of the policy uses specific words and concepts familiar to medical professionals. It is intended for providers. A provider can be a person, such as a doctor, nurse, psychologist, or dentist. A provider also can be a place where medical care is given, like a hospital, clinic, or lab. This policy informs them about when a service may be covered.

Policy Coverage Criteria

Device	Medical Necessity
<p>Percutaneous left atrial appendage closure device (eg, the Watchman)</p>	<p>The use of a device with U.S. Food and Drug Administration (FDA) approval for percutaneous left atrial appendage closure (eg, the Watchman) may be considered medically necessary for the prevention of stroke in patients with atrial fibrillation when the following criteria are met:</p> <ul style="list-style-type: none"> • There is an increased risk of stroke and systemic embolism based on CHADS₂ or CHA₂DS₂-VASc score and systemic anticoagulation therapy is recommended <p>AND</p> <ul style="list-style-type: none"> • The long-term risks of systemic anticoagulation outweigh the risks of the device implantation (see Related Information) <p>The use of a device with FDA approval for percutaneous left atrial appendage closure (eg, the Watchman) for stroke prevention in patients who do not meet the above criteria is considered investigational.</p>

Device	Investigational
<p>Other percutaneous left atrial appendage closure devices</p>	<p>The use of other percutaneous left atrial appendage closure devices, including but not limited to the Lariat and Amplatzer devices, for stroke prevention in patients with atrial fibrillation is considered investigational.</p>

Documentation Requirements

The patient's medical records submitted for review for all conditions should document that medical necessity criteria are met. The record should include ALL of the following:

- Name of the Food and Drug Administration (FDA) device to be used
- CHADS₂ or CHA₂DS₂-VASc score documenting patient's increased risk of stroke and systemic embolism
- Documentation that systemic anticoagulation therapy is recommended AND the long-term risks of systemic anticoagulation outweigh the risks of the device implantation



Coding

Code	Description
CPT	
33340	Percutaneous transcatheter closure of the left atrial appendage with endocardial implant, including fluoroscopy, transeptal puncture, catheter placement(s), left atrial angiography, left atrial appendage angiography, when performed, and radiological supervision and interpretation

Note: CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). HCPCS codes, descriptions and materials are copyrighted by Centers for Medicare Services (CMS).

Related Information

The balance of risks and benefits associated with implantation of the Watchman device for stroke prevention, as an alternative to systemic anticoagulation with warfarin, must be made on an individual basis.

Bleeding is the primary risk associated with systemic anticoagulation. A number of risk scores have been developed to estimate the risk of significant bleeding in patients treated with systemic anticoagulation. An example is the HAS-BLED score, which has been validated to assess the annual risk of significant bleeding in patients with atrial fibrillation treated with warfarin (Pisters et al, 2010). The score ranges from 0 to 9, based on a number of clinical characteristics (see [Table 1](#)).

Table 1: Clinical Components of the HAS-BLED Bleeding Risk Score

Letter	Clinical Characteristic	Points Awarded
H	Hypertension	1
A	Abnormal renal and liver function (1 point each)	1 or 2
S	Stroke	1
B	Bleeding	1
L	Labile international normalized ratios	1
E	Elderly (>65 y)	1



Letter	Clinical Characteristic	Points Awarded
D	Drugs or alcohol (1 point each)	1 or 2

Adapted from Pisters et al (2010)

Risk of major bleeding in patients with scores of 3, 4, and 5 has been reported at 3.74 per 100 patient-years, 8.70 per 100 patient-years, and 12.5 per 100 patient-years, respectively. Scores of 3 or greater are considered to be associated with a high risk of bleeding, potentially signaling the need for closer monitoring of patients for adverse risks, closer monitoring of international normalized ratio, or differential dose selections of oral anticoagulants or aspirin (January et al, 2014).

Evidence Review

Description

Stroke prevention in atrial fibrillation (AF) is an important goal of treatment. Treatment with anticoagulant medications is the most common approach to stroke prevention. Most embolic strokes originate from the left atrial appendage; therefore, occlusion of the left atrial appendage may offer a nonpharmacologic alternative to anticoagulant medications for this purpose. Multiple percutaneously deployed devices are being investigated for left atrial appendage closure (LAAC). One left atrial appendage device (the Watchman device) has approval from the U.S. Food and Drug Administration for stroke prevention in patients with AF.

Background

Stroke

Stroke is the most serious complication of atrial fibrillation (AF). The estimated incidence of stroke in nontreated patients with AF is 5% per year. Stroke associated with AF is primarily embolic in nature, tends to be more severe than the typical ischemic stroke, and causes higher rates of mortality and disability. As a result, stroke prevention is one of the main goals of AF treatment.



Stroke in AF occurs primarily as a result of thromboembolism from the left atrium. The lack of atrial contractions in AF leads to blood stasis in the left atrium, and this low flow state increases the risk for thrombosis. The area of the left atrium with the lowest blood flow in AF, and, therefore, the highest risk of thrombosis, is the left atrial appendage (LAA). It has been estimated that 90% of left atrial thrombi occur in the LAA.

Treatment

Pharmacologic

The main treatment for stroke prevention in AF is anticoagulation, which has proven efficacy. The risk for stroke among patients with AF is evaluated using several factors. Two commonly used scores, the CHADS₂ and the CHADS₂-VASc score, are described below in **Table 2**. Warfarin is the predominant anticoagulation agent in clinical use. A number of newer anticoagulant medications, including dabigatran, rivaroxaban, and apixaban, have recently received U.S. Food and Drug Administration (FDA) approval for stroke prevention in nonvalvular AF and have demonstrated noninferiority to warfarin in clinical trials. While anticoagulation is effective for stroke prevention, it carries an increased risk of bleeding. Also, warfarin requires frequent monitoring and adjustments, as well as lifestyle changes. Dabigatran does not require monitoring. However, unlike warfarin, the antithrombotic effects of dabigatran are not reversible with any currently available hemostatic drugs. Guidelines from the American College of Chest Physicians (2012) have recommended the use of oral anticoagulation for patients with AF who are at high risk of stroke (ie, CHADS₂ score ≥ 2), with more individualized choice of antithrombotic therapy in patients with lower stroke risk.¹

Table 2. CHADS₂ and CHADS₂-VASc Scores to Predict Ischemic Stroke Risk in Patients with Atrial Fibrillation

Letter	Clinical Characteristics	Points Awarded
C	Congestive heart failure (signs/symptoms of heart failure confirmed with objective evidence of cardiac dysfunction)	1
H	Hypertension (resting blood pressure >140/90 mmHg on at least 2 occasions or current antihypertensive pharmacologic treatment)	1
A	Age ≥ 75 y	2



Letter	Clinical Characteristics	Points Awarded
D	Diabetes (fasting glucose >125 mg/dL or treatment with oral hypoglycemic agent and/or insulin)	1
S	Stroke or transient ischemic attack (includes any history of cerebral ischemia)	2
V	Vascular disease (prior myocardial infarction, peripheral arterial disease, or aortic plaque)	1
A	Age 65-74 y	1
Sc	Sex category of female (female sex confers higher risk)	1

Adapted from You et al (2012)¹ and January et al (2014).²

Bleeding is the primary risk associated with systemic anticoagulation. Risk scores have been developed to estimate the risk of significant bleeding in patients treated with systemic anticoagulation, such as the HAS-BLED score, which has been validated to assess the annual risk of significant bleeding in patients with AF treated with warfarin.³ The score ranges from 0 to 9, based on a number of clinical characteristics, including the presence of hypertension, renal and liver function, history of stroke, bleeding, labile international normalized ratios, age, and drug/alcohol use. Scores of 3 or greater are considered to be associated with high risk of bleeding, potentially signaling the need for closer monitoring of patients for adverse risks, closer monitoring of international normalized ratios, or differential dose selections of oral anticoagulants or aspirin.²

Surgery

Surgical removal, or exclusion, of the LAA is often performed in patients with AF who are undergoing open heart surgery for other reasons. Percutaneous left atrial appendage closure (LAAC) closure devices have been developed as a nonpharmacologic alternative to anticoagulation for stroke prevention in AF. The devices may prevent stroke by occluding the LAA, thus preventing thrombus formation.

Several versions of LAA occlusion devices have been developed. The Watchman Left Atrial Appendage System (Boston Scientific) is a self-expanding nickel titanium device. It has a polyester covering and fixation barbs for attachment to the endocardium. Implantation is performed percutaneously through a catheter delivery system, using venous access and transseptal puncture to enter the left atrium. Following implantation, patients receive anticoagulation with warfarin or alternative agents for approximately 1 to 2 months. After this



period, patients are maintained on antiplatelet agents (ie, aspirin and/or clopidogrel) indefinitely. The Lariat Loop Applicator is a suture delivery device that is intended to close a variety of surgical wounds in addition to LAAC. The Cardioblade® closure device (Medtronic) is currently being tested in clinical studies. The Amplatzer cardiac plug (St. Jude Medical), is FDA-approved for closure of atrial septal defects but not for LAAC. A second-generation device, the Amplatzer Amulet, has been developed. The Percutaneous LAA Transcatheter Occlusion device (ev3) has also been evaluated in research studies but has not received FDA approval. The Occlutech® (Occlutech) Left Atrial Appendage Occluder has received a CE mark for coverage in Europe.

Outcome Measures

The optimal study design for evaluating the efficacy of percutaneous LAAC for the prevention of stroke in AF is a randomized controlled trial that includes clinically relevant measures of health outcomes. The rate of ischemic stroke during follow-up is the primary outcome of interest, along with rates of systemic embolization, cardiac events, bleeding complications, and death. For the LAAC devices, the appropriate comparison group could be oral anticoagulation, no therapy (for patients who have a prohibitive risk for oral anticoagulation), or open surgical repair.

Although the Watchman device and other LAAC devices would ideally represent an alternative to oral anticoagulation for the prevention of stroke in patients with AF, during the postimplantation period, the device may be associated with increased thrombogenicity and, therefore, anticoagulation is used during the periprocedural period. Most studies evaluating the Watchman device have included patients who are eligible for anticoagulation.

Summary of Evidence

For individuals who have AF who are at increased risk for embolic stroke who receive the Watchman percutaneous LAAC device, the evidence includes 2 RCTs and meta-analyses of these trials. Relevant outcomes are overall survival, morbid events, and treatment-related morbidity. The most relevant evidence comes from 2 industry-sponsored RCTs that compared the Watchman device with anticoagulation alone. One trial reported noninferiority on a composite outcome of stroke, cardiovascular/unexplained death, or systemic embolism after 2 years of follow-up, with continued benefits with the Watchman device after 4 years of follow-up. The second trial did not demonstrate noninferiority for the same composite outcome but did



demonstrate noninferiority of the Watchman device to warfarin for late ischemic stroke and systemic embolization. Patient-level meta-analyses at 5-year follow-up for the 2 trials reported that the Watchman device is noninferior to warfarin on the composite outcome of stroke, systemic embolism, and cardiovascular death. Also, the Watchman was associated with lower rates in major bleeding, particularly hemorrhagic stroke, and mortality over the long term. The evidence also indicates that the Watchman device is efficacious in preventing stroke in the subset of patients with AF who are at increased risk for embolic stroke. When it is determined on an individualized basis that the long-term risk of systemic anticoagulation exceeds the procedural risk of device implantation, the net health outcome will be improved. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have AF who are at increased risk for embolic stroke who receive a percutaneous LAAC device other than the Watchman device (eg, the Lariat or Amplatzer), the evidence includes uncontrolled case series. Relevant outcomes are overall survival, morbid events, and treatment-related morbidity. Case series of these devices have reported high procedural success, but also numerous complications. Also, these devices do not have Food and Drug Administration approval for LAAC. The evidence is insufficient to determine the effects of the technology on health outcomes.

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this policy are listed in [Table 3](#).

Table 3. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
Ongoing			
NCT02039167	WATCH Bleeding Episodes After Left Atrial Appendage Occlusion Versus Usual Care in Patients With Atrial Fibrillation and Severe to End-stage Chronic Kidney Disease (WatchAFIB in CKD)	300	Jun 2017 (ongoing)
NCT03276169	Left Atrial Function Changes after Left Atrial Appendage Closure in Patients with Persistent Atrial Fibrillation	105	Nov 2019
NCT02513797^a	aMAZE Study: LAA Ligation with the LARIAT Suture	600	Dec 2019



NCT No.	Trial Name	Planned Enrollment	Completion Date
	Delivery System as Adjunctive to Pulmonary Vein Isolation for Persistent Atrial Fibrillation (aMAZE)		
NCT02426944	Left Atrial Appendage Closure vs Novel Anticoagulation Agents in Atrial Fibrillation	400	May 2020
NCT02879448	AMPLATZER™ Amulet™ Left Atrial Appendage Occluder Randomized Controlled Trial	1600	Dec 2023
NCT03302494 ^a	WAVECrest Vs. Watchman Transseptal LAA Closure to Reduce AF-Mediated Stroke 2 (WAVECREST2)	1250	Dec 2025
Unpublished			
NCT01118299	AMPLATZER Cardiac Plug Clinical Trial	3000	Not approved/cleared
NCT01182441 ^a	Evaluation of the Watchman LAA closure device in patients with atrial fibrillation versus long term warfarin therapy	475	Aug 2017 (unknown)

NCT: national clinical trial

^a indicates industry-sponsored study

Clinical Input Received from Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

In response to requests, input was received from 1 physician specialty society (2 responses) and 4 academic medical centers, one of which provided 4 responses, for a total of 8 responses, while this policy was under review in 2015. Input generally supported the use of a left atrial appendage closure device approved by the Food and Drug Administration for patients with an increased risk of stroke and systemic embolism based on CHADS₂ or CHA₂DS₂-VASc score. Systemic anticoagulation therapy was recommended, but the long-term risks of systemic anticoagulation outweigh the risks of the device implantation.



Practice Guidelines and Position Statements

Guideline Comparison

Andrade et al (2017) provided the following summary (see [Table 4](#)) comparing guidelines by American, Canadian, and European societies on left atrial appendage exclusion and closure for the management of atrial fibrillation.⁵³

Table 4. Comparison of American, Canadian, and European Guidelines on LAA Exclusion/Closure

Procedure	AHA/ACC/HRS	CCS	ESC
Surgical LAA closure (excision or obliteration of LAA)	May be considered in patients undergoing cardiac surgery (IIb)	Should be considered as part of surgical ablation of AF associated with mitral, aortic valve, or coronary artery bypass surgery	May be considered in patients undergoing cardiac surgery (IIb) More data are needed to confirm safety and efficacy of thoroscopic exclusion
Percutaneous LAA exclusion	No recommendation	Not to be used, except in research or in systematically documented use protocols in patients at high risk of stroke (CHADS ₂ ≥2) and antithrombotic therapy precluded	May be considered in patients with contraindications for long term anticoagulant treatment (IIb)

Adapted from Andrade et al (2017).⁵³

ACC: American College of Cardiology; AF: atrial fibrillation; AHA: American Heart Association; CCS: Canadian Cardiovascular Society; CHADS₂: Congestive Heart Failure, Hypertension, Age, Diabetes, Stroke/Transient Ischemic Attack; ESC: European Society of Cardiology; HRS: Heart Rhythm Society; LAA: left atrial appendage

Medicare National Coverage

As of 2016, the Centers for Medicare & Medicaid Services has a national coverage determination (NCD) under coverage with evidence development for percutaneous LAAC in AF, as follows⁵⁴:

“LAAC devices are covered when the device has received Food and Drug Administration (FDA) Premarket Approval (PMA) for that device’s FDA-approved indication and meet all of the conditions specified below:

“The patient must have:



- “A CHADS2 score ≥ 2 (Congestive heart failure, Hypertension, Age > 75 , Diabetes, Stroke/transient ischemia attack/thromboembolism) or CHA2DS2-VASc score ≥ 3 (Congestive heart failure, Hypertension, Age ≥ 65 , Diabetes, Stroke/transient ischemia attack/thromboembolism, Vascular disease, Sex category)
- “A formal shared decision making interaction with an independent non-interventional physician using an evidence-based decision tool on oral anticoagulation in patients with NVAf [nonvalvular atrial fibrillation] prior to LAAC. Additionally, the shared decision making interaction must be documented in the medical record
- “A suitability for short-term warfarin but deemed unable to take long-term oral anticoagulation following the conclusion of shared decision making, as LAAC is only covered as a second line therapy to oral anticoagulants. The patient (preoperatively and postoperatively) is under the care of a cohesive, multidisciplinary team (MDT) of medical professionals. The procedure must be furnished in a hospital with an established structural heart disease (SHD) and/or electrophysiology (EP) program

“The procedure must be performed by an interventional cardiologist(s), electrophysiologist(s), or cardiovascular surgeon(s) that meets the following criteria:

- “Has received training prescribed by the manufacturer on the safe and effective use of the device prior to performing LAAC; and,
- “Has performed ≥ 25 interventional cardiac procedures that involve transseptal puncture through an intact septum; and,
- “Continues to perform ≥ 25 interventional cardiac procedures that involve transseptal puncture through an intact septum, of which at least 12 are LAAC, over a 2-year period.”

Patients must be enrolled in approved registries that track outcomes for procedures and devices.

Regulatory Status

In 2002, the PLAATO system (ev3 Endovascular) was the first device to be approved by FDA for LAA occlusion. The device was discontinued in 2007 for commercial reasons, and intellectual property was sold to manufacturers of the Watchman system.

In 2015, the Watchman™ Left Atrial Appendage Closure Technology (Boston Scientific) was approved by the FDA through the premarket approval process by the Left Atrial Appendage



Versus Warfarin Therapy for Prevention of Stroke in Patients with Atrial Fibrillation (PROTECT-AF) randomized controlled trial.⁴ This device is indicated to reduce the risk of thromboembolism from the LAA in patients with nonvalvular AF who:

- Are at increased risk for stroke and systemic embolism based on CHADS₂ or CHA₂DS₂-VASc scores and are recommended for anticoagulation therapy;
- Are deemed by their physicians to be suitable for warfarin; and
- Have an appropriate rationale to seek a nonpharmacologic alternative to warfarin, taking into account the safety and effectiveness of the device compared to warfarin.

FDA product code: NGV.

Several other devices are being evaluated for LAA occlusion but are not approved in the United States for percutaneous LAAC. In 2006, the Lariat® Loop Applicator device (SentreHEART), a suture delivery system, was cleared for marketing by FDA through the 510(k) process. The intended use is to facilitate suture placement and knot tying in surgical applications where soft tissues are being approximated or ligated with a pretied polyester suture. The Amplatzer Amulet® device (St. Jude Medical) and WaveCrest® (Coherex Medical) have CE approval in Europe for LAAC but are not currently approved in the United States for this indication.

References

1. You JJ, Singer DE, Howard PA, et al. Antithrombotic therapy for atrial fibrillation: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. Chest. Feb 2012;141(2 Suppl):e531S-575S. PMID 22315271
2. January CT, Wann LS, Alpert JS, et al. 2014 AHA/ACC/HRS guideline for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the Heart Rhythm Society. J Am Coll Cardiol. Dec 2 2014;64(21):e1-76. PMID 24685669
3. Pisters R, Lane DA, Nieuwlaat R, et al. A novel user-friendly score (HAS-BLED) to assess 1-year risk of major bleeding in patients with atrial fibrillation: the Euro Heart Survey. Chest. Nov 2010;138(5):1093-1100. PMID 20299623
4. Food and Drug Administration. Approval Letter: WATCHMAN LAA Closure Technology. 2015; http://www.accessdata.fda.gov/cdrh_docs/pdf13/p130013a.pdf Accessed July 2018.
5. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Percutaneous left atrial appendage closure therapy for prevention of stroke. TEC Assessments 2014;Volume 29:Tab 5.
6. Bode WD, Patel N, Gehi AK. Left atrial appendage occlusion for prevention of stroke in nonvalvular atrial fibrillation: a meta-analysis. J Interv Card Electrophysiol. Jun 2015;43(1):79-89. PMID 25711953



7. Briceno DF, Villablanca P, Cyrille N, et al. Left atrial appendage occlusion device and novel oral anticoagulants versus warfarin for stroke prevention in nonvalvular atrial fibrillation: systematic review and meta-analysis of randomized controlled trials. *Circ Arrhythm Electrophysiol*. Oct 2015;8(5):1057-1064. PMID 26226997
8. Holmes DR, Jr., Doshi SK, Kar S, et al. Left atrial appendage closure as an alternative to warfarin for stroke prevention in atrial fibrillation: a patient-level meta-analysis. *J Am Coll Cardiol*. Jun 23 2015;65(24):2614-2623. PMID 26088300
9. Li X, Wen SN, Li SN, et al. Over 1-year efficacy and safety of left atrial appendage occlusion versus novel oral anticoagulants for stroke prevention in atrial fibrillation: A systematic review and meta-analysis of randomized controlled trials and observational studies. *Heart Rhythm*. Jun 2016;13(6):1203-1214. PMID 26724488
10. Lip GY, Lane DA. Stroke prevention in atrial fibrillation: a systematic review. *JAMA*. May 19 2015;313(19):1950-1962. PMID 25988464
11. Price MJ, Reddy VY, Valderrabano M, et al. Bleeding outcomes after left atrial appendage closure compared with long-term warfarin: a pooled, patient-level analysis of the WATCHMAN randomized trial experience. *JACC Cardiovasc Interv*. Dec 28 2015;8(15):1925-1932. PMID 26627989
12. Noelck N, Papak J, Freeman M, et al. Effectiveness of left atrial appendage exclusion procedures to reduce the risk of stroke: a systematic review of the evidence. *Circ Cardiovasc Qual Outcomes*. Jul 2016;9(4):395-405. PMID 27407055
13. Sahay S, Nombela-Franco L, Rodes-Cabau J, et al. Efficacy and safety of left atrial appendage closure versus medical treatment in atrial fibrillation: a network meta-analysis from randomised trials. *Heart*. Jan 15 2017;103(2):139-147. PMID 27587437
14. Wei Z, Zhang X, Wu H, et al. A meta-analysis for efficacy and safety evaluation of transcatheter left atrial appendage occlusion in patients with nonvalvular atrial fibrillation. *Medicine (Baltimore)*. Aug 2016;95(31):e4382. PMID 27495048
15. Tereshchenko LG, Henrikson CA, Cigarroa J, et al. Comparative effectiveness of interventions for stroke prevention in atrial fibrillation: a network meta-analysis. *J Am Heart Assoc*. May 20 2016;5(5). PMID 27207998
16. Reddy VY, Doshi SK, Kar S, et al. 5-year outcomes after left atrial appendage closure: from the PREVAIL and PROTECT AF Trials. *J Am Coll Cardiol*. Dec 19 2017;70(24):2964-2975. PMID 29103847
17. Bajaj NS, Kalra R, Patel N, et al. Comparison of approaches for stroke prophylaxis in patients with non-valvular atrial fibrillation: network meta-analyses of randomized controlled trials. *PLoS One*. 2016;11(10):e0163608. PMID 27706224
18. Hanif H, Belley-Cote EP, Alotaibi A, et al. Left atrial appendage occlusion for stroke prevention in patients with atrial fibrillation: a systematic review and network meta-analysis of randomized controlled trials. *J Cardiovasc Surg (Torino)*. Feb 2018;59(1):128-139. PMID 28215062
19. Holmes DR, Reddy VY, Turi ZG, et al. Percutaneous closure of the left atrial appendage versus warfarin therapy for prevention of stroke in patients with atrial fibrillation: a randomised non-inferiority trial. *Lancet*. Aug 15 2009;374(9689):534-542. PMID 19683639
20. Reddy VY, Doshi SK, Sievert H, et al. Percutaneous left atrial appendage closure for stroke prophylaxis in patients with atrial fibrillation: 2.3-year follow-up of the PROTECT AF (Watchman Left Atrial Appendage System for Embolic Protection in Patients With Atrial Fibrillation) trial. *Circulation*. Feb 12 2013;127(6):720-729. PMID 23325525
21. Reddy VY, Sievert H, Halperin J, et al. Percutaneous left atrial appendage closure vs warfarin for atrial fibrillation: a randomized clinical trial. *JAMA*. Nov 19 2014;312(19):1988-1998. PMID 25399274
22. Alli O, Doshi S, Kar S, et al. Quality of life assessment in the randomized PROTECT AF (Percutaneous Closure of the Left Atrial Appendage Versus Warfarin Therapy for Prevention of Stroke in Patients With Atrial Fibrillation) trial of patients at risk for stroke with nonvalvular atrial fibrillation. *J Am Coll Cardiol*. Apr 30 2013;61(17):1790-1798. PMID 23500276
23. Holmes DR, Jr., Kar S, Price MJ, et al. Prospective randomized evaluation of the Watchman Left Atrial Appendage Closure device in patients with atrial fibrillation versus long-term warfarin therapy: the PREVAIL trial. *J Am Coll Cardiol*. Jul 8 2014;64(1):1-12. PMID 24998121
24. Lam YY, Yip GW, Yu CM, et al. Left atrial appendage closure with AMPLATZER cardiac plug for stroke prevention in atrial fibrillation: initial Asia-Pacific experience. *Catheter Cardiovasc Interv*. Apr 1 2012;79(5):794-800. PMID 21542102



25. Montenegro MJ, Quintella EF, Damonte A, et al. Percutaneous occlusion of left atrial appendage with the Amplatzer Cardiac Plug™ in atrial fibrillation. *Arq Bras Cardiol.* Jan 31 2012;98(2):143-150. PMID 22286325
26. Park JW, Bethencourt A, Sievert H, et al. Left atrial appendage closure with Amplatzer cardiac plug in atrial fibrillation: initial European experience. *Catheter Cardiovasc Interv.* Apr 1 2011;77(5):700-706. PMID 20824765
27. Reddy VY, Holmes D, Doshi SK, et al. Safety of percutaneous left atrial appendage closure: results from the Watchman Left Atrial Appendage System for Embolic Protection in Patients with AF (PROTECT AF) clinical trial and the Continued Access Registry. *Circulation.* Feb 1 2011;123(4):417-424. PMID 21242484
28. Swaans MJ, Post MC, Rensing BJ, et al. Percutaneous left atrial appendage closure for stroke prevention in atrial fibrillation. *Neth Heart J.* Jan 10 2012;20(4):161-166. PMID 22231152
29. Reddy VY, Möbius-Winkler S, Miller MA, et al. Left atrial appendage closure with the Watchman device in patients with a contraindication for oral anticoagulation: the ASAP study (ASA Plavix Feasibility Study With Watchman Left Atrial Appendage Closure Technology). *J Am Coll Cardiol.* Jun 25 2013;61(25):2551-2556. PMID 23583249
30. Chun KR, Bordignon S, Urban V, et al. Left atrial appendage closure followed by 6 weeks of antithrombotic therapy: a prospective single-center experience. *Heart Rhythm.* Dec 2013;10(12):1792-1799. PMID 23973952
31. Boersma LV, Schmidt B, Betts TR, et al. Implant success and safety of left atrial appendage closure with the WATCHMAN device: peri-procedural outcomes from the EWOLUTION registry. *Eur Heart J.* Aug 2016;37(31):2465-2474. PMID 26822918
32. Chatterjee S, Herrmann HC, Wilensky RL, et al. Safety and procedural success of left atrial appendage exclusion with the Lariat device: a systematic review of published reports and analytic review of the FDA MAUDE Database. *JAMA Intern Med.* Jul 2015;175(7):1104-1109. PMID 25938303
33. Lakkireddy D, Afzal MR, Lee RJ, et al. Short and long-term outcomes of percutaneous left atrial appendage suture ligation: Results from a US multicenter evaluation. *Heart Rhythm.* May 2016;13(5):1030-1036. PMID 26872554
34. Price MJ, Gibson DN, Yakubov SJ, et al. Early safety and efficacy of percutaneous left atrial appendage suture ligation: results from the U.S. transcatheter LAA ligation consortium. *J Am Coll Cardiol.* Aug 12 2014;64(6):565-572. PMID 25104525
35. Bartus K, Han FT, Bednarek J, et al. Percutaneous left atrial appendage suture ligation using the LARIAT device in patients with atrial fibrillation: initial clinical experience. *J Am Coll Cardiol.* Jul 9 2013;62(2):108-118. PMID 23062528
36. Bartus K, Bednarek J, Myc J, et al. Feasibility of closed-chest ligation of the left atrial appendage in humans. *Heart Rhythm.* Feb 2011;8(2):188-193. PMID 21050893
37. Stone D, Byrne T, Pershad A. Early results with the LARIAT device for left atrial appendage exclusion in patients with atrial fibrillation at high risk for stroke and anticoagulation. *Catheter Cardiovasc Interv.* Jul 2015;86(1):121-127. PMID 23765504
38. Massumi A, Chelu MG, Nazeri A, et al. Initial experience with a novel percutaneous left atrial appendage exclusion device in patients with atrial fibrillation, increased stroke risk, and contraindications to anticoagulation. *Am J Cardiol.* Jan 9 2013;111(6):869-873. PMID 23312129
39. Nietlispach F, Gloekler S, Krause R, et al. Amplatzer left atrial appendage occlusion: Single center 10-year experience. *Catheter Cardiovasc Interv.* Feb 14 2013;82(2):283-289. PMID 23412815
40. Kefer J, Vermeersch P, Budts W, et al. Transcatheter left atrial appendage closure for stroke prevention in atrial fibrillation with Amplatzer cardiac plug: the Belgian Registry. *Acta Cardiol.* Dec 2013;68(6):551-558. PMID 24579432
41. Guerios EE, Schmid M, Gloekler S, et al. Left atrial appendage closure with the Amplatzer cardiac plug in patients with atrial fibrillation. *Arq Bras Cardiol.* Jun 2012;98(6):528-536. PMID 22584492
42. Danna P, Proietti R, Sagone A, et al. Does left atrial appendage closure with a cardiac plug system reduce the stroke risk in nonvalvular atrial fibrillation patients? A single-center case series. *Pacing Clin Electrophysiol.* Mar 2013;36(3):347-353. PMID 23252940
43. Lopez-Minguez JR, Eldoayen-Gragera J, Gonzalez-Fernandez R, et al. Immediate and one-year results in 35 consecutive patients after closure of left atrial appendage with the Amplatzer Cardiac Plug. *Rev Esp Cardiol.* Feb 2013;66(2):90-97. PMID 22939161



44. Streb W, Szymala M, Kukulski T, et al. Percutaneous closure of the left atrial appendage using the Amplatzer Cardiac Plug in patients with atrial fibrillation: evaluation of safety and feasibility. *Kardiol Pol.* Jan 2013;71(1):8-16. PMID 23348528
45. Santoro G, Meucci F, Stolcova M, et al. Percutaneous left atrial appendage occlusion in patients with non-valvular atrial fibrillation: implantation and up to four years follow-up of the AMPLATZER Cardiac Plug. *EuroIntervention.* Feb 20 2016;11(10):1188-1194. PMID 25354761
46. Meerkin D, Butnaru A, Dratva D, et al. Early safety of the Amplatzer Cardiac Plug for left atrial appendage occlusion. *Int J Cardiol.* Oct 9 2013;168(4):3920-3925. PMID 23890886
47. Wiebe J, Bertog S, Franke J, et al. Safety of percutaneous left atrial appendage closure with the Amplatzer cardiac plug in patients with atrial fibrillation and contraindications to anticoagulation. *Catheter Cardiovasc Interv.* Apr 1 2014;83(5):796-802. PMID 24327462
48. Urena M, Rodes-Cabau J, Freixa X, et al. Percutaneous left atrial appendage closure with the AMPLATZER cardiac plug device in patients with nonvalvular atrial fibrillation and contraindications to anticoagulation therapy. *J Am Coll Cardiol.* Jul 9 2013;62(2):96-102. PMID 23665098
49. Gloekler S, Shakir S, Doblies J, et al. Early results of first versus second generation Amplatzer occluders for left atrial appendage closure in patients with atrial fibrillation. *Clin Res Cardiol.* Aug 2015;104(8):656-665. PMID 25736061
50. Bayard YL, Omran H, Neuzil P, et al. PLAATO (Percutaneous Left Atrial Appendage Transcatheter Occlusion) for prevention of cardioembolic stroke in non-anticoagulation eligible atrial fibrillation patients: results from the European PLAATO study. *EuroIntervention.* Jun 2010;6(2):220-226. PMID 20562072
51. Cruz-Gonzalez I, Martin Moreiras J, Garcia E. Thrombus formation after left atrial appendage exclusion using an amplatzer cardiac plug device. *Catheter Cardiovasc Interv.* Nov 15 2011;78(6):970-973. PMID 21523900
52. Viles-Gonzalez JF, Reddy VY, Petru J, et al. Incomplete occlusion of the left atrial appendage with the percutaneous left atrial appendage transcatheter occlusion device is not associated with increased risk of stroke. *J Interv Card Electrophysiol.* Jan 2012;33(1):69-75. PMID 21947786
53. Andrade JG, Macle L, Nattel S, et al. Contemporary atrial fibrillation management: a comparison of the current AHA/ACC/HRS, CCS, and ESC guidelines. *Can J Cardiol.* Aug 2017;33(8):965-976. PMID 28754397

Center for Medicare & Medicaid Services. National Coverage Determination (NCD) for Percutaneous Left Atrial Appendage Closure (LAAC) (20.34). 2016; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=367&ncdver=1&NCAId=281&bc=AAAAAAAAAAAA%3d%3d&>. Accessed July 2018.

History

Date	Comments
06/13/11	Add to Cardiology Section - New medical policy created with literature search; procedure considered investigational.
12/29/11	Code 0281T added.
05/22/12	Replace policy. Policy updated with literature review, references 2-4, 6-9, 11, 12 added. Policy title and policy statements revised to include percutaneous – no other change to policy statement.
09/17/12	Update Coding Section – ICD-10 codes are now effective 10/01/2014.



Date	Comments
05/28/13	Replace policy. Policy updated with literature review through January 2013, references 2, 12-20 added. Policy statement unchanged.
09/23/14	Annual Review. Policy updated with literature review through June 5, 2014. References 1-3, 12-14, 17, 25-29, 31-32, and 34 added. Policy statement unchanged.
12/08/15	Annual Review. Policy updated with literature review through May 29, 2015; references 2-3, 6, 9, 12, 21, 33, and 43 added; clinical input reviewed. An FDA-approved left atrial appendage closure device is considered medically necessary with conditions.
08/01/16	Annual Review, approved July 12, 2016. Policy updated with literature review through April 30, 2016; references 6-7, 9-11, and 25-27 added. Policy statements unchanged.
01/01/17	Coding update, added new code 33340 effective 1/1/17.
07/01/17	Annual Review, approved June 22, 2017. Policy moved into new format. Policy updated with literature review through March 23, 2017; references 13-18 and 55-56 added. Policy statements unchanged.
01/01/18	Removed code CPT code 0281T as it was terminated 1/1/17 and replaced with 33340.
08/01/18	Annual Review, approved July 13, 2018. Policy updated with literature review through March 2018; references 16 and 53 added. PLAATO device removed from the policy statement; statements otherwise unchanged. Removed CPT codes 33999 and 93799.

Disclaimer: This medical policy is a guide in evaluating the medical necessity of a particular service or treatment. The Company adopts policies after careful review of published peer-reviewed scientific literature, national guidelines and local standards of practice. Since medical technology is constantly changing, the Company reserves the right to review and update policies as appropriate. Member contracts differ in their benefits. Always consult the member benefit booklet or contact a member service representative to determine coverage for a specific medical service or supply. CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). ©2018 Premera All Rights Reserved.

Scope: Medical policies are systematically developed guidelines that serve as a resource for Company staff when determining coverage for specific medical procedures, drugs or devices. Coverage for medical services is subject to the limits and conditions of the member benefit plan. Members and their providers should consult the member benefit booklet or contact a customer service representative to determine whether there are any benefit limitations applicable to this service or supply. This medical policy does not apply to Medicare Advantage.



Discrimination is Against the Law

Premera Blue Cross complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex. Premera does not exclude people or treat them differently because of race, color, national origin, age, disability or sex.

Premera:

- Provides free aids and services to people with disabilities to communicate effectively with us, such as:
 - Qualified sign language interpreters
 - Written information in other formats (large print, audio, accessible electronic formats, other formats)
- Provides free language services to people whose primary language is not English, such as:
 - Qualified interpreters
 - Information written in other languages

If you need these services, contact the Civil Rights Coordinator.

If you believe that Premera has failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, or sex, you can file a grievance with:

Civil Rights Coordinator - Complaints and Appeals
PO Box 91102, Seattle, WA 98111
Toll free 855-332-4535, Fax 425-918-5592, TTY 800-842-5357
Email AppealsDepartmentInquiries@Premera.com

You can file a grievance in person or by mail, fax, or email. If you need help filing a grievance, the Civil Rights Coordinator is available to help you.

You can also file a civil rights complaint with the U.S. Department of Health and Human Services, Office for Civil Rights, electronically through the Office for Civil Rights Complaint Portal, available at <https://ocrportal.hhs.gov/ocr/portal/lobby.jsf>, or by mail or phone at: U.S. Department of Health and Human Services
200 Independence Avenue SW, Room 509F, HHH Building
Washington, D.C. 20201, 1-800-368-1019, 800-537-7697 (TDD)
Complaint forms are available at <http://www.hhs.gov/ocr/office/file/index.html>.

Getting Help in Other Languages

This Notice has Important Information. This notice may have important information about your application or coverage through Premera Blue Cross. There may be key dates in this notice. You may need to take action by certain deadlines to keep your health coverage or help with costs. You have the right to get this information and help in your language at no cost. Call 800-722-1471 (TTY: 800-842-5357).

አማርኛ (Amharic):

ይህ ማስታወቂያ አስፈላጊ መረጃ ይዟል። ይህ ማስታወቂያ ስለ ማመልከቻዎ ወይም የ Premera Blue Cross ሽፋን አስፈላጊ መረጃ ሊኖረው ይችላል። በዚህ ማስታወቂያ ውስጥ ቁልፍ ቀናት ሊኖሩ ይችላሉ። የጤና ሽፋንዎን ለመጠበቅና በአስፈላጊ እርዳታ ለማግኘት በተውሰኑ የጊዜ ገደቦች እርምጃ መውሰድ ይገባዎት ይሆናል። ይህን መረጃ እንዲያገኙ እና የለምንም ክፍያ በቋንቋዎ እርዳታ እንዲያገኙ መሰታወቅ አለዎት። በስልክ ቁጥር 800-722-1471 (TTY: 800-842-5357) ይደውሉ።

العربية (Arabic):

يحتوي هذا الإشعار على معلومات هامة. قد يحتوي هذا الإشعار على معلومات مهمة بخصوص طلبك أو التغطية التي تزيد الحصول عليها من خلال Premera Blue Cross. قد تكون هناك تواريخ مهمة في هذا الإشعار. وقد تحتاج لاتخاذ إجراء في تاريخ معينه للحفاظ على تغطيتك الصحية أو للمساعدة في دفع التكاليف. يحق لك الحصول على هذه المعلومات والمساعدة بلغتك دون تكبد أية تكلفة. اتصل بـ 800-722-1471 (TTY: 800-842-5357)

中文 (Chinese):

本通知有重要的訊息。本通知可能有關於您透過 Premera Blue Cross 提交的申請或保險的重要訊息。本通知內可能有重要日期。您可能需要在截止日期之前採取行動，以保留您的健康保險或者費用補貼。您有權利免費以您的母語得到本訊息和幫助。請撥電話 800-722-1471 (TTY: 800-842-5357)。

Oromoo (Cushite):

Beeksisni kun odeeffannoo barbaachisaa qaba. Beeksisti kun sagantaa yookan karaa Premera Blue Cross tiin tajaajila keessan ilaalchisee odeeffannoo barbaachisaa qabaachuu danda'a. Guyyaawwan murteessaa ta'an beeksisa kana keessatti ilaalaa. Tarii kaffaltiidhaan deeggaramuuf yookan tajaajila fayyaa keessaniif guyyaa dhumaa irratti wanti raawwattan jiraachuu danda'a. Kaffaltii irraa bilisa haala ta'een afaan keessaniin odeeffannoo argachuu fi deeggarsa argachuuf mirga ni qabaattu. Lakkoofsa bilbilaa 800-722-1471 (TTY: 800-842-5357) tii bilbilaa.

Français (French):

Cet avis a d'importantes informations. Cet avis peut avoir d'importantes informations sur votre demande ou la couverture par l'intermédiaire de Premera Blue Cross. Le présent avis peut contenir des dates clés. Vous devez peut-être prendre des mesures par certains délais pour maintenir votre couverture de santé ou d'aide avec les coûts. Vous avez le droit d'obtenir cette information et de l'aide dans votre langue à aucun coût. Appelez le 800-722-1471 (TTY: 800-842-5357).

Kreyòl ayisyen (Creole):

Avi sila a gen Enfòmasyon Enpòtan ladann. Avi sila a kapab genyen enfòmasyon enpòtan konsènan aplikasyon w lan oswa konsènan kouvèti asirans lan atravè Premera Blue Cross. Kapab genyen dat ki enpòtan nan avi sila a. Ou ka gen pou pran kèk aksyon avan sèten dat limit pou ka kenbe kouvèti asirans sante w la oswa pou yo ka ede w avèk depans yo. Se dwa w pou resewva enfòmasyon sa a ak asistans nan lang ou pale a, san ou pa gen pou peye pou sa. Rele nan 800-722-1471 (TTY: 800-842-5357).

Deutsche (German):

Diese Benachrichtigung enthält wichtige Informationen. Diese Benachrichtigung enthält unter Umständen wichtige Informationen bezüglich Ihres Antrags auf Krankenversicherungsschutz durch Premera Blue Cross. Suchen Sie nach eventuellen wichtigen Terminen in dieser Benachrichtigung. Sie könnten bis zu bestimmten Stichtagen handeln müssen, um Ihren Krankenversicherungsschutz oder Hilfe mit den Kosten zu behalten. Sie haben das Recht, kostenlose Hilfe und Informationen in Ihrer Sprache zu erhalten. Rufen Sie an unter 800-722-1471 (TTY: 800-842-5357).

Hmoob (Hmong):

Tsab ntawv tshaj xo no muaj cov ntshiab lus tseem ceeb. Tej zaum tsab ntawv tshaj xo no muaj cov ntshiab lus tseem ceeb txog koj daim ntawv thov kev pab los yog koj qhov kev pab cuam hnuv ntawm Premera Blue Cross. Tej zaum muaj cov hnuv tseem ceeb uas sau rau hauv daim ntawv no. Tej zaum koj kuj yuav tau ua qee yam uas peb kom koj ua tsis pub dhau cov caij nyoog uas teev tseg rau hauv daim ntawv no mas koj thiaj yuav tau txais kev pab cuam kho mob los yog kev pab them tej nqi kho mob ntawd. Koj muaj cai kom lawv muab cov ntshiab lus no uas tau muab sau ua koj hom lus pub dawb rau koj. Hu rau 800-722-1471 (TTY: 800-842-5357).

Iloko (Ilocano):

Daytoy a Pakdaar ket naglaon iti Napateg nga Impormasion. Daytoy a pakdaar mabalin nga adda ket naglaon iti napateg nga impormasion maipanggep iti aplikasyonyo wenna coverage babaen iti Premera Blue Cross. Daytoy ket mabalin dagiti importante a petsa iti daytoy a pakdaar. Mabalin nga adda rumbeng nga aramidenyo nga addang sakbay dagiti partikular a naituding nga aldaw tapno mapagtalinaedyo ti coverage ti salun-atyto wenna tulong kadagiti gastos. Adda karbenganyo a mangala iti daytoy nga impormasion ken tulong iti bukodyo a pagsasao nga awan ti bayadanyo. Tumawag iti numero nga 800-722-1471 (TTY: 800-842-5357).

Italiano (Italian):

Questo avviso contiene informazioni importanti. Questo avviso può contenere informazioni importanti sulla tua domanda o copertura attraverso Premera Blue Cross. Potrebbero esserci date chiave in questo avviso. Potrebbe essere necessario un tuo intervento entro una scadenza determinata per consentirti di mantenere la tua copertura o sovvenzione. Hai il diritto di ottenere queste informazioni e assistenza nella tua lingua gratuitamente. Chiama 800-722-1471 (TTY: 800-842-5357).

日本語 (Japanese):

この通知には重要な情報が含まれています。この通知には、Premera Blue Cross の申請または補償範囲に関する重要な情報が含まれている場合があります。この通知に記載されている可能性がある重要な日付をご確認ください。健康保険や有料サポートを維持するには、特定の期日までに行動を取らなければならない場合があります。ご希望の言語による情報とサポートが無料で提供されます。800-722-1471 (TTY: 800-842-5357)までお電話ください。

한국어 (Korean):

본 통지서에는 중요한 정보가 들어 있습니다. 즉 이 통지서는 귀하의 신청에 관하여 그리고 Premera Blue Cross 를 통한 커버리지에 관한 정보를 포함하고 있을 수 있습니다. 본 통지서에는 핵심이 되는 날짜들이 있을 수 있습니다. 귀하의 건강 커버리지를 계속 유지하거나 비용을 절감하기 위해서 일정한 마감일까지 조치를 취해야 할 필요가 있을 수 있습니다. 귀하의 이러한 정보와 도움을 귀하의 언어로 비용 부담없이 얻을 수 있는 권리가 있습니다. 800-722-1471 (TTY: 800-842-5357) 로 전화하십시오.

ລາວ (Lao):

ແຈງການນີ້ມີຂໍ້ມູນສໍາຄັນ. ແຈງການນີ້ອາດຈະມີຂໍ້ມູນສໍາຄັນກ່ຽວກັບຄໍາຮ້ອງສະໝັກ ຫຼື ຄວາມຄົມຄອງປະກັນໄພຂອງທ່ານຜ່ານ Premera Blue Cross. ອາດຈະມີວັນທີ່ສໍາຄັນໃນແຈງການນີ້. ທ່ານອາດຈະຈໍາເປັນຕ້ອງດໍາເນີນການຕາມກຳນົດ ເວລາສະເພາະເພື່ອຮັກສາຄວາມຄົມຄອງປະກັນສະພາບ ຫຼື ຄວາມຊ່ວຍເຫຼືອເວັ້ນເວີ້ ຄ່າໃຊ້ຈ່າຍຂອງທ່ານໄດ້. ທ່ານມີສິດໄດ້ຮັບຂໍ້ມູນນີ້ ແລະ ຄວາມຊ່ວຍເຫຼືອເປັນພາສາຂອງທ່ານໂດຍບໍ່ເສຍຄ່າ. ໃຫ້ໃບທາ 800-722-1471 (TTY: 800-842-5357).

ភាសាខ្មែរ (Khmer):

សេចក្តីជូនដំណឹងនេះមានព័ត៌មានយ៉ាងសំខាន់។ សេចក្តីជូនដំណឹងនេះប្រហែលជាមានព័ត៌មានយ៉ាងសំខាន់អំពីទម្រង់បែបបទ ឬការរៀបចំរបស់អ្នកកាមរយ: Premera Blue Cross ។ ប្រហែលជាមាន កាលបរិច្ឆេទសំខាន់នៅក្នុងសេចក្តីជូនដំណឹងនេះ។ អ្នកប្រហែលជាត្រូវការបញ្ជាក់សមត្ថភាព ដល់កំណត់ថ្លៃជាតំបន់នានា ដើម្បីនឹងរក្សាទុកការធានារ៉ាប់រងអន្តរជាតិរបស់អ្នក ឬប្រាក់ដុល្លារចេញថ្លៃ។ អ្នកមានសិទ្ធិទទួលព័ត៌មាននេះ និងដុល្លារនៅក្នុងភាសារបស់អ្នកដោយមិនអស់លុយឡើយ។ សូមទូរស័ព្ទ 800-722-1471 (TTY: 800-842-5357)។

ਪੰਜਾਬੀ (Punjabi):

ਇਸ ਨੋਟਿਸ ਵਿਚ ਖਾਸ ਜਾਣਕਾਰੀ ਹੈ. ਇਸ ਨੋਟਿਸ ਵਿਚ Premera Blue Cross ਵਲੋਂ ਤੁਹਾਡੀ ਕਵਰੇਜ ਅਤੇ ਅਰਜੀ ਬਾਰੇ ਮਹੱਤਵਪੂਰਨ ਜਾਣਕਾਰੀ ਹੋ ਸਕਦੀ ਹੈ . ਇਸ ਨੋਟਿਸ ਨਵ ਖਾਸ ਤਾਰੀਖਾਂ ਹੋ ਸਕਦੀਆਂ ਹਨ. ਜੇਕਰ ਤੁਸੀਂ ਜਸਰਤ ਕਵਰੇਜ ਰਿੱਖਣੀ ਹੋਵੇ ਜਾਂ ਓਸ ਦੀ ਲਾਗਤ ਜਵਿੱਚ ਮਦਦ ਦੇ ਇਛੁੱਕ ਹੋ ਤਾਂ ਤੁਹਾਨੂੰ ਅੰਤਮ ਤਾਰੀਖ ਤੋਂ ਪਹਿਲਾਂ ਢੁੱਝ ਖਾਸ ਕਦਮ ਚੁੱਕਣ ਦੀ ਲੋੜ ਹੋ ਸਕਦੀ ਹੈ ,ਤੁਹਾਨੂੰ ਮੁਫਤ ਵਿੱਚ ਤੋਂ ਅਪਣੀ ਭਾਸ਼ਾ ਵਿੱਚ ਜਾਣਕਾਰੀ ਅਤੇ ਮਦਦ ਪ੍ਰਾਪਤ ਕਰਨ ਦਾ ਅਧਿਕਾਰ ਹੈ ,ਕਾਲ 800-722-1471 (TTY: 800-842-5357).

فارسی (Farsi):

این اعلامیه حاوی اطلاعات مهم میباشد. این اعلامیه ممکن است حاوی اطلاعات مهم درباره فرم تقاضا و یا پوشش بیمه ای شما از طریق Premera Blue Cross باشد. به تاریخ های مهم در این اعلامیه توجه نمایید. شما ممکن است برای حفظ پوشش بیمه تان یا کمک در پرداخت هزینه های درمانی تان، به تاریخ های مشخصی برای انجام کارهای خاصی احتیاج داشته باشید. شما حق این را دارید که این اطلاعات و کمک را به زبان خود به طور رایگان دریافت نمایید. برای کسب اطلاعات با شماره 800-722-1471 (کلیران TTY تماس باشماره 800-842-5357) تماس برقرار نمایید.

Polskie (Polish):

To ogłoszenie może zawierać ważne informacje. To ogłoszenie może zawierać ważne informacje odnośnie Państwa wniosku lub zakresu świadczeń poprzez Premera Blue Cross. Prosimy zwrócić uwagę na kluczowe daty, które mogą być zawarte w tym ogłoszeniu aby nie przekroczyć terminów w przypadku utrzymania polisy ubezpieczeniowej lub pomocy związanej z kosztami. Macie Państwo prawo do bezpłatnej informacji we własnym języku. Zadzwońcie pod 800-722-1471 (TTY: 800-842-5357).

Português (Portuguese):

Este aviso contém informações importantes. Este aviso poderá conter informações importantes a respeito de sua aplicação ou cobertura por meio do Premera Blue Cross. Poderão existir datas importantes neste aviso. Talvez seja necessário que você tome providências dentro de determinados prazos para manter sua cobertura de saúde ou ajuda de custos. Você tem o direito de obter esta informação e ajuda em seu idioma e sem custos. Ligue para 800-722-1471 (TTY: 800-842-5357).

Română (Romanian):

Prezenta notificare conține informații importante privind cererea sau acoperirea asigurării dumneavoastră de sănătate prin Premera Blue Cross. Pot exista date cheie în această notificare. Este posibil să fie nevoie să acționați până la anumite termene limită pentru a vă menține acoperirea asigurării de sănătate sau asistența provizorie la costuri. Aveți dreptul de a obține gratuit aceste informații și ajutor în limba dumneavoastră. Sunați la 800-722-1471 (TTY: 800-842-5357).

Русский (Russian):

Настоящее уведомление содержит важную информацию. Это уведомление может содержать важную информацию о вашем заявлении или страховом покрытии через Premera Blue Cross. В настоящем уведомлении могут быть указаны ключевые даты. Вам, возможно, потребуется принять меры к определенным предельным срокам для сохранения страхового покрытия или помощи с расходами. Вы имеете право на бесплатное получение этой информации и помощь на вашем языке. Звоните по телефону 800-722-1471 (TTY: 800-842-5357).

Fa'asamoa (Samoan):

Atonu ua iai i lenei fa'asilasilaga ni fa'amatalaga e sili ona taua e tatau ona e malamalama i ai. O lenei fa'asilasilaga o se fesoasoani e fa'amatala atili i ai i le tulaga o le polokalame, Premera Blue Cross, ua e tau fia maua atu i ai. Fa'amolemole, ia e iloilo fa'alelei i aso fa'apitoa olo'o iai i lenei fa'asilasilaga taua. Masalo o le'a iai ni feau e tatau ona e faia ao le'i aulia le aso ua ta'ua i lenei fa'asilasilaga ina ia e iai pea ma maua fesoasoani mai ai i le polokalame a le Malo olo'o e iai i ai. Olo'o iai iate oe le aia tatau e maua atu i lenei fa'asilasilaga ma lenei fa'matalaga i legagana e te malamalama i ai aunoa ma se togiga tupe. Vili atu i le telefoni 800-722-1471 (TTY: 800-842-5357).

Español (Spanish):

Este Aviso contiene información importante. Es posible que este aviso contenga información importante acerca de su solicitud o cobertura a través de Premera Blue Cross. Es posible que haya fechas clave en este aviso. Es posible que deba tomar alguna medida antes de determinadas fechas para mantener su cobertura médica o ayuda con los costos. Usted tiene derecho a recibir esta información y ayuda en su idioma sin costo alguno. Llame al 800-722-1471 (TTY: 800-842-5357).

Tagalog (Tagalog):

Ang Paunawa na ito ay naglalaman ng mahalagang impormasyon tungkol sa iyong aplikasyon o pagsakop sa pamamagitan ng Premera Blue Cross. Maaaring may mga mahalagang petsa dito sa paunawa. Maaring mangailangan ka na magsagawa ng hakbang sa ilang mga itinakdang panahon upang mapanatili ang iyong pagsakop sa kalusugan o tulong na walang gastos. May karapatan ka na makakuha ng ganiitong impormasyon at tulong sa iyong wika ng walang gastos. Tumawag sa 800-722-1471 (TTY: 800-842-5357).

ไทย (Thai):

ประกาศนี้มีข้อมูลสำคัญ ประกาศนี้อาจมีข้อมูลที่สำคัญเกี่ยวกับกาการสมัครหรือขอบเขตประกันสุขภาพของคุณผ่าน Premera Blue Cross และอาจมีกำหนดการในประกาศนี้ คุณอาจจะต้องดำเนินการภายในกำหนดระยะเวลาที่แน่นอนเพื่อจะรักษาการประกันสุขภาพของคุณหรือการช่วยเหลือที่มีค่าใช้จ่าย คุณมีสิทธิที่จะได้รับข้อมูลและความช่วยเหลือนี้ในภาษาของคุณโดยไม่มีค่าใช้จ่าย โทร 800-722-1471 (TTY: 800-842-5357)

Український (Ukrainian):

Це повідомлення містить важливу інформацію. Це повідомлення може містити важливу інформацію про Ваше звернення щодо страховального покриття через Premera Blue Cross. Зверніть увагу на ключові дати, які можуть бути вказані у цьому повідомленні. Існує імовірність того, що Вам треба буде здійснити певні кроки у конкретні кінцеві строки для того, щоб зберегти Ваше медичне страхування або отримати фінансову допомогу. У Вас є право на отримання цієї інформації та допомоги безкоштовно на Вашій рідній мові. Дзвоніть за номером телефону 800-722-1471 (TTY: 800-842-5357).

Tiếng Việt (Vietnamese):

Thông báo này cung cấp thông tin quan trọng. Thông báo này có thông tin quan trọng về đơn xin tham gia hoặc hợp đồng bảo hiểm của quý vị qua chương trình Premera Blue Cross. Xin xem ngày quan trọng trong thông báo này. Quý vị có thể phải thực hiện theo thông báo đúng trong thời hạn để duy trì bảo hiểm sức khỏe hoặc được trợ giúp thêm về chi phí. Quý vị có quyền được biết thông tin này và được trợ giúp bằng ngôn ngữ của mình miễn phí. Xin gọi số 800-722-1471 (TTY: 800-842-5357).