

MEDICAL POLICY – 6.01.25

Percutaneous Vertebroplasty and Sacroplasty

BCBSA Ref. Policy: 6.01.25

Effective Date: July 1, 2018

Last Revised: June 22, 2018


Replaces: 6.01.520

RELATED MEDICAL POLICIES:

6.01.38 Percutaneous Balloon Kyphoplasty and Mechanical Vertebral Augmentation

Select a hyperlink below to be directed to that section.

[POLICY CRITERIA](#) | [DOCUMENTATION REQUIREMENTS](#) | [CODING](#)
[RELATED INFORMATION](#) | [EVIDENCE REVIEW](#) | [REFERENCES](#) | [HISTORY](#)

 Clicking this icon returns you to the hyperlinks menu above.

Introduction

Osteoporosis or cancer in the bones can cause the vertebrae (the bone in the spine) to weaken. They may become so weak that they collapse. This is known as a compression fracture. The collapse usually happens at the front side of the vertebra, creating a vertebra that looks a bit like a wedge. Percutaneous vertebroplasty is a non-surgical procedure to stabilize a spinal compression fracture. A hollow needle is inserted through the skin and into the damaged bone. Bone cement is then injected into the bone. This policy describes when this procedure may be considered medically necessary. Percutaneous sacroplasty is a similar procedure, but the bone cement is placed in the sacrum. The sacrum is the bone at the bottom of the spine and forms the back of the pelvis. Using this technique for the sacrum is investigational. There are not yet enough medical studies to show whether percutaneous sacroplasty is effective.

Note: The Introduction section is for your general knowledge and is not to be taken as policy coverage criteria. The rest of the policy uses specific words and concepts familiar to medical professionals. It is intended for providers. A provider can be a person, such as a doctor, nurse, psychologist, or dentist. A provider also can be a place where medical care is given, like a hospital, clinic, or lab. This policy informs them about when a service may be covered.

Policy Coverage Criteria

Service	Medical Necessity
<p>Percutaneous vertebroplasty</p>	<p>Percutaneous vertebroplasty may be considered medically necessary for the treatment of:</p> <ul style="list-style-type: none"> • Symptomatic osteoporotic vertebral fractures that have failed to respond to conservative treatment (eg, analgesics, physical therapy, and rest) for at least 6 weeks <p>OR</p> <ul style="list-style-type: none"> • Symptomatic osteoporotic vertebral fractures that happened less than 6 weeks ago and have led to hospitalization or persist at a level that prevents ambulation <p>OR</p> <ul style="list-style-type: none"> • Severe pain due to osteolytic lesions of the spine related to multiple myeloma or metastatic malignancies <p>Note: There is considerable variability in pain scores based on the literature review. If the patient is in intractable pain that cannot be managed safely with conservative treatment for at least 1 week, then percutaneous vertebroplasty surgery may be considered sooner than 6 weeks.</p> <p>Percutaneous vertebroplasty is considered investigational for all other indications, including use in acute vertebral fractures due to osteoporosis or trauma that have not led to hospitalization or prevent ambulation.</p>

Service	Investigational
<p>Percutaneous sacroplasty</p>	<p>Percutaneous sacroplasty is considered investigational for all indications, including use in sacral insufficiency fractures due to osteoporosis and spinal lesions due to metastatic malignancies or multiple myeloma.</p>

Documentation Requirements
<p>Clinical documentation of one of the following conditions:</p> <ul style="list-style-type: none"> • Symptomatic osteoporotic vertebral fractures that have failed to respond to conservative treatment (eg, analgesics, physical therapy, and rest) for at least 6 weeks <p>OR</p> <ul style="list-style-type: none"> • Symptomatic osteoporotic vertebral fractures that happened less than 6 weeks ago and have



Documentation Requirements

led to hospitalization or persist at a level that prevents walking

OR

- Severe pain due to osteolytic lesions of the spine related to multiple myeloma or metastatic malignancies

Coding

Code	Description
CPT	
0200T	Percutaneous sacral augmentation (sacroplasty), unilateral injection(s), including the use of a balloon or mechanical device, when used, 1 or more needles
0201T	Percutaneous sacral augmentation (sacroplasty), bilateral injections, including the use of a balloon or mechanical device, when used, 2 or more needles
22510	Percutaneous vertebroplasty (bone biopsy included when performed), 1 vertebral body, unilateral or bilateral injection, inclusive of all imaging guidance; cervicothoracic
22511	Percutaneous vertebroplasty (bone biopsy included when performed), 1 vertebral body, unilateral or bilateral injection, inclusive of all imaging guidance; lumbosacral
22512	Percutaneous vertebroplasty (bone biopsy included when performed), 1 vertebral body, unilateral or bilateral injection, inclusive of all imaging guidance; each additional cervicothoracic or lumbosacral vertebral body (List separately in addition to code for primary procedure)
22513	Percutaneous vertebral augmentation, including cavity creation (fracture reduction and bone biopsy included when performed) using mechanical device (eg, kyphoplasty), 1 vertebral body, unilateral or bilateral cannulation, inclusive of all imaging guidance; thoracic
22514	Percutaneous vertebral augmentation, including cavity creation (fracture reduction and bone biopsy included when performed) using mechanical device (eg, kyphoplasty), 1 vertebral body, unilateral or bilateral cannulation, inclusive of all imaging guidance; lumbar
22515	Percutaneous vertebral augmentation, including cavity creation (fracture reduction and bone biopsy included when performed) using mechanical device (eg, kyphoplasty), 1 vertebral body, unilateral or bilateral cannulation, inclusive of all imaging guidance; each additional thoracic or lumbar vertebral body (List separately in addition to code for primary procedure)

Note: CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). HCPCS codes, descriptions and materials are copyrighted by Centers for Medicare Services (CMS).



Related Information

Benefit Application

Percutaneous vertebroplasty/sacroplasty may be performed by interventional radiologists or orthopedic surgeons.

Evidence Review

Description

Percutaneous vertebroplasty is an interventional technique involving the fluoroscopically guided injection of polymethylmethacrylate (PMMA) into a weakened vertebral body. The technique has been investigated to provide mechanical support and symptomatic relief in patients with osteoporotic vertebral compression fractures or those with osteolytic lesions of the spine (eg, multiple myeloma, metastatic malignancies), as a treatment for sacral insufficiency fractures, and as a technique to limit blood loss related to surgery.

Background

Osteoporotic Fracture

Vertebral Compression Fracture

Osteoporotic compression fractures are common. It is estimated that up to one-half of women and approximately one-quarter of men will have a vertebral fracture at some point in their lives. However, only about one-third of vertebral fractures actually reach clinical diagnosis, and most symptomatic fractures will heal within a month. Nonetheless, some individuals with acute fractures will have severe pain and decreased function that interferes with their ability to ambulate and is not responsive to the usual medical management. Also, a minority of patients will exhibit chronic pain following osteoporotic compression fracture that presents challenges for medical management.



Treatment

Chronic symptoms do not tend to respond to the management strategies for acute pain, such as bed rest, immobilization or bracing device, and analgesic medication, sometimes including narcotic analgesics. The source of chronic pain after a vertebral compression fracture may not be from the vertebra itself but may be predominantly related to strain on muscles and ligaments secondary to kyphosis. This type of pain frequently does not improve with analgesics and may be better addressed through exercise. Improvements in pain and ability to function are the principal outcomes of interest for treatment of osteoporotic fractures.

Sacral Insufficiency Fractures

Sacral insufficiency fractures (SIFs) the consequences of stress on the weakened bone and often cause low back pain in the elderly population.¹ Osteoporosis is the most common risk factor for SIF. Spontaneous fracture of the sacrum in patients with osteoporosis was described by Lourie (1982) and presents as lower back and buttock pain with or without referred pain into the legs.² Although common, SIFs can escape detection due to low provider suspicion and poor sensitivity on plain radiographs, slowing the application of appropriate intervention.

Treatment

Similar interventions are used for sacral and vertebral fractures and include bedrest, bracing, and analgesics. Initial clinical improvements may occur quickly; however, resolution of all symptoms may not occur for 9 to 12 months.^{1,3}

Vertebral and Sacral Body Metastasis

Metastatic malignant disease of the spine generally involves the vertebrae/sacrum, with pain being the most frequent complaint.

Treatment

While radiotherapy and chemotherapy are frequently effective in reducing tumor burden and associated symptoms, pain relief may be delayed days to weeks, depending on tumor response.



Further, these approaches rely on bone remodeling to regain strength in the vertebrae/sacrum, which may necessitate supportive bracing to minimize the risk of vertebral/sacral collapse during healing. Improvements in pain and function are the primary outcomes of interest for treatment of bone malignancy with percutaneous vertebroplasty or sacroplasty.

Surgical Treatment Options

Percutaneous Vertebroplasty

Vertebroplasty is a surgical procedure that involves the injection of synthetic cement (eg, polymethylmethacrylate [PMMA], bis-glycidal dimethacrylate [Cortoss]⁴) into a fractured vertebra. It has been suggested that vertebroplasty may provide an analgesic effect through mechanical stabilization of a fractured or otherwise weakened vertebral body. However, other mechanisms of effect have been postulated, including thermal damage to intraosseous nerve fibers.

Percutaneous Sacroplasty

Sacroplasty evolved from the treatment of insufficiency fractures in the thoracic and lumbar vertebrae with vertebroplasty. The procedure, essentially identical to vertebroplasty, entails guided injection of polymethylmethacrylate (PMMA) through a needle inserted into the fracture zone. While first described in 2000 as a treatment for symptomatic sacral metastatic lesions,^{5,6} it is most often described as a minimally invasive alternative to conservative management⁷⁻⁹ for SIFs.

Pain and function are subjective outcomes and, thus, may be susceptible to placebo effects. Furthermore, the natural history of pain and disability associated with these conditions may vary. Therefore, controlled comparison studies would be valuable to demonstrate the clinical effectiveness of vertebroplasty and sacroplasty over and above any associated nonspecific or placebo effects and to demonstrate the effect of treatment compared with alternatives such as continued medical management.

In all clinical situations, adverse effects related to complications from vertebroplasty and sacroplasty are the primary harms to be considered. Principal safety concerns relate to the incidence and consequences of leakage of the injected PMMA or another injectate.⁴



Summary of Evidence

For individuals who have symptomatic osteoporotic vertebral fractures that are between 6 weeks and 1 year old who receive vertebroplasty, the evidence includes 2 randomized sham-controlled trials, nonblinded randomized controlled trials (RCTs) comparing vertebroplasty with conservative management, and systematic reviews of these RCTs. Relevant outcomes are symptoms, functional outcomes, quality of life, hospitalizations, medication use, and treatment-related morbidity. Despite the completion of numerous RCTs, including 2 with sham controls, the efficacy of vertebroplasty for painful osteoporotic compression fractures remains uncertain. Two meta-analysis studies, which included the 2 sham-controlled trials, have demonstrated mixed results. The two studies had methodologic issues, including the choice of sham procedure and the potential effect of the sham procedure to have a therapeutic effect by reducing pain. Questions have also been raised about the low percentage of patients screened who participated in the trial, the volume of polymethylmethacrylate injected, and the inclusion of patients with chronic pain. Overall, conclusions about the effect of vertebroplasty remain unclear. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals with symptomatic osteoporotic vertebral fractures that are less than 6 weeks old who receive vertebroplasty, the evidence includes a randomized sham-controlled trial and other nonblinded RCTs comparing vertebroplasty with conservative management. Relevant outcomes are symptoms, functional outcomes, quality of life, hospitalizations, medication use, and treatment-related morbidity. For acute fractures, conservative therapy consisting of rest, analgesics, and physical therapy is an option, and symptoms will resolve in a large percentage of patients with conservative treatment only. However, a sham-controlled randomized trial in patients who had severe pain of fewer than 6 weeks duration found a significant benefit using vertebroplasty for the treatment of osteoporotic vertebral fractures at the thoracolumbar junction. Other RCTs without sham controls have reported that vertebroplasty is associated with significant improvements in pain and reductions in the duration of bed rest. Given the high morbidity associated with extended bed rest in older adults, this procedure is considered to have a significant health benefit. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals with sacral insufficiency fractures who receive sacroplasty, the evidence includes two prospective cohort studies and a case series. Relevant outcomes are symptoms, functional outcomes, quality of life, hospitalizations, medication use, and treatment-related morbidity. No RCTs have been reported. The available evidence includes a prospective cohort study and a retrospective series with 243 patients. These studies have reported rapid and sustained decreases in pain following percutaneous sacroplasty. Additional literature has mostly reported



immediate improvements following the procedure. However, due to the small size of the evidence base, the harms associated with sacroplasty have not been adequately studied. The evidence is insufficient to determine the effects of the technology on health outcomes.

Ongoing and Unpublished Clinical Trials

Ongoing trials that might influence this policy are listed in [Table 1](#).

Table 1. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
Ongoing			
NCT02370628	Vertebroplasty in the treatment of acute fracture trial (The VITTA Trial)	495	Apr 2018
NCT02902250	The Comparative Study About the Effect of Vertebral Body Decompression Procedure and Conservative Treatment for Benign Vertebral Compression Fracture - Prospective Randomized Control Study	80	Apr 2018
NCT02489825	Pilot Study: Does Preventive Adjacent Level Cement Augmentation Positively Affect Reoperation Rates After Osteoporotic Vertebral Compression Fractures?	100	Dec 2018

NCT: national clinical trial

Clinical Input Received from Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.



2014 Input

In response to requests, input was received from 2 physician specialty societies and 3 academic medical centers while this policy was under review in 2014. Input was sought on the treatment of acute vertebral fractures when there is severe pain that has led to hospitalization or persists at a level that prevents ambulation, and on the treatment of traumatic fractures that have remained symptomatic after 6 weeks of conservative treatment. Input on these issues was mixed.

2008 Input

In response to requests, input was received from 5 physician specialty societies and 2 academic medical centers while this policy was under review in 2008. Unsolicited input was received from a sixth physician specialty society. All reviewers disagreed with the proposed policy and provided references in support of the use of vertebroplasty.

Practice Guidelines and Position Statements

American College of Radiology et al

The American College of Radiology and 4 other medical specialty associations updated a 2012 joint position statement on percutaneous vertebral augmentation in 2014.⁴⁰ The statement indicated that percutaneous vertebral augmentation with the use of vertebroplasty or kyphoplasty is a safe, efficacious, and durable procedure in appropriate patients with symptomatic osteoporotic and neoplastic fractures, when performed in accordance with public standards. The document also stated that these procedures are offered only when nonoperative medical therapy has not provided adequate pain relief or pain is significantly altering the patients' quality of life.

Society for Interventional Radiology

In a 2014 quality improvement guideline from Society for Interventional Radiology, failure of medical therapy is defined as follows⁴¹:



1. For a patient rendered nonambulatory as a result of pain from a weakened or fractured vertebral body, pain persisting at a level that prevents ambulation despite 24 hours of analgesic therapy;
2. For a patient with sufficient pain from a weakened or fractured vertebral body that physical therapy is intolerable, pain persisting at that level despite 24 hours of analgesic therapy; or
3. For any patient with a weakened or fractured vertebral body, unacceptable side effects such as excessive sedation, confusion, or constipation as a result of the analgesic therapy necessary to reduce pain to a tolerable level.

American Academy of Orthopaedic Surgeons

The American Academy of Orthopaedic Surgeons (AAOS) approved practice guidelines (2010) on the treatment of osteoporotic spinal compression fractures.⁴² AAOS approved a strong recommendation against the use of vertebroplasty for patients who “present with an osteoporotic spinal compression fracture on imaging with correlating clinical signs and symptoms and who are neurologically intact.” With this recommendation, AAOS expressed its confidence that future evidence is unlikely to overturn the existing evidence. As a note, these recommendations were based on a literature review through September 2009; therefore, the 2010 Klazen trial was not included in the systematic review.

National Institute for Health and Care Excellence

The National Institute for Health and Care Excellence (NICE) concluded in its 2003 guidance on percutaneous vertebroplasty that the current evidence on the safety and efficacy of vertebroplasty for vertebral compression fractures appeared “adequate to support the use of this procedure” to “provide pain relief for people with severe painful osteoporosis with loss of height and/or compression fractures of the vertebral body....”⁴³ The guidance also recommended that the procedure be limited to patients whose pain is refractory to more conservative treatment. A 2013 NICE guidance indicated that percutaneous vertebroplasty and percutaneous balloon kyphoplasty “are recommended as options for treating osteoporotic vertebral compression fractures” in persons having “severe, ongoing pain after a recent, unhealed vertebral fracture despite optimal pain management” and whose “pain has been confirmed to be at the level of the fracture by physical examination and imaging.”⁴⁴



In 2008, NICE issued guidance on the diagnosis and management of adults with metastatic spinal cord compression.⁴⁵ This guidance indicated that vertebroplasty or kyphoplasty should be considered for “patients who have vertebral metastases and no evidence of MSCC [metastatic spinal cord compression] or spinal instability if they have: mechanical pain resistant to conventional pain management, or vertebral body collapse.”

Medicare National Coverage

There is no national coverage determination (NCD). In the absence of an NCD, coverage decisions are left to the discretion of local Medicare carriers.

Regulatory Status

Vertebroplasty is a surgical procedure and, as such, is not subject to U.S. Food and Drug Administration (FDA) approval.

Polymethylmethacrylate (PMMA) bone cement was available as a drug product before enactment of FDA’s device regulation and was at first considered what FDA terms a “transitional device.” It was transitioned to a class III device requiring premarketing applications. Several orthopedic companies have received approval of their bone cement products since 1976. In 1999, PMMA was reclassified from class III to class II, which requires future 510(k) submissions to meet “special controls” instead of “general controls” to assure safety and effectiveness. Thus, use of PMMA in vertebroplasty represented an off-label use of an FDA-regulated product before 2005. In 2005, PMMA bone cements such as Spine-Fix® Biomimetic Bone Cement and Osteopal® V were cleared for marketing by FDA through the 510(k) process for the fixation of pathologic fractures of the vertebral body using vertebroplasty procedures.

The use of PMMA in sacroplasty is an off-label use of an FDA-regulated product (bone cements such as Spine-Fix® Biomimetic Bone Cement [Teknimed] and Osteopal® V [Heraeus]), because the 510(k) approval was for the fixation of pathologic fractures of the vertebral body using vertebroplasty procedures. Sacroplasty was not included. FDA product code: NDN.

In May 2009, Cortoss® (Stryker) Bone Augmentation Material was cleared for marketing by FDA through the 510(k) process. Cortoss® is a nonresorbable synthetic material that is a composite resin-based, bis-glycidal dimethacrylate. FDA classifies this product as a PMMA bone cement.



In 2010, the Parallax® Contour® Vertebral Augmentation Device (ArthroCare) was cleared for marketing by FDA through the 510(k) process. The device creates a void in cancellous bone that can then be filled with bone cement. FDA product code: HXG.

References

1. Gotis-Graham I, McGuigan L, Diamond T, et al. Sacral insufficiency fractures in the elderly. *J Bone Joint Surg Br.* Nov 1994;76(6):882-886. PMID 7983111
2. Lourie H. Spontaneous osteoporotic fracture of the sacrum. An unrecognized syndrome of the elderly. *JAMA.* Aug 13 1982;248(6):715-717. PMID 7097924
3. Lin J, Lachmann E, Nagler W. Sacral insufficiency fractures: a report of two cases and a review of the literature. *J Womens Health Gend Based Med.* Sep 2001;10(7):699-705. PMID 11571100
4. Bae H, Hatten HP, Jr., Linovitz R, et al. A prospective randomized FDA-IDE trial comparing Cortoss with PMMA for vertebroplasty: a comparative effectiveness research study with 24-month follow-up. *Spine (Phila Pa 1976).* Apr 1 2012;37(7):544-550. PMID 21738093
5. Dehdashti AR, Martin JB, Jean B, et al. PMMA cementoplasty in symptomatic metastatic lesions of the S1 vertebral body. *Cardiovasc Intervent Radiol.* May-Jun 2000;23(3):235-237. PMID 10821903
6. Marcy PY, Palussiere J, Descamps B, et al. Percutaneous cementoplasty for pelvic bone metastasis. *Support Care Cancer.* Nov 2000;8(6):500-503. PMID 11094996
7. Aretxabala I, Fraiz E, Perez-Ruiz F, et al. Sacral insufficiency fractures. High association with pubic rami fractures. *Clin Rheumatol.* Oct 2000;19(5):399-401. PMID 11055834
8. Leroux JL, Denat B, Thomas E, et al. Sacral insufficiency fractures presenting as acute low-back pain. Biomechanical aspects. *Spine (Phila Pa 1976).* Dec 1993;18(16):2502-2506. PMID 8303454
9. Newhouse KE, el-Khoury GY, Buckwalter JA. Occult sacral fractures in osteopenic patients. *J Bone Joint Surg Am.* Dec 1992;74(10):1472-1477. PMID 1364816
10. Blue Cross and Blue Shield Technology Evaluation Center (TEC). Percutaneous Vertebroplasty. *TEC Assessments.* 2000;Volume 15:Tab 21.
11. Blue Cross and Blue Shield Technology Evaluation Center (TEC). Percutaneous vertebroplasty for vertebral fractures caused by osteoporosis, malignancy, or hemangioma. *TEC Assessments.* 2004;Volume 19:Tab 13.
12. Blue Cross and Blue Shield Technology Evaluation Center (TEC). Percutaneous vertebroplasty for vertebral fractures caused by osteoporosis or malignancy. *TEC Assessments.* 2005;Volume 20:Tab 6.
13. Blue Cross and Blue Shield Technology Evaluation Center (TEC). Percutaneous vertebroplasty or kyphoplasty for vertebral fractures caused by osteoporosis or malignancy. *TEC Assessments.* 2008;Volume 23:Tab 5.
14. Blue Cross and Blue Shield Technology Evaluation Center (TEC). Percutaneous vertebroplasty or kyphoplasty for vertebral fractures caused by osteoporosis. *TEC Assessments.* 2009;Volume 24:Tab 7.
15. Blue Cross and Blue Shield Technology Evaluation Center (TEC). Percutaneous vertebroplasty for vertebral fractures caused by osteoporosis. *TEC Assessments.* 2010;Volume 25:Tab 9.
16. Buchbinder R, Golmohammadi K, Johnston RV, et al. Percutaneous vertebroplasty for osteoporotic vertebral compression fracture. *Cochrane Database Syst Rev.* Apr 30 2015(4):CD006349. PMID 25923524



17. Buchbinder R, Osborne RH, Ebeling PR, et al. A randomized trial of vertebroplasty for painful osteoporotic vertebral fractures. *N Engl J Med.* Aug 6 2009;361(6):557-568. PMID 19657121
18. Kallmes DF, Comstock BA, Heagerty PJ, et al. A randomized trial of vertebroplasty for osteoporotic spinal fractures. *N Engl J Med.* Aug 6 2009;361(6):569-579. PMID 19657122
19. Staples MP, Kallmes DF, Comstock BA, et al. Effectiveness of vertebroplasty using individual patient data from two randomised placebo controlled trials: meta-analysis. *BMJ.* Jul 12 2011;343:d3952. PMID 21750078
20. Xie L, Zhao ZG, Zhang SJ, et al. Percutaneous vertebroplasty versus conservative treatment for osteoporotic vertebral compression fractures: An updated meta-analysis of prospective randomized controlled trials. *Int J Surg.* Nov 2017;47:25-32. PMID 28939236
21. Kroon F, Staples M, Ebeling PR, et al. Two-year results of a randomized placebo-controlled trial of vertebroplasty for acute osteoporotic vertebral fractures. *J Bone Miner Res.* Jun 2014;29(6):1346-1355. PMID 24967454
22. Ostelo RW, Deyo RA, Stratford P, et al. Interpreting change scores for pain and functional status in low back pain: towards international consensus regarding minimal important change. *Spine (Phila Pa 1976).* Jan 1 2008;33(1):90-94. PMID 18165753
23. Comstock BA, Sitlani CM, Jarvik JG, et al. Investigational vertebroplasty safety and efficacy trial (INVEST): patient-reported outcomes through 1 year. *Radiology.* Oct 2013;269(1):224-231. PMID 23696683
24. Chen D, An ZQ, Song S, et al. Percutaneous vertebroplasty compared with conservative treatment in patients with chronic painful osteoporotic spinal fractures. *J Clin Neurosci.* Mar 2014;21(3):473-477. PMID 24315046
25. Farrokhi MR, Alibai E, Maghami Z. Randomized controlled trial of percutaneous vertebroplasty versus optimal medical management for the relief of pain and disability in acute osteoporotic vertebral compression fractures. *J Neurosurg Spine.* May 2011;14(5):561-569. PMID 21375382
26. Edidin AA, Ong KL, Lau E, et al. Mortality risk for operated and nonoperated vertebral fracture patients in the Medicare population. *J Bone Miner Res.* Jul 2011;26(7):1617-1626. PMID 21308780
27. Edidin AA, Ong KL, Lau E, et al. Morbidity and mortality after vertebral fractures: comparison of vertebral augmentation and nonoperative management in the Medicare population. *Spine (Phila Pa 1976).* Aug 01 2015;40(15):1228-1241. PMID 26020845
28. Lin JH, Chien LN, Tsai WL, et al. Early vertebroplasty associated with a lower risk of mortality and respiratory failure in aged patients with painful vertebral compression fractures: a population-based cohort study in Taiwan. *Spine J.* Sep 2017;17(9):1310-1318. PMID 28483705
29. Clark W, Bird P, Gonski P, et al. Safety and efficacy of vertebroplasty for acute painful osteoporotic fractures (VAPOUR): a multicentre, randomised, double-blind, placebo-controlled trial. *Lancet.* Oct 01 2016;388(10052):1408-1416. PMID 27544377
30. Klazen CA, Lohle PN, de Vries J, et al. Vertebroplasty versus conservative treatment in acute osteoporotic vertebral compression fractures (Vertos II): an open-label randomised trial. *Lancet.* Sep 25 2010;376(9746):1085-1092. PMID 20701962
31. Yi X, Lu H, Tian F, et al. Recompression in new levels after percutaneous vertebroplasty and kyphoplasty compared with conservative treatment. *Arch Orthop Trauma Surg.* Jan 2014;134(1):21-30. PMID 24287674
32. Leali PT, Solla F, Maestretti G, et al. Safety and efficacy of vertebroplasty in the treatment of osteoporotic vertebral compression fractures: a prospective multicenter international randomized controlled study. *Clin Cases Miner Bone Metab.* Sep-Dec 2016;13(3):234-236. PMID 28228788
33. Yang EZ, Xu JG, Huang GZ, et al. Percutaneous Vertebroplasty versus conservative treatment in aged patients with acute osteoporotic vertebral compression fractures: a prospective randomized controlled clinical study. *Spine (Phila Pa 1976).* Apr 2016;41(8):653-660. PMID 26630417
34. Frey ME, Depalma MJ, Cifu DX, et al. Percutaneous sacroplasty for osteoporotic sacral insufficiency fractures: a prospective, multicenter, observational pilot study. *Spine J.* Mar-Apr 2008;8(2):367-373. PMID 17981097
35. Kortman K, Ortiz O, Miller T, et al. Multicenter study to assess the efficacy and safety of sacroplasty in patients with osteoporotic sacral insufficiency fractures or pathologic sacral lesions. *J Neurointerv Surg.* Sep 1 2013;5(5):461-466. PMID 22684691



36. Frey ME, Warner C, Thomas SM, et al. Sacroplasty: a ten-year analysis of prospective patients treated with percutaneous sacroplasty: literature review and technical considerations. *Pain Physician*. Nov 2017;20(7):E1063-E1072. PMID 29149151
37. Dougherty RW, McDonald JS, Cho YW, et al. Percutaneous sacroplasty using CT guidance for pain palliation in sacral insufficiency fractures. *J Neurointerv Surg*. Jan 2014;6(1):57-60. PMID 23345629
38. Zaman FM, Frey M, Slipman CW. Sacral stress fractures. *Curr Sports Med Rep*. Feb 2006;5(1):37-43. PMID 16483515
39. Denis F, Davis S, Comfort T. Sacral fractures: an important problem. Retrospective analysis of 236 cases. *Clin Orthop Relat Res*. Feb 1988;227:67-81. PMID 3338224
40. Barr JD, Jensen ME, Hirsch JA, et al. Position statement on percutaneous vertebral augmentation: a consensus statement developed by the Society of Interventional Radiology (SIR), American Association of Neurological Surgeons (AANS) and the Congress of Neurological Surgeons (CNS), American College of Radiology (ACR), American Society of Neuroradiology (ASNR), American Society of Spine Radiology (ASSR), Canadian Interventional Radiology Association (CIRA), and the Society of NeuroInterventional Surgery (SNIS). *J Vasc Interv Radiol*. Feb 2014;25(2):171-181. PMID 24325929
41. Baerlocher MO, Saad WE, Dariushnia S, et al. Quality improvement guidelines for percutaneous vertebroplasty. *J Vasc Interv Radiol*. Feb 2014;25(2):165-170. PMID 24238815
42. McGuire RAJ. AAOS Now: Treating spinal compression fractures. 2010; <https://www.aaos.org/AAOSNow/2010/Oct/cover/cover1/?ssopc=1>. Accessed June 2018.
43. National Institute for Health and Care Excellence (NICE). Percutaneous vertebroplasty [IPG12]. 2003; <https://www.nice.org.uk/guidance/ipg12>. Accessed June 2018.
44. National Institute for Health and Care Excellence (NICE). Percutaneous vertebroplasty and percutaneous balloon kyphoplasty for treating osteoporotic vertebral compression fractures [TA279]. 2013; <https://www.nice.org.uk/guidance/ta279> Accessed June 2018.
45. National Institute for Health and Care Excellence (NICE). Metastatic spinal cord compression in adults: risk assessment, diagnosis and management [CG75]. 2008; <https://www.nice.org.uk/guidance/cg75/chapter/1-guidance>. Accessed June 2018.

History

Date	Comments
06/12/12	New policy, add to Radiology section. Policy replaces 6.01.520 in conjunction with 6.01.38.
09/25/12	Update Coding Section – ICD-10 codes are now effective 10/01/2014.
07/24/13	Replace policy. Rationale updated based on a literature review through March 2013. ACR 2012 practice guideline added. References 31, 33, 34, 37 added; others renumbered or removed. Policy statements unchanged.
06/19/14	Annual Review. Policy updated with literature review through March 25, 2014; references 22, 31, 40-42, and 45-46 added; policy statements unchanged. ICD-9 procedure code 81.65 removed; this is performed outpatient – ICD-10 procedure codes also removed, along with both sets of diagnosis codes.
01/12/15	Coding update. New CPT codes 22510-22515, effective 1/1/15, added to policy;



Date	Comments
	notation made regarding CPT codes 22520-22522 and 72291-72292, deleted as of 12/31/14.
08/11/15	Annual Review. Policy updated with literature review through March 3, 2015; references 18 and 27 added; Reworded the third policy statement for clarity: Percutaneous vertebroplasty is considered investigational for all other indications not listed above.
08/01/16	Annual Review, approved July 12, 2016. No change to policy statements. No new RCTs identified.
06/06/17	Coding update, removed HCPCS codes S2360 and S2361 as they were terminated 01/01/16.
08/01/17	Annual Review, approved July 18, 2017. Policy moved into the new format. Policy updated with literature review through March 23, 2017; references 9, 16, 26-27, and 30-31 added; vertebroplasty may be medically necessary in vertebral fractures of less than 6 weeks in duration that prevent ambulation.
07/01/18	Annual Review, approved June 22, 2018. Policy updated with literature review through February 2018; references 20, 28, and 36 added. Policy statements unchanged.

Disclaimer: This medical policy is a guide in evaluating the medical necessity of a particular service or treatment. The Company adopts policies after careful review of published peer-reviewed scientific literature, national guidelines and local standards of practice. Since medical technology is constantly changing, the Company reserves the right to review and update policies as appropriate. Member contracts differ in their benefits. Always consult the member benefit booklet or contact a member service representative to determine coverage for a specific medical service or supply. CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). ©2018 Premera All Rights Reserved.

Scope: Medical policies are systematically developed guidelines that serve as a resource for Company staff when determining coverage for specific medical procedures, drugs or devices. Coverage for medical services is subject to the limits and conditions of the member benefit plan. Members and their providers should consult the member benefit booklet or contact a customer service representative to determine whether there are any benefit limitations applicable to this service or supply. This medical policy does not apply to Medicare Advantage.



Discrimination is Against the Law

Premera Blue Cross complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex. Premera does not exclude people or treat them differently because of race, color, national origin, age, disability or sex.

Premera:

- Provides free aids and services to people with disabilities to communicate effectively with us, such as:
 - Qualified sign language interpreters
 - Written information in other formats (large print, audio, accessible electronic formats, other formats)
- Provides free language services to people whose primary language is not English, such as:
 - Qualified interpreters
 - Information written in other languages

If you need these services, contact the Civil Rights Coordinator.

If you believe that Premera has failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, or sex, you can file a grievance with:

Civil Rights Coordinator - Complaints and Appeals
PO Box 91102, Seattle, WA 98111
Toll free 855-332-4535, Fax 425-918-5592, TTY 800-842-5357
Email AppealsDepartmentInquiries@Premera.com

You can file a grievance in person or by mail, fax, or email. If you need help filing a grievance, the Civil Rights Coordinator is available to help you.

You can also file a civil rights complaint with the U.S. Department of Health and Human Services, Office for Civil Rights, electronically through the Office for Civil Rights Complaint Portal, available at <https://ocrportal.hhs.gov/ocr/portal/lobby.jsf>, or by mail or phone at: U.S. Department of Health and Human Services
200 Independence Avenue SW, Room 509F, HHH Building
Washington, D.C. 20201, 1-800-368-1019, 800-537-7697 (TDD)
Complaint forms are available at <http://www.hhs.gov/ocr/office/file/index.html>.

Getting Help in Other Languages

This Notice has Important Information. This notice may have important information about your application or coverage through Premera Blue Cross. There may be key dates in this notice. You may need to take action by certain deadlines to keep your health coverage or help with costs. You have the right to get this information and help in your language at no cost. Call 800-722-1471 (TTY: 800-842-5357).

አማርኛ (Amharic):

ይህ ማስታወቂያ አስፈላጊ መረጃ ይዟል። ይህ ማስታወቂያ ስለ ማመልከቻዎ ወይም የ Premera Blue Cross ሽፋን አስፈላጊ መረጃ ሊኖረው ይችላል። በዚህ ማስታወቂያ ውስጥ ቁልፍ ቀናት ሊኖሩ ይችላሉ። የጤና ሽፋንዎን ለማመልከት በአስፈላጊ እርዳታ ለማግኘት በተውሰኑ የጊዜ ገደቦች እርምጃ መውሰድ ይገባዎት ይሆናል። ይህን መረጃ እንዲያገኙ እና የለምንም ክፍያ በቋንቋዎ እርዳታ እንዲያገኙ መሰታወቅ አለዎት። በስልክ ቁጥር 800-722-1471 (TTY: 800-842-5357) ይደውሉ።

العربية (Arabic):

يحتوي هذا الإشعار على معلومات هامة. قد يحتوي هذا الإشعار على معلومات مهمة بخصوص طلبك أو التخطيط التي تزيد الحصول عليها من خلال Premera Blue Cross. قد تكون هناك تواريخ مهمة في هذا الإشعار. وقد تحتاج لاتخاذ إجراء في تاريخ معينة للحفاظ على تغطيتك الصحية أو للمساعدة في دفع التكاليف. يحق لك الحصول على هذه المعلومات والمساعدة بلغتك دون تكبد أية تكلفة. اتصل بـ 800-722-1471 (TTY: 800-842-5357)

中文 (Chinese):

本通知有重要的訊息。本通知可能有關於您透過 Premera Blue Cross 提交的申請或保險的重要訊息。本通知內可能有重要日期。您可能需要在截止日期之前採取行動，以保留您的健康保險或者費用補貼。您有權利免費以您的母語得到本訊息和幫助。請撥電話 800-722-1471 (TTY: 800-842-5357)。

Oromoo (Cushite):

Beeksisni kun odeeffannoo barbaachisaa qaba. Beeksisni kun sagantaa yookan karaa Premera Blue Cross tiin tajaajila keessan ilaalchisee odeeffannoo barbaachisaa qabaachuu danda'a. Guyyaawwan murteessaa ta'an beeksisa kana keessatti ilaalaa. Tarii kaffaltiidhaan deeggaramuuf yookan tajaajila fayyaa keessaniif guyyaa dhumaa irratti wanti raawwattan jiraachuu danda'a. Kaffaltii irraa bilisa haala ta'een afaan keessaniin odeeffannoo argachuu fi deeggarsa argachuuf mirga ni qabaattu. Lakkoofsa bilbilaa 800-722-1471 (TTY: 800-842-5357) tii bilbilaa.

Français (French):

Cet avis a d'importantes informations. Cet avis peut avoir d'importantes informations sur votre demande ou la couverture par l'intermédiaire de Premera Blue Cross. Le présent avis peut contenir des dates clés. Vous devez peut-être prendre des mesures par certains délais pour maintenir votre couverture de santé ou d'aide avec les coûts. Vous avez le droit d'obtenir cette information et de l'aide dans votre langue à aucun coût. Appelez le 800-722-1471 (TTY: 800-842-5357).

Kreyòl ayisyen (Creole):

Avi sila a gen Enfòmasyon Enpòtan ladann. Avi sila a kapab genyen enfòmasyon enpòtan konsènan aplikasyon w lan oswa konsènan kouvèti asirans lan atravè Premera Blue Cross. Kapab genyen dat ki enpòtan nan avi sila a. Ou ka gen pou pran kèk aksyon avan sèten dat limit pou ka kenbe kouvèti asirans sante w la oswa pou yo ka ede w avèk depans yo. Se dwa w pou resewva enfòmasyon sa a ak asistans nan lang ou pale a, san ou pa gen pou peye pou sa. Rele nan 800-722-1471 (TTY: 800-842-5357).

Deutsche (German):

Diese Benachrichtigung enthält wichtige Informationen. Diese Benachrichtigung enthält unter Umständen wichtige Informationen bezüglich Ihres Antrags auf Krankenversicherungsschutz durch Premera Blue Cross. Suchen Sie nach eventuellen wichtigen Terminen in dieser Benachrichtigung. Sie könnten bis zu bestimmten Stichtagen handeln müssen, um Ihren Krankenversicherungsschutz oder Hilfe mit den Kosten zu behalten. Sie haben das Recht, kostenlose Hilfe und Informationen in Ihrer Sprache zu erhalten. Rufen Sie an unter 800-722-1471 (TTY: 800-842-5357).

Hmoob (Hmong):

Tsab ntawv tshaj xo no muaj cov ntshiab lus tseem ceeb. Tej zaum tsab ntawv tshaj xo no muaj cov ntshiab lus tseem ceeb txog koj daim ntawv thov kev pab los yog koj qhov kev pab cuam hnu ntawm Premera Blue Cross. Tej zaum muaj cov hnuv tseem ceeb uas sau rau hauv daim ntawv no. Tej zaum koj kuj yuav tau ua qee yam uas pab kom koj ua tsis pub dhau cov caij nyuog uas teev tseg rau hauv daim ntawv no mas koj thiaj yuav tau txais kev pab cuam kho mob los yog kev pab them tej nqi kho mob ntawd. Koj muaj cai kom lawv muab cov ntshiab lus no uas tau muab sau ua koj hom lus pub dawb rau koj. Hu rau 800-722-1471 (TTY: 800-842-5357).

Iloko (Ilocano):

Daytoy a Pakdaar ket naglaon iti Napateg nga Impormasion. Daytoy a pakdaar mabalin nga adda ket naglaon iti napateg nga impormasion maipanggep iti aplikasyonyo wenna coverage babaen iti Premera Blue Cross. Daytoy ket mabalin dagiti importante a petsa iti daytoy a pakdaar. Mabalin nga adda rumbeng nga aramidenyo nga addang sakbay dagiti partikular a naituding nga aldaw tapno mapagtalinaedyo ti coverage ti salun-atyto wenna tulong kadagiti gastos. Adda karbenganyo a mangala iti daytoy nga impormasion ken tulong iti bukodyo a pagsasao nga awan ti bayadanyo. Tumawag iti numero nga 800-722-1471 (TTY: 800-842-5357).

Italiano (Italian):

Questo avviso contiene informazioni importanti. Questo avviso può contenere informazioni importanti sulla tua domanda o copertura attraverso Premera Blue Cross. Potrebbero esserci date chiave in questo avviso. Potrebbe essere necessario un tuo intervento entro una scadenza determinata per consentirti di mantenere la tua copertura o sovvenzione. Hai il diritto di ottenere queste informazioni e assistenza nella tua lingua gratuitamente. Chiama 800-722-1471 (TTY: 800-842-5357).

日本語 (Japanese):

この通知には重要な情報が含まれています。この通知には、Premera Blue Cross の申請または補償範囲に関する重要な情報が含まれている場合があります。この通知に記載されている可能性がある重要な日付をご確認ください。健康保険や有料サポートを維持するには、特定の期日までに行動を取らなければならない場合があります。ご希望の言語による情報とサポートが無料で提供されます。800-722-1471 (TTY: 800-842-5357)までお電話ください。

한국어 (Korean):

본 통지서에는 중요한 정보가 들어 있습니다. 즉 이 통지서는 귀하의 신청에 관하여 그리고 Premera Blue Cross 를 통한 커버리지에 관한 정보를 포함하고 있을 수 있습니다. 본 통지서에는 핵심이 되는 날짜들이 있을 수 있습니다. 귀하의 귀하의 건강 커버리지를 계속 유지하거나 비용을 절감하기 위해서 일정한 마감일까지 조치를 취해야 할 필요가 있을 수 있습니다. 귀하의 이러한 정보와 도움을 귀하의 언어로 비용 부담없이 얻을 수 있는 권리가 있습니다. 800-722-1471 (TTY: 800-842-5357) 로 전화하십시오.

ລາວ (Lao):

ແຈງການນີ້ມີຂໍ້ມູນສໍາຄັນ. ແຈງການນີ້ອາດຈະມີຂໍ້ມູນສໍາຄັນກ່ຽວກັບຄໍາຮ້ອງສະໝັກ ຫຼື ຄວາມຄົມຄອງປະກັນໄພຂອງທ່ານຜ່ານ Premera Blue Cross. ອາດຈະມີວັນທີ່ສໍາຄັນໃນແຈງການນີ້. ທ່ານອາດຈະຈໍາເປັນຕ້ອງດໍາເນີນການຕາມກຳນົດ ເວລາສະເພາະເພື່ອຮັກສາຄວາມຄົມຄອງປະກັນສະພາບ ຫຼື ຄວາມຊ່ວຍເຫຼືອເວັ້ນເວີ້ ຄ່າໃຊ້ຈ່າຍຂອງທ່ານໄດ້. ທ່ານມີສິດໄດ້ຮັບຂໍ້ມູນນີ້ ແລະ ຄວາມຊ່ວຍເຫຼືອເປັນພາສາຂອງທ່ານໂດຍບໍ່ເສຍຄ່າ. ໃຫ້ໃບທາ 800-722-1471 (TTY: 800-842-5357).

ភាសាខ្មែរ (Khmer):

សេចក្តីជូនដំណឹងនេះមានព័ត៌មានយ៉ាងសំខាន់។ សេចក្តីជូនដំណឹងនេះប្រហែលជាមានព័ត៌មានយ៉ាងសំខាន់អំពីទម្រង់បែបបទ ឬការរៀបចំរបស់អ្នកកាមរយ: Premera Blue Cross ។ ប្រហែលជាមាន កាលបរិច្ឆេទសំខាន់នៅក្នុងសេចក្តីជូនដំណឹងនេះ។ អ្នកប្រហែលជាត្រូវការបញ្ជាក់សមត្ថភាព ដល់កិច្ចការច្នៃផ្ទះធានា ដើម្បីនឹងរក្សាទុកការធានារ៉ាប់រងអនាគតរបស់អ្នក ឬប្រាក់ដុល្លារចេញថ្លៃ។ អ្នកមានសិទ្ធិទទួលព័ត៌មាននេះ និងដុល្លារនៅក្នុងភាសារបស់អ្នកដោយមិនអស់លុយឡើយ។ សូមទូរស័ព្ទ 800-722-1471 (TTY: 800-842-5357)។

ਪੰਜਾਬੀ (Punjabi):

ਇਸ ਨੋਟਿਸ ਵਿਚ ਖਾਸ ਜਾਣਕਾਰੀ ਹੈ. ਇਸ ਨੋਟਿਸ ਵਿਚ Premera Blue Cross ਵਲੋਂ ਤੁਹਾਡੀ ਕਵਰੇਜ ਅਤੇ ਅਰਜੀ ਬਾਰੇ ਮਹੱਤਵਪੂਰਨ ਜਾਣਕਾਰੀ ਹੋ ਸਕਦੀ ਹੈ . ਇਸ ਨੋਟਿਸ ਨਵ ਖਾਸ ਤਾਰੀਖਾਂ ਹੋ ਸਕਦੀਆਂ ਹਨ. ਜੇਕਰ ਤੁਸੀਂ ਜਸਰਤ ਕਵਰੇਜ ਰਿੱਖਣੀ ਹੋਵੇ ਜਾਂ ਓਸ ਦੀ ਲਾਗਤ ਜਵਿੱਚ ਮਦਦ ਦੇ ਇਕੱਠ ਹੋ ਤਾਂ ਤੁਹਾਨੂੰ ਅੰਤਮ ਤਾਰੀਖ ਤੋਂ ਪਹਿਲਾਂ ਢੁੱਝ ਖਾਸ ਕਦਮ ਚੁੱਕਣ ਦੀ ਲੋੜ ਹੋ ਸਕਦੀ ਹੈ ,ਤੁਹਾਨੂੰ ਮੁਫਤ ਵਿੱਚ ਤੋਂ ਅਪਣੀ ਭਾਸ਼ਾ ਵਿੱਚ ਜਾਣਕਾਰੀ ਅਤੇ ਮਦਦ ਪ੍ਰਾਪਤ ਕਰਨ ਦਾ ਅਧਿਕਾਰ ਹੈ ,ਕਾਲ 800-722-1471 (TTY: 800-842-5357).

فارسی (Farsi):

این اعلامیه حاوی اطلاعات مهم میباشد. این اعلامیه ممکن است حاوی اطلاعات مهم درباره فرم تقاضا و یا پوشش بیمه ای شما از طریق Premera Blue Cross باشد. به تاریخ های مهم در این اعلامیه توجه نمایید. شما ممکن است برای حفظ پوشش بیمه تان یا کمک در پرداخت هزینه های درمانی تان، به تاریخ های مشخصی برای انجام کارهای خاصی احتیاج داشته باشید. شما حق این را دارید که این اطلاعات و کمک را به زبان خود به طور رایگان دریافت نمایید. برای کسب اطلاعات با شماره 800-722-1471 (کلیران TTY تماس باشماره 800-842-5357) تماس برقرار نمایید.

Polskie (Polish):

To ogłoszenie może zawierać ważne informacje. To ogłoszenie może zawierać ważne informacje odnośnie Państwa wniosku lub zakresu świadczeń poprzez Premera Blue Cross. Prosimy zwrócić uwagę na kluczowe daty, które mogą być zawarte w tym ogłoszeniu aby nie przekroczyć terminów w przypadku utrzymania polisy ubezpieczeniowej lub pomocy związanej z kosztami. Macie Państwo prawo do bezpłatnej informacji we własnym języku. Zadzwońcie pod 800-722-1471 (TTY: 800-842-5357).

Português (Portuguese):

Este aviso contém informações importantes. Este aviso poderá conter informações importantes a respeito de sua aplicação ou cobertura por meio do Premera Blue Cross. Poderão existir datas importantes neste aviso. Talvez seja necessário que você tome providências dentro de determinados prazos para manter sua cobertura de saúde ou ajuda de custos. Você tem o direito de obter esta informação e ajuda em seu idioma e sem custos. Ligue para 800-722-1471 (TTY: 800-842-5357).

Română (Romanian):

Prezenta notificare conține informații importante privind cererea sau acoperirea asigurării dumneavoastră de sănătate prin Premera Blue Cross. Pot exista date cheie în această notificare. Este posibil să fie nevoie să acționați până la anumite termene limită pentru a vă menține acoperirea asigurării de sănătate sau asistența provizorie la costuri. Aveți dreptul de a obține gratuit aceste informații și ajutor în limba dumneavoastră. Sunați la 800-722-1471 (TTY: 800-842-5357).

Русский (Russian):

Настоящее уведомление содержит важную информацию. Это уведомление может содержать важную информацию о вашем заявлении или страховом покрытии через Premera Blue Cross. В настоящем уведомлении могут быть указаны ключевые даты. Вам, возможно, потребуется принять меры к определенным предельным срокам для сохранения страхового покрытия или помощи с расходами. Вы имеете право на бесплатное получение этой информации и помощь на вашем языке. Звоните по телефону 800-722-1471 (TTY: 800-842-5357).

Fa'asamoa (Samoan):

Atonu ua iai i lenei fa'asilasilaga ni fa'amatalaga e sili ona taua e tatau ona e malamalama i ai. O lenei fa'asilasilaga o se fesoasoani e fa'amatala atili i ai i le tulaga o le polokalame, Premera Blue Cross, ua e tau fia maua atu i ai. Fa'amolemole, ia e iloilo fa'alelei i aso fa'apitoa olo'o iai i lenei fa'asilasilaga taua. Masalo o le'a iai ni feau e tatau ona e faia ao le'i aulia le aso ua ta'ua i lenei fa'asilasilaga ina ia e iai pea ma maua fesoasoani mai ai i le polokalame a le Malo olo'o e iai i ai. Olo'o iai iate oe le aia tatau e maua atu i lenei fa'asilasilaga ma lenei fa'matalaga i legagana e te malamalama i ai aunoa ma se togiga tupe. Vili atu i le telefoni 800-722-1471 (TTY: 800-842-5357).

Español (Spanish):

Este Aviso contiene información importante. Es posible que este aviso contenga información importante acerca de su solicitud o cobertura a través de Premera Blue Cross. Es posible que haya fechas clave en este aviso. Es posible que deba tomar alguna medida antes de determinadas fechas para mantener su cobertura médica o ayuda con los costos. Usted tiene derecho a recibir esta información y ayuda en su idioma sin costo alguno. Llame al 800-722-1471 (TTY: 800-842-5357).

Tagalog (Tagalog):

Ang Paunawa na ito ay naglalaman ng mahalagang impormasyon tungkol sa iyong aplikasyon o pagsakop sa pamamagitan ng Premera Blue Cross. Maaaring may mga mahalagang petsa dito sa paunawa. Maaring mangailangan ka na magsagawa ng hakbang sa ilang mga itinakdang panahon upang mapanatili ang iyong pagsakop sa kalusugan o tulong na walang gastos. May karapatan ka na makakuha ng ganiitong impormasyon at tulong sa iyong wika ng walang gastos. Tumawag sa 800-722-1471 (TTY: 800-842-5357).

ไทย (Thai):

ประกาศนี้มีข้อมูลสำคัญ ประกาศนี้อาจมีข้อมูลที่สำคัญเกี่ยวกับกาการสมัครหรือขอบเขตประกันสุขภาพของคุณผ่าน Premera Blue Cross และอาจมีกำหนดการในประกาศนี้ คุณอาจจะต้องดำเนินการภายในกำหนดระยะเวลาที่แน่นอนเพื่อจะรักษาการประกันสุขภาพของคุณหรือการช่วยเหลือที่มีค่าใช้จ่าย คุณมีสิทธิที่จะได้รับข้อมูลและความช่วยเหลือนี้ในภาษาของคุณโดยไม่มีค่าใช้จ่าย โทร 800-722-1471 (TTY: 800-842-5357)

Український (Ukrainian):

Це повідомлення містить важливу інформацію. Це повідомлення може містити важливу інформацію про Ваше звернення щодо страховального покриття через Premera Blue Cross. Зверніть увагу на ключові дати, які можуть бути вказані у цьому повідомленні. Існує імовірність того, що Вам треба буде здійснити певні кроки у конкретні кінцеві строки для того, щоб зберегти Ваше медичне страхування або отримати фінансову допомогу. У Вас є право на отримання цієї інформації та допомоги безкоштовно на Вашій рідній мові. Дзвоніть за номером телефону 800-722-1471 (TTY: 800-842-5357).

Tiếng Việt (Vietnamese):

Thông báo này cung cấp thông tin quan trọng. Thông báo này có thông tin quan trọng về đơn xin tham gia hoặc hợp đồng bảo hiểm của quý vị qua chương trình Premera Blue Cross. Xin xem ngày quan trọng trong thông báo này. Quý vị có thể phải thực hiện theo thông báo đúng trong thời hạn để duy trì bảo hiểm sức khỏe hoặc được trợ giúp thêm về chi phí. Quý vị có quyền được biết thông tin này và được trợ giúp bằng ngôn ngữ của mình miễn phí. Xin gọi số 800-722-1471 (TTY: 800-842-5357).