

## MEDICAL POLICY – 7.01.131

## Transcatheter Pulmonary Valve Implantation

BCBSA Ref. Policy: 7.01.131

Effective Date: Sept. 1, 2020

Last Revised: Aug. 4, 2020


Replaces: N/A

RELATED MEDICAL POLICIES:

None

Select a hyperlink below to be directed to that section.

[POLICY CRITERIA](#) | [DOCUMENTATION REQUIREMENTS](#) | [CODING](#)  
[RELATED INFORMATION](#) | [EVIDENCE REVIEW](#) | [REFERENCES](#) | [HISTORY](#)

 Clicking this icon returns you to the hyperlinks menu above.

## Introduction

Congenital heart disease is a term that means a person was born with heart problems. These difficulties affect the heart's function and structure. Congenital heart disease can range from mild, which may not need treatment, to severe, which often does. One congenital heart defect is known as right ventricular outflow tract (RVOT) dysfunction. Essentially, it's a problem with how the blood flows as it leaves the heart and goes to the lungs. Repairing it requires reconstructing certain areas of the heart and placing a tube (conduit) to allow the blood to flow correctly. Over a long period of time the conduit can become narrowed or a specific valve can become leaky. A second valve replacement surgery may be needed in this situation. This second surgery is usually done as an open surgery. However, surgery using a long, thin tube (a heart catheter) instead of open heart surgery can be done in certain situations. This policy describes when an additional RVOT surgery using a catheter may be considered medically necessary.

**Note:** The Introduction section is for your general knowledge and is not to be taken as policy coverage criteria. The rest of the policy uses specific words and concepts familiar to medical professionals. It is intended for providers. A provider can be a person, such as a doctor, nurse, psychologist, or dentist. A provider also can be a place where medical care is given, like a hospital, clinic, or lab. This policy informs them about when a service may be covered.

Service	Medical Necessity
<b>Transcatheter pulmonary valve implantation (TPVI)</b>	<p><b>Transcatheter pulmonary valve implantation is considered medically necessary for patients with congenital heart disease and current right ventricular outflow tract obstruction (RVOT) or regurgitation including the following indications:</b></p> <ul style="list-style-type: none"> <li>• Individuals with right ventricle-to-pulmonary artery conduit with or without bioprosthetic valve with at least moderate pulmonic regurgitation</li> </ul> <p><b>OR</b></p> <ul style="list-style-type: none"> <li>• Individuals with native or patched RVOT with at least moderate pulmonic regurgitation</li> </ul> <p><b>OR</b></p> <ul style="list-style-type: none"> <li>• Individuals with right ventricle-to-pulmonary artery conduit with or without bioprosthetic valve with pulmonic stenosis (mean RVOT gradient at least 35 mm Hg)</li> </ul> <p><b>OR</b></p> <ul style="list-style-type: none"> <li>• Individuals with native or patched RVOT with pulmonic stenosis (mean RVOT gradient at least 35 mm Hg).</li> </ul> <p><b>Transcatheter pulmonary valve implantation is considered investigational for all other indications.</b></p>

Documentation Requirements
<p><b>The patient’s medical records submitted for review should document that medical necessity criteria are met. The record should include clinical documentation of:</b></p> <ul style="list-style-type: none"> <li>• Diagnosis/condition</li> <li>• History and physical examination documenting the severity of the condition</li> <li>• Right ventricular outflow tract (RVOT) gradient</li> <li>• Pulmonic regurgitation (if present)</li> </ul>

## Coding



Code	Description
<b>CPT</b>	
33477	Transcatheter pulmonary valve implantation, percutaneous approach, including pre-stenting of the valve delivery site, when performed

**Note:** CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). HCPCS codes, descriptions and materials are copyrighted by Centers for Medicare Services (CMS).

## Related Information

N/A

## Evidence Review

### Description

Transcatheter pulmonary valve implantation (TPVI) is a less invasive alternative to open surgical pulmonary valve replacement or reconstruction for right ventricular outflow tract (RVOT) obstruction. Percutaneous pulmonary valve replacement may be indicated for congenital pulmonary stenosis. Pulmonary stenosis or regurgitation in a patient with congenital heart disease (CHD) who has previously undergone RVOT surgery are additional indications. Patients with prior CHD repair are at risk of needing repeated reconstruction procedures.

### Background

#### Congenital Heart Disease

Congenital heart disease, including tetralogy of Fallot, pulmonary atresia, and transposition of the great arteries, is generally treated by surgical repair at an early age. This involves reconstruction of the right ventricular outflow tract (RVOT) and pulmonary valve using a surgical homograft or a bovine-derived valved conduit. These repairs are prone to development of pulmonary stenosis or regurgitation over long periods of follow-up.



Because individuals with surgically corrected congenital heart disease are living into adulthood, RVOT dysfunction following initial repair has become more common. Calcification of the RVOT conduit can lead to pulmonary stenosis, while aneurysmal dilatation can result in pulmonary regurgitation. RVOT dysfunction can lead to decreased exercise tolerance, potentially fatal arrhythmias, and/or irreversible right ventricular dysfunction.<sup>1</sup>

## **Treatment**

Interventions for RVOT dysfunction often require numerous repeat open heart procedures for patients who live into adulthood. Treatment options for pulmonary stenosis are open surgery with valve replacement, balloon dilatation, or percutaneous stenting.<sup>1</sup> Interventions for pulmonary regurgitation are primarily surgical, either reconstruction of the RVOT conduit or replacement of the pulmonary valve. The optimal timing of these interventions is not well understood.<sup>2</sup>

Transcatheter pulmonary valve replacement offers a less invasive treatment option for patients with prior surgery for congenital heart disease and RVOT dysfunction. It is possible that a less invasive valve replacement technique could spare patients from multiple repeat open heart procedures over long periods of follow-up.

## **Summary of Evidence**

For individuals who have a history of congenital heart disease (CHD) and current right ventricular outflow tract (RVOT) obstruction who receive transcatheter pulmonary valve implantation (TPVI) with a U.S. Food and Drug Administration (FDA)-approved device and indication, the evidence includes prospective, interventional, noncomparative studies and multiple prospective and retrospective case or cohort series. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, hospitalizations, and treatment-related morbidity and mortality. Results of the case series have indicated that there is a high rate of procedural success and low procedural mortality, although the rates of serious procedural adverse events reported ranges from 3.0% to 7.4%. Most valves have demonstrated competent functioning by Doppler echocardiography at 6- to 12-month follow-ups, but complications (eg, stent fractures, need for re-interventions) were reported in an FDA analysis at rates of 18% and 7%, respectively. Other publications with longer follow-up have reported stent fractures in up to 26% of patients; however, most stent fractures did not required reintervention. Studies with follow-up extending to a maximum of 7 years postprocedure have suggested that the functional and hemodynamic improvements are durable, but a relatively high proportion of patients (20%-



30%) have required reintervention on the pulmonary valve. No comparative studies were identified, and there is no direct evidence that TPVI reduces future open heart procedures. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have a history of CHD and current RVOT obstruction who receive TPVI with a non-FDA-approved device or indication, the evidence includes case series. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, hospitalizations, and treatment-related morbidity and mortality. There is limited evidence on the off-label use of TPVI, including the use of a non-FDA-approved valve, or use of an approved valve for a non-FDA-approved indication. The published case series enrolled relatively few patients and are heterogeneous regarding devices used and indications for TPVI. The evidence is insufficient to determine the effects of the technology on health outcomes.

## Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in [Table 1](#).

**Table 1. Summary of Key Trials**

NCT No.	Trial Name	Planned Enrollment	Completion Date
<b>Ongoing</b>			
<a href="#">NCT00740870</a> <sup>a</sup>	Implantation of the Medtronic Melody Transcatheter Pulmonary Valve in Patients With Dysfunctional RVOT Conduits: A Feasibility Study	171	Jul 2020
<a href="#">NCT02744677</a> <sup>a</sup>	COngenital Multicenter Trial of Pulmonic vAlve Dysfunction Studying the SAPIEN 3 interventIOnal THV	108	Dec 2027
<a href="#">NCT02979587</a>	The Medtronic Harmony™ Transcatheter Pulmonary Valve Clinical Study	184	Dec 2024
<a href="#">NCT02987387</a> <sup>a</sup>	New Enrollment SAPIEN XT Post-Approval Study	191	Aug 2027

NCT: national clinical trial

<sup>a</sup> Denotes industry-sponsored or cosponsored trial



## Clinical Input Received from Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

### 2018 Input

In response to requests, clinical input on the use of transcatheter pulmonary valve implantation (TPVI) was received from two specialty society-level respondents while this policy was under review in 2018. The combined clinical input response incorporated input from a panel including physicians affiliated with academic medical centers.

Based on the evidence and independent clinical input, the clinical input supports that the following indications provide a clinically meaningful improvement in the net health outcome and are consistent with generally accepted medical practice:

- Use of TPVI for individuals with right ventricle-to-pulmonary artery conduit with or without bioprosthetic valve with at least moderate pulmonic regurgitation;
- Use of TPVI for individuals with native or patched right ventricular outflow tract (RVOT) with at least moderate pulmonic regurgitation;
- Use of TPVI for individuals with right ventricle-to-pulmonary artery conduit with or without bioprosthetic valve with pulmonic stenosis (mean RVOT gradient at least 35 mm Hg); or
- Use of TPVI for individuals with native or patched RVOT with pulmonic stenosis (mean RVOT gradient at least 35 mm Hg)

### 2011 Input

In response to requests, input was received from six academic medical centers while this policy was under review in 2011. Overall response to whether TPVI was investigational was mixed, with two of five reviewers agreeing with the investigational status, and three reviewers indicating partial support. Most reviewers (4/5) indicated that there is a subpopulation of patients who are at high risk for surgery or who are not candidates for surgery, and for whom there are no other



available options. These reviewers felt TPVI was a viable alternative that offered potential benefit for these patients.

## **Practice Guidelines and Position Statements**

### **Society for Cardiovascular Angiography and Interventions et al**

The Society for Cardiovascular Angiography and Interventions, American Association for Thoracic Surgery, American College of Cardiology (ACC) and the Society of Thoracic Surgeons (2015) published a consensus-based report on operator and institutional requirements for transcatheter pulmonary valve implantation (TPVI).<sup>44</sup> Recommendations to qualify for a TPVI program included 150 catheterizations per year, association with a surgical program, submission of all cases to a national registry, and, for patients, 80% freedom from re-intervention at 1 year.

### **American Heart Association and American College of Cardiology**

The 2014 American Heart Association (AHA) and the ACC guidelines on the management of patients with valvular disease and the 2017 AHA and ACC focused update<sup>45,46</sup> do not make specific recommendations on the treatment of primary pulmonary valve disease (stenosis or regurgitation), but instead referred to the 2008 guidelines for the management of adults with congenital heart disease.

In 2008, the AHA and ACC (in collaboration with other medical societies) issued guidelines for the management of adults with congenital heart disease.<sup>47</sup> For patients with isolated valvular pulmonary stenosis, the guidelines make recommendations on balloon valvulotomy or surgical intervention; however, TPVI was not addressed. In 2015, an AHA scientific statement on congenital heart disease in older adults was published and meant to complement the 2008 ACC and AHA guidelines for adults with congestive heart disease.<sup>48</sup> The intent was to outline the natural history, ramifications of childhood repair, and late initial diagnosis of congestive heart disease in older adults. The statement commented on the emerging use of the currently available transcatheter valve repair devices for both pulmonary stenosis and pulmonary valve regurgitation primarily after repair of tetralogy of Fallot. There was a specific comment that contemporary morbidity, mortality, and durability of surgical pulmonary valve regurgitation are unknown and, therefore, no contemporaneous benchmark against which to compare transcatheter valve implantation.



## Regulatory Status

Devices for transcatheter pulmonary valve implantation were initially cleared from marketing by the U.S. Food and Drug Administration (FDA) through the humanitarian device exemption (HDE) process or used off-label until approved by FDA through the premarket approval (PMA) process between 2015 and 2016 (see [Table 2](#)).

**Table 2. Regulatory Status of Transcatheter Pulmonary Valve Implantation Devices**

Device	Manufacturer	Date Approved	PMA No.	Indications
Melody® Transcatheter Pulmonary Valve (TPV)	Medtronic	Jan 2010	H080002 (HDE)	Pulmonary valve replacement for pediatric and adult patients with a dysfunctional, noncompliant RVOT conduit
Melody TPV	Medtronic	Jan 2015	P140017	Pulmonary valve replacement for pediatric and adult patients with a dysfunctional, noncompliant RVOT conduit
Melody TPV	Medtronic	Feb 2017	P140017/S005	Valve-in-valve for patients with a dysfunctional surgical bioprosthetic pulmonary valve
SAPIEN XT™ Transcatheter Heart Valve (pulmonic)	Edwards Lifesciences	Feb 2016	P130009/S037	Pulmonary valve replacement for pediatric and adult patients with a dysfunctional, noncompliant RVOT conduit

HDE: humanitarian device exemption; PMA: premarket approval; RVOT: right ventricular outflow tract.

The Melody® Transcatheter Pulmonary Valve (TPV) and the Ensemble® Transcatheter Valve Delivery System are used together for percutaneous replacement of a dysfunctional pulmonary valve. The Melody® valve consists of a section of bovine jugular vein with an intact native venous valve. The valve and surrounding tissue are sutured within a platinum-iridium stent scaffolding. The transcatheter delivery system consists of a balloon-in-balloon catheter with a retractable sheath and distal cup into which the valve is placed. The procedure is performed on a beating heart without the use of cardiopulmonary bypass.





The Melody® valve is first crimped to fit into the delivery system. It is introduced through the femoral vein and advanced into the right side of the heart and put into place at the site of the pulmonary valve. The inner balloon is inflated to open the artificial valve, and then the outer balloon is inflated to position the valve into place.

In January 2010, the Melody® TPV and the Ensemble® Transcatheter Valve Delivery System (Medtronic) were approved by FDA under the HDE program for use as an adjunct to surgery in the management of pediatric and adult patients with the following clinical conditions:

- Existence of a full (circumferential) RVOT conduit that is 16 mm or greater in diameter when originally implanted, and
- Dysfunctional RVOT conduits with clinical indication for intervention, and either:
  - regurgitation: moderate-to-severe regurgitation, or
  - stenosis: mean RVOT gradient  $\geq 35$  mm Hg

On January 27, 2015, approval of the Melody® system was amended to a PMA because FDA determined that the device represented a breakthrough technology.<sup>3</sup> The PMA was based, in part, on two prospective clinical studies, the Melody TPV Long-term Follow-up Post Approval Study and the Melody TPV New Enrollment Post Approval Study.

On February 24, 2017, approval of the Melody® system was expanded to include patients with a dysfunctional surgical bioprosthetic valve (valve-in-valve).

The Edwards SAPIEN XT™ Transcatheter Heart Valve (Pulmonic) (Edwards Lifesciences) is composed of a stainless steel frame with bovine pericardial tissue leaflets and available in 23- and 26-mm sizes. It includes a delivery accessories system. On February 29, 2016, it was approved by FDA as a supplement “for use in pediatric and adult patients with a dysfunctional, noncompliant Right Ventricular Outflow Tract (RVOT) conduit with a clinical indication for intervention and:

- pulmonary regurgitation  $\geq$  moderate and/or
- mean RVOT gradient  $\geq 35$  mmHg”

The approval for the pulmonic valve indication is a supplement to the 2014 PMA for use of the Edwards SAPIEN XT™ Transcatheter Heart Valve System for relief of aortic stenosis in patients with symptomatic heart disease due to severe native calcific aortic stenosis and who are judged by a heart team, including a cardiac surgeon, to be at high or greater risk for open surgical



therapy (ie, Society of Thoracic Surgeons operative risk score  $\geq 8\%$  or at a  $\geq 15\%$  risk of mortality at 30 days).

FDA product code: NPV

## References

1. Khambadkone S, Nordmeyer J, Bonhoeffer P. Percutaneous implantation of the pulmonary and aortic valves: indications and limitations. *J Cardiovasc Med (Hagerstown)*. Jan 2007; 8(1): 57-61. PMID 17255818
2. McElhinney DB, Hellenbrand WE, Zahn EM, et al. Short- and medium-term outcomes after transcatheter pulmonary valve placement in the expanded multicenter US melody valve trial. *Circulation*. Aug 03 2010; 122(5): 507-16. PMID 20644013
3. Food and Drug Administration. Summary of Safety and Effectiveness Data: Melody™ Transcatheter Pulmonary Valve. 2015; [https://www.accessdata.fda.gov/cdrh\\_docs/pdf14/p140017b.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf14/p140017b.pdf). Accessed July, 2020.
4. Zahn EM, Hellenbrand WE, Lock JE, et al. Implantation of the melody transcatheter pulmonary valve in patients with a dysfunctional right ventricular outflow tract conduit early results from the u.s. Clinical trial. *J Am Coll Cardiol*. Oct 27 2009; 54(18): 1722-9. PMID 19850214
5. Cheatham JP, Hellenbrand WE, Zahn EM, et al. Clinical and hemodynamic outcomes up to 7 years after transcatheter pulmonary valve replacement in the US melody valve investigational device exemption trial. *Circulation*. Jun 02 2015; 131(22): 1960-70. PMID 25944758
6. Batra AS, McElhinney DB, Wang W, et al. Cardiopulmonary exercise function among patients undergoing transcatheter pulmonary valve implantation in the US Melody valve investigational trial. *Am Heart J*. Feb 2012; 163(2): 280-7. PMID 22305848
7. Armstrong AK, Balzer DT, Cabalka AK, et al. One-year follow-up of the Melody transcatheter pulmonary valve multicenter post-approval study. *JACC Cardiovasc Interv*. Nov 2014; 7(11): 1254-62. PMID 25459038
8. Gillespie MJ, McElhinney DB, Kreutzer J, et al. Transcatheter Pulmonary Valve Replacement for Right Ventricular Outflow Tract Conduit Dysfunction After the Ross Procedure. *Ann Thorac Surg*. Sep 2015; 100(3): 996-1002; discussion 1002-3. PMID 26190388
9. Food and Drug Administration. Conditions for Approval for an HDE: Medtronic Melody Transcatheter Pulmonary Valve (Model PB 10) and Medtronic Ensemble Transcatheter Valve Delivery System (NU10) (H080002). 2010; [https://www.accessdata.fda.gov/cdrh\\_docs/pdf8/H080002A.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf8/H080002A.pdf). Accessed July, 2020.
10. Food and Drug Administration. Summary of Safety and Effectiveness Data: Melody™ Transcatheter Pulmonary Valve, models PB1016 and PB1018; Ensemble™ Transcatheter Valve Delivery System 2017; [https://www.accessdata.fda.gov/cdrh\\_docs/pdf14/p140017s005b.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf14/p140017s005b.pdf). Accessed July, 2020.
11. Lurz P, Coats L, Khambadkone S, et al. Percutaneous pulmonary valve implantation: impact of evolving technology and learning curve on clinical outcome. *Circulation*. Apr 15 2008; 117(15): 1964-72. PMID 18391109
12. Eicken A, Ewert P, Hager A, et al. Percutaneous pulmonary valve implantation: two-centre experience with more than 100 patients. *Eur Heart J*. May 2011; 32(10): 1260-5. PMID 21273201
13. Khambadkone S, Coats L, Taylor A, et al. Percutaneous pulmonary valve implantation in humans: results in 59 consecutive patients. *Circulation*. Aug 23 2005; 112(8): 1189-97. PMID 16103239
14. Momenah TS, El Oakley R, Al Najashi K, et al. Extended application of percutaneous pulmonary valve implantation. *J Am Coll Cardiol*. May 19 2009; 53(20): 1859-63. PMID 19442885



15. Nordmeyer J, Coats L, Bonhoeffer P. Current experience with percutaneous pulmonary valve implantation. *Semin Thorac Cardiovasc Surg.* 2006; 18(2): 122-5. PMID 17157232
16. Nordmeyer J, Coats L, Lurz P, et al. Percutaneous pulmonary valve-in-valve implantation: a successful treatment concept for early device failure. *Eur Heart J.* Mar 2008; 29(6): 810-5. PMID 18316357
17. Vezmar M, Chaturvedi R, Lee KJ, et al. Percutaneous pulmonary valve implantation in the young 2-year follow-up. *JACC Cardiovasc Interv.* Apr 2010; 3(4): 439-48. PMID 20398873
18. Muller J, Engelhardt A, Fratz S, et al. Improved exercise performance and quality of life after percutaneous pulmonary valve implantation. *Int J Cardiol.* May 15 2014; 173(3): 388-92. PMID 24713459
19. Butera G, Milanesi O, Spadoni I, et al. Melody transcatheter pulmonary valve implantation. Results from the registry of the Italian Society of Pediatric Cardiology. *Catheter Cardiovasc Interv.* Feb 2013; 81(2): 310-6. PMID 22718682
20. Fraisse A, Aldebert P, Malekzadeh-Milani S, et al. Melody (R) transcatheter pulmonary valve implantation: results from a French registry. *Arch Cardiovasc Dis.* Nov 2014; 107(11): 607-14. PMID 25453718
21. Borik S, Crean A, Horlick E, et al. Percutaneous pulmonary valve implantation: 5 years of follow-up: does age influence outcomes?. *Circ Cardiovasc Interv.* Feb 2015; 8(2): e001745. PMID 25652317
22. Markham R, Challa A, Kyranis S, et al. Outcomes Following Melody Transcatheter Pulmonary Valve Implantation for Right Ventricular Outflow Tract Dysfunction in Repaired Congenital Heart Disease: First Reported Australian Single Centre Experience. *Heart Lung Circ.* Oct 2017; 26(10): 1085-1093. PMID 28185817
23. Food and Drug Administration. Summary of Safety and Effectiveness Data (SSED): Edwards SAPIEN XTTM Transcatheter Heart Valve. 2016; [https://www.accessdata.fda.gov/cdrh\\_docs/pdf13/p130009s037b.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf13/p130009s037b.pdf). Accessed July, 2020.
24. Boshoff DE, Cools BL, Heying R, et al. Off-label use of percutaneous pulmonary valved stents in the right ventricular outflow tract: time to rewrite the label?. *Catheter Cardiovasc Interv.* May 2013; 81(6): 987-95. PMID 22887796
25. Faza N, Kenny D, Kavinsky C, et al. Single-center comparative outcomes of the Edwards SAPIEN and Medtronic Melody transcatheter heart valves in the pulmonary position. *Catheter Cardiovasc Interv.* Oct 01 2013; 82(4): E535-41. PMID 23008193
26. Haas NA, Moysich A, Neudorf U, et al. Percutaneous implantation of the Edwards SAPIEN() pulmonic valve: initial results in the first 22 patients. *Clin Res Cardiol.* Feb 2013; 102(2): 119-28. PMID 22932954
27. Kenny D, Hijazi ZM, Kar S, et al. Percutaneous implantation of the Edwards SAPIEN transcatheter heart valve for conduit failure in the pulmonary position: early phase 1 results from an international multicenter clinical trial. *J Am Coll Cardiol.* Nov 15 2011; 58(21): 2248-56. PMID 22078433
28. Odemis E, Guzeltas A, Saygi M, et al. Percutaneous pulmonary valve implantation using Edwards SAPIEN transcatheter heart valve in different types of conduits: initial results of a single center experience. *Congenit Heart Dis.* Sep-Oct 2013; 8(5): 411-7. PMID 23448542
29. Demkow M, Ruzyllo W, Biernacka EK, et al. Percutaneous Edwards SAPIEN() valve implantation for significant pulmonary regurgitation after previous surgical repair with a right ventricular outflow patch. *Catheter Cardiovasc Interv.* Feb 15 2014; 83(3): 474-81. PMID 23804542
30. McElhinney DB, Cabalka AK, Aboulhosn JA, et al. Transcatheter Tricuspid Valve-in-Valve Implantation for the Treatment of Dysfunctional Surgical Bioprosthetic Valves: An International, Multicenter Registry Study. *Circulation.* Apr 19 2016; 133(16): 1582-93. PMID 26994123
31. Cheatham SL, Holzer RJ, Chisolm JL, et al. The Medtronic Melody(R) transcatheter pulmonary valve implanted at 24-mm diameter--it works. *Catheter Cardiovasc Interv.* Nov 01 2013; 82(5): 816-23. PMID 23359563
32. Meadows JJ, Moore PM, Berman DP, et al. Use and performance of the Melody Transcatheter Pulmonary Valve in native and postsurgical, nonconduit right ventricular outflow tracts. *Circ Cardiovasc Interv.* Jun 2014; 7(3): 374-80. PMID 24867892
33. Malekzadeh-Milani S, Ladouceur M, Cohen S, et al. Results of transcatheter pulmonary valvulation in native or patched right ventricular outflow tracts. *Arch Cardiovasc Dis.* Nov 2014; 107(11): 592-8. PMID 25218009



34. Food and Drug Administration. Summary of Safety and Probable Benefit: Melody Transcatheter Pulmonary Valve and Ensemble Transcatheter Valve Delivery System. 2010; [https://www.accessdata.fda.gov/cdrh\\_docs/pdf8/H080002b.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf8/H080002b.pdf). Accessed July, 2020.
35. McElhinney DB, Cheatham JP, Jones TK, et al. Stent fracture, valve dysfunction, and right ventricular outflow tract reintervention after transcatheter pulmonary valve implantation: patient-related and procedural risk factors in the US Melody Valve Trial. *Circ Cardiovasc Interv.* Dec 01 2011; 4(6): 602-14. PMID 22075927
36. McElhinney DB, Benson LN, Eicken A, et al. Infective endocarditis after transcatheter pulmonary valve replacement using the Melody valve: combined results of 3 prospective North American and European studies. *Circ Cardiovasc Interv.* Jun 2013; 6(3): 292-300. PMID 23735475
37. Malekzadeh-Milani S, Ladouceur M, Iserin L, et al. Incidence and outcomes of right-sided endocarditis in patients with congenital heart disease after surgical or transcatheter pulmonary valve implantation. *J Thorac Cardiovasc Surg.* Nov 2014; 148(5): 2253-9. PMID 25218536
38. Boudjemline Y, Malekzadeh-Milani S, Patel M, et al. Predictors and outcomes of right ventricular outflow tract conduit rupture during percutaneous pulmonary valve implantation: a multicentre study. *EuroIntervention.* Jan 22 2016; 11(9): 1053-62. PMID 25244126
39. Morray BH, McElhinney DB, Cheatham JP, et al. Risk of coronary artery compression among patients referred for transcatheter pulmonary valve implantation: a multicenter experience. *Circ Cardiovasc Interv.* Oct 01 2013; 6(5): 535-42. PMID 24065444
40. Fraisse A, Assaidi A, Mauri L, et al. Coronary artery compression during intention to treat right ventricle outflow with percutaneous pulmonary valve implantation: incidence, diagnosis, and outcome. *Catheter Cardiovasc Interv.* Jun 01 2014; 83(7): E260-8. PMID 24619978
41. Van Dijk I, Budts W, Cools B, et al. Infective endocarditis of a transcatheter pulmonary valve in comparison with surgical implants. *Heart.* May 15 2015; 101(10): 788-93. PMID 25539944
42. Malekzadeh-Milani S, Ladouceur M, Patel M, et al. Incidence and predictors of Melody(R) valve endocarditis: a prospective study. *Arch Cardiovasc Dis.* Feb 2015; 108(2): 97-106. PMID 25445752
43. O'Donnell C, Holloway R, Tilton E, et al. Infective endocarditis following Melody valve implantation: comparison with a surgical cohort. *Cardiol Young.* Mar 2017; 27(2): 294-301. PMID 27161361
44. Hijazi ZM, Ruiz CE, Zahn E, et al. SCAI/AATS/ACC/STS Operator and Institutional Requirements for Transcatheter Valve Repair and Replacement, Part III: Pulmonic Valve. *J Am Coll Cardiol.* Jun 16 2015; 65(23): 2556-63. PMID 25819263
45. Nishimura RA, Otto CM, Bonow RO, et al. 2014 AHA/ACC guideline for the management of patients with valvular heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol.* Jun 10 2014; 63(22): e57-185. PMID 24603191
46. Nishimura RA, Otto CM, Bonow RO, et al. 2017 AHA/ACC Focused Update of the 2014 AHA/ACC Guideline for the Management of Patients With Valvular Heart Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol.* Jul 11 2017; 70(2): 252-289. PMID 28315732
47. Warnes CA, Williams RG, Bashore TM, et al. ACC/AHA 2008 guidelines for the management of adults with congenital heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Develop Guidelines on the Management of Adults With Congenital Heart Disease). Developed in Collaboration With the American Society of Echocardiography, Heart Rhythm Society, International Society for Adult Congenital Heart Disease, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. *J Am Coll Cardiol.* Dec 02 2008; 52(23): e143-e263. PMID 19038677
48. Bhatt AB, Foster E, Kuehl K, et al. Congenital heart disease in the older adult: a scientific statement from the American Heart Association. *Circulation.* May 26 2015; 131(21): 1884-931. PMID 25896865



## History

Date	Comments
01/10/12	New Policy – Policy created with literature search through June 15, 2011; considered medically necessary for patients who are high risk for open surgery and are poor surgical candidates due to multiple prior thoracotomies for open heart surgery. Considered investigational for all other indications. Clinical vetting information added.
09/27/12	Update Coding Section – ICD-10 codes are now effective 10/01/2014.
01/29/13	Replace policy. Policy updated with literature review, references 4, 5, 13-15, 17 added. Medically necessary statement amended to include “when performed according to FDA-approved indications”.
01/21/14	Replace policy. Policy updated with literature review, references 4, 5, 13-15, 17 added. Medically necessary statement amended to include “when performed according to FDA-approved indications”. Policy updated with literature review through September 30, 2013. References 13, 14, 16, 18, 21 added. No change to policy statement. Remove ICD-9 procedure and diagnosis codes; remove all ICD-10 codes except 02RH4JZ (which specifically applies) – these will not be used for adjudication.
09/23/14	Update Related Policies. Add 2.02.30.
01/28/15	Annual Review. Policy updated with literature review through September 23, 2014. References 13, 16-19, and 26-31 added; others renumbered. Policy statement unchanged.
01/19/16	Coding update. New CPT code 33477, effective 1/1/16, added to policy.
02/01/16	Coding update. Added 93799.
09/01/16	Annual Review, approved August 9, 2016. Policy updated with literature review through April 28, 2016; references 3, 5, 7-8, 16, 19, and 36-37 added. Policy statement unchanged. CPT coding updated.
12/01/17	Annual Review, approved November 14, 2017. Policy updated with literature review through October 2017. References 42-44 added. Policy statement changed to include specific FDA approved device indications for pulmonary valve regurgitation and stenosis. Removed CPT codes 0262T and 93799.
08/01/18	Annual Review, approved July 10, 2018. Policy updated with literature review through May 2017; references 9, 22, and 43 added. Clinical input was obtained and policy statement changed to: Transcatheter pulmonary valve implantation is considered medically necessary for patients with congenital heart disease and current right ventricular outflow tract obstruction or regurgitation when specified indications are met.
04/01/19	Minor update, added Documentation Requirements section.



Date	Comments
09/01/19	Annual Review, approved August 22, 2019. Policy updated with literature review through April 2019, no references added. Policy statements unchanged.
09/01/20	Annual Review, approved August 4, 2020. Policy updated with literature review through April, 2020, no references added. Policy statements unchanged.

**Disclaimer:** This medical policy is a guide in evaluating the medical necessity of a particular service or treatment. The Company adopts policies after careful review of published peer-reviewed scientific literature, national guidelines and local standards of practice. Since medical technology is constantly changing, the Company reserves the right to review and update policies as appropriate. Member contracts differ in their benefits. Always consult the member benefit booklet or contact a member service representative to determine coverage for a specific medical service or supply. CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). ©2020 Premera All Rights Reserved.

**Scope:** Medical policies are systematically developed guidelines that serve as a resource for Company staff when determining coverage for specific medical procedures, drugs or devices. Coverage for medical services is subject to the limits and conditions of the member benefit plan. Members and their providers should consult the member benefit booklet or contact a customer service representative to determine whether there are any benefit limitations applicable to this service or supply. This medical policy does not apply to Medicare Advantage.



**Discrimination is Against the Law**

Premera Blue Cross complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex. Premera does not exclude people or treat them differently because of race, color, national origin, age, disability or sex.

Premera:

- Provides free aids and services to people with disabilities to communicate effectively with us, such as:
  - Qualified sign language interpreters
  - Written information in other formats (large print, audio, accessible electronic formats, other formats)
- Provides free language services to people whose primary language is not English, such as:
  - Qualified interpreters
  - Information written in other languages

If you need these services, contact the Civil Rights Coordinator.

If you believe that Premera has failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, or sex, you can file a grievance with:

Civil Rights Coordinator - Complaints and Appeals  
PO Box 91102, Seattle, WA 98111  
Toll free 855-332-4535, Fax 425-918-5592, TTY 800-842-5357  
Email AppealsDepartmentInquiries@Premera.com

You can file a grievance in person or by mail, fax, or email. If you need help filing a grievance, the Civil Rights Coordinator is available to help you.

You can also file a civil rights complaint with the U.S. Department of Health and Human Services, Office for Civil Rights, electronically through the Office for Civil Rights Complaint Portal, available at <https://ocrportal.hhs.gov/ocr/portal/lobby.jsf>, or by mail or phone at: U.S. Department of Health and Human Services  
200 Independence Avenue SW, Room 509F, HHH Building  
Washington, D.C. 20201, 1-800-368-1019, 800-537-7697 (TDD)  
Complaint forms are available at <http://www.hhs.gov/ocr/office/file/index.html>.

**Getting Help in Other Languages**

**This Notice has Important Information.** This notice may have important information about your application or coverage through Premera Blue Cross. There may be key dates in this notice. You may need to take action by certain deadlines to keep your health coverage or help with costs. You have the right to get this information and help in your language at no cost. Call 800-722-1471 (TTY: 800-842-5357).

**አማርኛ (Amharic):**

ይህ ማስታወቂያ አስፈላጊ መረጃ ይዟል። ይህ ማስታወቂያ ስለ ማመልከቻዎ ወይም የ Premera Blue Cross ሽፋን አስፈላጊ መረጃ ሊኖረው ይችላል። በዚህ ማስታወቂያ ውስጥ ቁልፍ ቀዳሾች ሊኖሩ ይችላሉ። የጤና ሽፋንዎን ለመጠበቅና በአስፋፈል እርዳታ ለማግኘት በተውሰኑ የጊዜ ገደቦች እርምጃ መውሰድ ይገባዎት ይሆናል። ይህን መረጃ እንዲያገኙ እና የለምንም ክፍያ በቋንቋዎ እርዳታ እንዲያገኙ መሰታወቅ አለዎት። በስልክ ቁጥር 800-722-1471 (TTY: 800-842-5357) ይደውሉ።

**العربية (Arabic):**

يحتوي هذا الإشعار معلومات هامة. قد يحوي هذا الإشعار معلومات مهمة بخصوص طلبك أو التغطية التي تزيد الحصول عليها من خلال Premera Blue Cross. قد تكون هناك تواريخ مهمة في هذا الإشعار. وقد تحتاج لاتخاذ إجراء في تاريخ معينة للحفاظ على تغطيتك الصحية أو للمساعدة في دفع التكاليف. يحق لك الحصول على هذه المعلومات والمساعدة بلغتك دون تكبد أية تكلفة. اتصل بـ 800-722-1471 (TTY: 800-842-5357)

**中文 (Chinese):**

**本通知有重要的訊息。**本通知可能有關於您透過 Premera Blue Cross 提交的申請或保險的重要訊息。本通知內可能有重要日期。您可能需要在截止日期之前採取行動，以保留您的健康保險或者費用補貼。您有權利免費以您的母語得到本訊息和幫助。請撥電話 800-722-1471 (TTY: 800-842-5357)。

**Oromoo (Cushite):**

**Beeksisni kun odeeffannoo barbaachisaa qaba.** Beeksisni kun sagantaa yookan karaa Premera Blue Cross tiin tajaajila keessan ilaalchisee odeeffannoo barbaachisaa qabaachuu danda'a. Guyyaawwan murteessaa ta'an beeksisa kana keessatti ilaalaa. Tarii kaffaltiidhaan deeggaramuuf yookan tajaajila fayyaa keessaniif guyyaa dhumaa irratti wanti raawwattan jiraachuu danda'a. Kaffaltii irraa bilisa haala ta'een afaan keessaniin odeeffannoo argachuu fi deeggarsa argachuuf mirga ni qabaattu. Lakkoofsa bilbilaa 800-722-1471 (TTY: 800-842-5357) tii bilbilaa.

**Français (French):**

**Cet avis a d'importantes informations.** Cet avis peut avoir d'importantes informations sur votre demande ou la couverture par l'intermédiaire de Premera Blue Cross. Le présent avis peut contenir des dates clés. Vous devez peut-être prendre des mesures par certains délais pour maintenir votre couverture de santé ou d'aide avec les coûts. Vous avez le droit d'obtenir cette information et de l'aide dans votre langue à aucun coût. Appelez le 800-722-1471 (TTY: 800-842-5357).

**Kreyòl ayisyen (Creole):**

**Avi sila a gen Enfòmasyon Enpòtan ladann.** Avi sila a kapab genyen enfòmasyon enpòtan konsènan aplikasyon w lan oswa konsènan kouvèti asirans lan atravè Premera Blue Cross. Kapab genyen dat ki enpòtan nan avi sila a. Ou ka gen pou pran kèk aksyon avan sèten dat limit pou ka kenbe kouvèti asirans sante w la oswa pou yo ka ede w avèk depans yo. Se dwa w pou resewva enfòmasyon sa a ak asistans nan lang ou pale a, san ou pa gen pou peye pou sa. Rele nan 800-722-1471 (TTY: 800-842-5357).

**Deutsche (German):**

**Diese Benachrichtigung enthält wichtige Informationen.** Diese Benachrichtigung enthält unter Umständen wichtige Informationen bezüglich Ihres Antrags auf Krankenversicherungsschutz durch Premera Blue Cross. Suchen Sie nach eventuellen wichtigen Terminen in dieser Benachrichtigung. Sie könnten bis zu bestimmten Stichtagen handeln müssen, um Ihren Krankenversicherungsschutz oder Hilfe mit den Kosten zu behalten. Sie haben das Recht, kostenlose Hilfe und Informationen in Ihrer Sprache zu erhalten. Rufen Sie an unter 800-722-1471 (TTY: 800-842-5357).

**Hmoob (Hmong):**

**Tsab ntawv tshaj xo no muaj cov ntshiab lus tseem ceeb.** Tej zaum tsab ntawv tshaj xo no muaj cov ntshiab lus tseem ceeb txog koj daim ntawv thov kev pab los yog koj qhov kev pab cuam hnuv ntawm Premera Blue Cross. Tej zaum muaj cov hnuv tseem ceeb uas sau rau hauv daim ntawv no. Tej zaum koj kuj yuav tau ua qee yam uas peb kom koj ua tsis pub dhau cov caij nyoog uas teev tseg rau hauv daim ntawv no mas koj thiaj yuav tau txais kev pab cuam kho mob los yog kev pab them tej nqi kho mob ntawd. Koj muaj cai kom lawv muab cov ntshiab lus no uas tau muab sau ua koj hom lus pub dawb rau koj. Hu rau 800-722-1471 (TTY: 800-842-5357).

**Iloko (Ilocano):**

**Daytoy a Pakdaar ket naglaon iti Napateg nga Impormasion.** Daytoy a pakdaar mabalin nga adda ket naglaon iti napateg nga impormasion maipanggep iti aplikasyonyo wenno coverage babaen iti Premera Blue Cross. Daytoy ket mabalin dagiti importante a petsa iti daytoy a pakdaar. Mabalin nga adda rumbeng nga aramidenyo nga addang sakbay dagiti partikular a naituding nga aldaw tapno mapagtalinaedyo ti coverage ti salun-atyto wenno tulong kadagiti gastos. Adda karbenganyo a mangala iti daytoy nga impormasion ken tulong iti bukodyo a pagsasao nga awan ti bayadanyo. Tumawag iti numero nga 800-722-1471 (TTY: 800-842-5357).

**Italiano (Italian):**

**Questo avviso contiene informazioni importanti.** Questo avviso può contenere informazioni importanti sulla tua domanda o copertura attraverso Premera Blue Cross. Potrebbero esserci date chiave in questo avviso. Potrebbe essere necessario un tuo intervento entro una scadenza determinata per consentirti di mantenere la tua copertura o sovvenzione. Hai il diritto di ottenere queste informazioni e assistenza nella tua lingua gratuitamente. Chiama 800-722-1471 (TTY: 800-842-5357).

**日本語 (Japanese):**

この通知には重要な情報が含まれています。この通知には、Premera Blue Cross の申請または補償範囲に関する重要な情報が含まれている場合があります。この通知に記載されている可能性がある重要な日付をご確認ください。健康保険や有料サポートを維持するには、特定の期日までに行動を取らなければならない場合があります。ご希望の言語による情報とサポートが無料で提供されます。800-722-1471 (TTY: 800-842-5357)までお電話ください。

**한국어 (Korean):**

본 통지서에는 중요한 정보가 들어 있습니다. 즉 이 통지서는 귀하의 신청에 관하여 그리고 Premera Blue Cross 를 통한 커버리지에 관한 정보를 포함하고 있을 수 있습니다. 본 통지서에는 핵심이 되는 날짜들이 있을 수 있습니다. 귀하의 건강 커버리지를 계속 유지하거나 비용을 절감하기 위해서 일정한 마감일까지 조치를 취해야 할 필요가 있을 수 있습니다. 귀하의 이러한 정보와 도움을 귀하의 언어로 비용 부담없이 얻을 수 있는 권리가 있습니다. 800-722-1471 (TTY: 800-842-5357) 로 전화하십시오.

**ລາວ (Lao):**

ແຈ້ງການນີ້ມີຂໍ້ມູນສໍາຄັນ. ແຈ້ງການນີ້ອາດຈະມີຂໍ້ມູນສໍາຄັນກ່ຽວກັບຄໍາອ້ອງສະໝັກ ຫຼື ຄວາມຄົມຄອງປະກັນໄພຂອງທ່ານຜ່ານ Premera Blue Cross. ອາດຈະມີວັນທີ່ສໍາຄັນໃນແຈ້ງການນີ້. ທ່ານອາດຈະຈໍາເປັນຕ້ອງດໍາເນີນການຕາມກຳນົດ ເວລາສະເພາະເພື່ອຮັກສາຄວາມຄົມຄອງປະກັນສະພາບ ຫຼື ຄວາມຊ່ວຍເຫຼືອເວັ້ນເວີ້ ຄ່າໃຊ້ຈ່າຍຂອງທ່ານໄດ້. ທ່ານມີສິດໄດ້ຮັບຂໍ້ມູນນີ້ ແລະ ຄວາມຊ່ວຍເຫຼືອເປັນພາສາຂອງທ່ານໂດຍບໍ່ເສຍຄ່າ. ໃຫ້ໃບທາ 800-722-1471 (TTY: 800-842-5357).

**ភាសាខ្មែរ (Khmer):**

សេចក្តីជូនដំណឹងនេះមានព័ត៌មានយ៉ាងសំខាន់។ សេចក្តីជូនដំណឹងនេះប្រហែលជាមានព័ត៌មានយ៉ាងសំខាន់អំពីទម្រង់បែបបទ ឬការរៀបចំរបស់អ្នកកាមរយ: Premera Blue Cross ។ ប្រហែលជាមាន កាលបរិច្ឆេទសំខាន់នៅក្នុងសេចក្តីជូនដំណឹងនេះ។ អ្នកប្រហែលជាត្រូវការបញ្ជាក់សមត្ថភាព ដល់កិច្ចការផ្ទៃក្នុងនានា នៃស្ត្រីនិងក្មេងៗក្នុងគ្រួសាររបស់អ្នក ឬប្រាក់ចំណូលចេញថ្លៃ។ អ្នកមានសិទ្ធិទទួលបានព័ត៌មាននេះ និងដំណោះស្រាយនៅក្នុងភាសារបស់អ្នកដោយមិនអស់លុយឡើយ។ សូមទូរស័ព្ទ 800-722-1471 (TTY: 800-842-5357)។

**ਪੰਜਾਬੀ (Punjabi):**

ਇਸ ਨੋਟਿਸ ਵਿਚ ਖਾਸ ਜਾਣਕਾਰੀ ਹੈ. ਇਸ ਨੋਟਿਸ ਵਿਚ Premera Blue Cross ਵਲੋਂ ਤੁਹਾਡੀ ਕਵਰੇਜ ਅਤੇ ਅਰਜੀ ਬਾਰੇ ਮਹੱਤਵਪੂਰਨ ਜਾਣਕਾਰੀ ਹੋ ਸਕਦੀ ਹੈ . ਇਸ ਨੋਟਿਸ ਨਵ ਖਾਸ ਤਾਰੀਖਾਂ ਹੋ ਸਕਦੀਆਂ ਹਨ. ਜੇਕਰ ਤੁਸੀਂ ਜਸਰਤ ਕਵਰੇਜ ਰਿੱਖਣੀ ਹੋਵੇ ਜਾਂ ਓਸ ਦੀ ਲਾਗਤ ਜਵਿੱਚ ਮਦਦ ਦੇ ਇਕੱਠ ਹੋ ਤਾਂ ਤੁਹਾਨੂੰ ਅੰਤਮ ਤਾਰੀਖ ਤੋਂ ਪਹਿਲਾਂ ਢੁੱਝ ਖਾਸ ਕਦਮ ਚੁੱਕਣ ਦੀ ਲੋੜ ਹੋ ਸਕਦੀ ਹੈ ,ਤੁਹਾਨੂੰ ਮੁਫਤ ਵਿੱਚ ਤੋਂ ਅਪਣੀ ਭਾਸ਼ਾ ਵਿੱਚ ਜਾਣਕਾਰੀ ਅਤੇ ਮਦਦ ਪ੍ਰਾਪਤ ਕਰਨ ਦਾ ਅਧਿਕਾਰ ਹੈ ,ਕਾਲ 800-722-1471 (TTY: 800-842-5357).

**فارسی (Farsi):**

این اعلامیه حاوی اطلاعات مهم میباشد. این اعلامیه ممکن است حاوی اطلاعات مهم درباره فرم تقاضا و یا پوشش بیمه ای شما از طریق Premera Blue Cross باشد. به تاریخ های مهم در این اعلامیه توجه نمایید. شما ممکن است برای حفظ پوشش بیمه تان یا کمک در پرداخت هزینه های درمانی تان، به تاریخ های مشخصی برای انجام کارهای خاصی احتیاج داشته باشید. شما حق این را دارید که این اطلاعات و کمک را به زبان خود به طور رایگان دریافت نمایید. برای کسب اطلاعات با شماره 800-722-1471 (کلیر بران TTY تماس باشماره 800-842-5357) تماس برقرار نمایید.

**Polskie (Polish):**

To ogłoszenie może zawierać ważne informacje. To ogłoszenie może zawierać ważne informacje odnośnie Państwa wniosku lub zakresu świadczeń poprzez Premera Blue Cross. Prosimy zwrócić uwagę na kluczowe daty, które mogą być zawarte w tym ogłoszeniu aby nie przekroczyć terminów w przypadku utrzymania polisy ubezpieczeniowej lub pomocy związanej z kosztami. Macie Państwo prawo do bezpłatnej informacji we własnym języku. Zadzwońcie pod 800-722-1471 (TTY: 800-842-5357).

**Português (Portuguese):**

Este aviso contém informações importantes. Este aviso poderá conter informações importantes a respeito de sua aplicação ou cobertura por meio do Premera Blue Cross. Poderão existir datas importantes neste aviso. Talvez seja necessário que você tome providências dentro de determinados prazos para manter sua cobertura de saúde ou ajuda de custos. Você tem o direito de obter esta informação e ajuda em seu idioma e sem custos. Ligue para 800-722-1471 (TTY: 800-842-5357).

**Română (Romanian):**

Prezenta notificare conține informații importante privind cererea sau acoperirea asigurării dumneavoastră de sănătate prin Premera Blue Cross. Pot exista date cheie în această notificare. Este posibil să fie nevoie să acționați până la anumite termene limită pentru a vă menține acoperirea asigurării de sănătate sau asistența provizorie la costuri. Aveți dreptul de a obține gratuit aceste informații și ajutor în limba dumneavoastră. Sunați la 800-722-1471 (TTY: 800-842-5357).

**Русский (Russian):**

Настоящее уведомление содержит важную информацию. Это уведомление может содержать важную информацию о вашем заявлении или страховом покрытии через Premera Blue Cross. В настоящем уведомлении могут быть указаны ключевые даты. Вам, возможно, потребуется принять меры к определенным предельным срокам для сохранения страхового покрытия или помощи с расходами. Вы имеете право на бесплатное получение этой информации и помощь на вашем языке. Звоните по телефону 800-722-1471 (TTY: 800-842-5357).

**Fa'asamoa (Samoan):**

Atonu ua iai i lenei fa'asilasilaga ni fa'amatalaga e sili ona taua e tatau ona e malamalama i ai. O lenei fa'asilasilaga o se fesoasoani e fa'amatala atili i ai i le tulaga o le polokalame, Premera Blue Cross, ua e tau fia maua atu i ai. Fa'amolemole, ia e iloilo fa'alelei i aso fa'apitoa olo'o iai i lenei fa'asilasilaga taua. Masalo o le'a iai ni feau e tatau ona e faia ao le'i aulia le aso ua ta'ua i lenei fa'asilasilaga ina ia e iai pea ma maua fesoasoani mai ai i le polokalame a le Malo olo'o e iai i ai. Olo'o iai iate oe le aia tatau e maua atu i lenei fa'asilasilaga ma lenei fa'matalaga i legagana e te malamalama i ai aunoa ma se togiga tupe. Vili atu i le telefoni 800-722-1471 (TTY: 800-842-5357).

**Español (Spanish):**

Este Aviso contiene información importante. Es posible que este aviso contenga información importante acerca de su solicitud o cobertura a través de Premera Blue Cross. Es posible que haya fechas clave en este aviso. Es posible que deba tomar alguna medida antes de determinadas fechas para mantener su cobertura médica o ayuda con los costos. Usted tiene derecho a recibir esta información y ayuda en su idioma sin costo alguno. Llame al 800-722-1471 (TTY: 800-842-5357).

**Tagalog (Tagalog):**

Ang Paunawa na ito ay naglalaman ng mahalagang impormasyon tungkol sa iyong aplikasyon o pagsakop sa pamamagitan ng Premera Blue Cross. Maaaring may mga mahalagang petsa dito sa paunawa. Maaring mangailangan ka na magsagawa ng hakbang sa ilang mga itinakdang panahon upang mapanatili ang iyong pagsakop sa kalusugan o tulong na walang gastos. May karapatan ka na makakuha ng ganiitong impormasyon at tulong sa iyong wika ng walang gastos. Tumawag sa 800-722-1471 (TTY: 800-842-5357).

**ไทย (Thai):**

ประกาศนี้มีข้อมูลสำคัญ ประกาศนี้อาจมีข้อมูลที่สำคัญเกี่ยวกับกาการสมัครหรือขอบเขตประกันสุขภาพของคุณผ่าน Premera Blue Cross และอาจมีกำหนดการในประกาศนี้ คุณอาจจะต้องดำเนินการภายในกำหนดระยะเวลาที่แน่นอนเพื่อจะรักษาการประกันสุขภาพของคุณหรือการช่วยเหลือที่มีค่าใช้จ่าย คุณมีสิทธิที่จะได้รับข้อมูลและความช่วยเหลือนี้ในภาษาของคุณโดยไม่มีค่าใช้จ่าย โทร 800-722-1471 (TTY: 800-842-5357)

**Український (Ukrainian):**

Це повідомлення містить важливу інформацію. Це повідомлення може містити важливу інформацію про Ваше звернення щодо страховального покриття через Premera Blue Cross. Зверніть увагу на ключові дати, які можуть бути вказані у цьому повідомленні. Існує імовірність того, що Вам треба буде здійснити певні кроки у конкретні кінцеві строки для того, щоб зберегти Ваше медичне страхування або отримати фінансову допомогу. У Вас є право на отримання цієї інформації та допомоги безкоштовно на Вашій рідній мові. Дзвоніть за номером телефону 800-722-1471 (TTY: 800-842-5357).

**Tiếng Việt (Vietnamese):**

Thông báo này cung cấp thông tin quan trọng. Thông báo này có thông tin quan trọng về đơn xin tham gia hoặc hợp đồng bảo hiểm của quý vị qua chương trình Premera Blue Cross. Xin xem ngày quan trọng trong thông báo này. Quý vị có thể phải thực hiện theo thông báo đúng trong thời hạn để duy trì bảo hiểm sức khỏe hoặc được trợ giúp thêm về chi phí. Quý vị có quyền được biết thông tin này và được trợ giúp bằng ngôn ngữ của mình miễn phí. Xin gọi số 800-722-1471 (TTY: 800-842-5357).