

## MEDICAL POLICY – 7.01.136

Radiofrequency Ablation of the Renal Sympathetic Nerves  
as a Treatment for Resistant Hypertension

BCBSA Ref. Policy: 7.01.136

Effective Date: Dec. 1, 2018

Last Revised: Nov. 6, 2018


Replaces: N/A

RELATED MEDICAL POLICIES:

None

Select a hyperlink below to be directed to that section.

[POLICY CRITERIA](#) | [CODING](#) | [RELATED INFORMATION](#)  
[EVIDENCE REVIEW](#) | [REFERENCES](#) | [HISTORY](#)

 Clicking this icon returns you to the hyperlinks menu above.

---

## Introduction

Treating high blood pressure often involves making lifestyle changes and using medications. A newer technique that destroys parts of certain nerves has been proposed as a way to lower blood pressure when medication doesn't work. For this procedure, a narrow tube (catheter) is inserted in an artery near the groin. The catheter is threaded up to the artery that supplies blood to the kidney and then maneuvered to the nerves in the wall of this kidney artery. A weak electrical current is released, and the heat kills parts of these nerves. It's thought that because these nerves are damaged they will reduce the kidney's production of specific hormones. The decreased amount of this hormone is thought to decrease blood pressure over time. Medical studies have conflicting results about whether this treatment works. This service is considered investigational (unproven).

**Note:** The Introduction section is for your general knowledge and is not to be taken as policy coverage criteria. The rest of the policy uses specific words and concepts familiar to medical professionals. It is intended for providers. A provider can be a person, such as a doctor, nurse, psychologist, or dentist. A provider also can be a place where medical care is given, like a hospital, clinic, or lab. This policy informs them about when a service may be covered.

## Policy Coverage Criteria

Service	Investigational
<b>Radiofrequency ablation of the renal sympathetic nerves</b>	<b>Radiofrequency ablation of the renal sympathetic nerves is considered investigational for the treatment of resistant hypertension.</b>

## Coding

Code	Description
<b>CPT</b>	
0338T	Transcatheter renal sympathetic denervation, percutaneous approach including arterial puncture, selective catheter placement(s) renal artery(ies), fluoroscopy, contrast injection(s), intraprocedural roadmapping and radiological supervision and interpretation, including pressure gradient measurements, flush aortogram and diagnostic renal angiography when performed; unilateral
0339T	Transcatheter renal sympathetic denervation, percutaneous approach including arterial puncture, selective catheter placement(s) renal artery(ies), fluoroscopy, contrast injection(s), intraprocedural roadmapping and radiological supervision and interpretation, including pressure gradient measurements, flush aortogram and diagnostic renal angiography when performed; bilateral

**Note:** CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). HCPCS codes, descriptions and materials are copyrighted by Centers for Medicare Services (CMS).

## Related Information

N/A

## Evidence Review



## Description

Radiofrequency ablation (RFA) of the renal sympathetic nerves is thought to decrease both the afferent sympathetic signals from the kidney to the brain and the efferent signals from the brain to the kidney. This procedure decreases sympathetic activation, decreases vasoconstriction, and decreases activation of the renin-angiotensin system. RFA of the renal sympathetic nerves may act as a nonpharmacologic treatment for hypertension and has been proposed as a treatment option for patients with resistant hypertension.

## Background

### *Resistant Hypertension*

Hypertension is estimated to affect approximately 30% of the population in the United States.<sup>1</sup> It accounts for a high burden of morbidity related to strokes, ischemic heart disease, kidney disease, and peripheral arterial disease. Resistant hypertension is defined as elevated blood pressure, despite treatment with at least 3 antihypertensive agents at optimal doses. Resistant hypertension is also a relatively common condition, given the large number of individuals with hypertension. In large clinical trials of hypertension treatment, 20% to 30% of participants meet the definition for resistant hypertension, and in tertiary care hypertension clinics, the prevalence is estimated at 11% to 18%.<sup>1</sup> Resistant hypertension is associated with a higher risk for adverse outcomes such as stroke, myocardial infarction, heart failure, and kidney failure.

A number of factors may contribute to uncontrolled hypertension, and they should be considered and addressed in all patients with hypertension before labeling a patient resistant. They include nonadherence to medications, excessive salt intake, inadequate doses of medications, excess alcohol intake, volume overload, drug-induced hypertension, and other forms of secondary hypertension.<sup>2</sup> Also, sometimes it is necessary to address comorbid conditions (ie, obstructive sleep apnea) to control blood pressure adequately.

## Treatment

Treatment for resistant hypertension is mainly intensified drug therapy, sometimes with the use of nontraditional antihypertensive medications such as spironolactone and/or minoxidil. However, control of resistant hypertension with additional medications is often challenging and can lead to high costs and frequent adverse effects of treatment. As a result, there is a large unmet need for additional treatments that can control resistant hypertension.



Nonpharmacologic interventions for resistant hypertension include modulation of the baroreflex receptor and/or radiofrequency (RF) denervation of the renal nerves.

### ***RF Denervation of the Renal Sympathetic Nerves***

Increased sympathetic nervous system activity has been linked to essential hypertension. Surgical sympathectomy has been shown to be effective in reducing blood pressure but is limited by the adverse effects of surgery and was largely abandoned after effective medications for hypertension became available. The renal sympathetic nerves arise from the thoracic nerve roots and innervate the renal artery, the renal pelvis, and the renal parenchyma. Radiofrequency ablation (RFA) is thought to decrease both the afferent sympathetic signals from the kidney to the brain and the efferent signals from the brain to the kidney. This procedure decreases sympathetic activation, decreases vasoconstriction, and decreases activation of the renin-angiotensin system.<sup>3</sup>

The procedure is performed percutaneously with access at the femoral artery. A flexible catheter is threaded into the renal artery, and a controlled energy source, most commonly low-power RF energy, is delivered to the arterial walls where the renal sympathetic nerves are located. Once adequate RF energy has been delivered to ablate the sympathetic nerves, the catheter is removed.

## **Summary of Evidence**

For individuals who have hypertension resistant to standard medical management who receive RFA of the renal sympathetic nerves, the evidence includes at least 10 randomized controlled trials (RCTs), numerous systematic reviews of the RCTs, as well as multiple nonrandomized comparative studies and case series. Relevant outcomes are symptoms, change in disease status, morbid events, medication use, and treatment-related morbidity. The largest trial, the Symplicity HTN-3 trial, used a sham-controlled design to reduce the likelihood of placebo effect and demonstrated no significant differences between renal denervation and sham-control patients in office-based or ambulatory blood pressure at 6-month follow-up. Results from Symplicity HTN-3 are supported by a subsequent sham-controlled trial. The Symplicity HTN-3 results were in contrast to other studies, including Symplicity HTN-2 and Renal Denervation for Hypertension (DENERHTN) trial, which reported efficacy in reducing blood pressure over a 6-month time period compared with a control group. Additional smaller randomized controlled trials, some of which were stopped early after results of the Symplicity HTN-3 trial became available, did not demonstrate significantly improved outcomes with renal denervation. Single-arm studies with



overlapping populations have reported improvements in blood pressure and related physiologic parameters, such as echocardiographic measures of left ventricular hypertrophy, that appear to be durable for up to 24 months of follow-up. The strongest evidence comes from sham-controlled trials, the largest of which found no significant benefits with renal denervation. Meta-analyses of the systematic reviews have also reported inconsistent findings, with most analyses showing no significant benefit in blood pressure measurements following RFA. The evidence is insufficient to determine the effects of the technology on health outcomes.

## Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in [Table 1](#).

**Table 1. Summary of Key Trials**

NCT No.	Trial Name	Planned Enrollment	Completion Date
<b>Ongoing</b>			
<a href="#">NCT02439775</a>	Global Clinical Study of Renal Denervation With the Symplicity Spyral™ Multi-electrode Renal Denervation System in Patients With Uncontrolled Hypertension on Standard Medical Therapy (SPYRAL HTN-ON MED)	106	Feb 2021
<a href="#">NCT01673516</a>	Effect of Renal Sympathetic Denervation on Resistant Hypertension and Cardiovascular Hemodynamic in Comparison to Intensive Medical Therapy Utilizing Impedance Cardiography	60	Aug 2022
<a href="#">NCT02439749</a>	Global Clinical Study of Renal Denervation With the Symplicity Spyral™ Multi-electrode Renal Denervation System in Patients With Uncontrolled Hypertension in the Absence of Antihypertensive Medications (SPYRAL HTN-OFF MED)	433	Dec 2022
<b>Unpublished</b>			
<a href="#">NCT02029885<sup>a</sup></a>	Wave IV Study: Phase II Randomized Sham Controlled Study of Renal Denervation for Subjects With Uncontrolled Hypertension	132	Mar 2018 (unknown)
<a href="#">NCT01911078</a>	Renal Sympathetic Denervation in Metabolic Syndrome (Metabolic Syndrome Study)	20	Jun 2016 (completed)



NCT No.	Trial Name	Planned Enrollment	Completion Date
<a href="#">NCT01901549</a>	Renal Denervation in Patients After Acute Coronary Syndrome	80	Jun 2016 (unknown)
<a href="#">NCT02041130</a>	Renal Sympathectomy in Heart Failure (the RESPECT-HF Study) - a Study of Renal Denervation for Heart Failure With Preserved Ejection Fraction	144	Dec 2016 (unknown)
<a href="#">NCT01522430</a>	Denervation of Renal Sympathetic Activity and Hypertension Study	120	Dec 2016 (unknown)
<a href="#">NCT02021019</a>	Renal Denervation to Improve Outcomes in Patients With End-stage Renal	100	Dec 2016 (unknown)
<a href="#">NCT01932450</a>	A Randomized, Open-label Study Investigating the Effect of Bilateral Renal Artery Sympathetic Denervation by Catheter-based Radiofrequency Ablation on Blood Pressure and Disease Progression in Autosomal Dominant Polycystic Kidney Disease	100	Jul 2015 (unknown)
<a href="#">NCT01895140</a> <sup>a</sup>	A Pragmatic Randomized Clinical Evaluation of Renal Denervation for Treatment Resistant Hypertension	104	Oct 2014 (terminated)
<a href="#">NCT01628172</a> <sup>a</sup>	Renal Sympathetic Denervation for the Management of Chronic Hypertension	96	Mar 2014 (completed)

NCT: national clinical trial

<sup>a</sup> Denotes industry-sponsored or cosponsored trial

## Practice Guidelines and Position Statements

### *American Heart Association et al*

The American Heart Association, American College of Cardiology, and American Society of Hypertension (2015) issued guidelines on the treatment of hypertension in patients with coronary artery disease.<sup>44</sup> The guidelines noted that the Symplicity HTN-3 trial did not find a significant benefit from renal denervation and stated that additional randomized controlled trials would be needed.

The American Heart Association, American College of Cardiology, and 9 additional specialty societies (2018) published joint guidelines on the prevention, detection, evaluation, and management of high blood pressure in adults.<sup>45</sup> In discussing resistant hypertension, the guidelines indicated that studies using catheter ablation of renal sympathetic nerves “have not provided sufficient evidence to recommend the use of these devices.”



## ***Joint UK Societies***

The British Hypertension Society and 3 other British medical societies (collectively, the Joint UK Societies) issued an expert consensus statement (2014) on renal denervation for resistant hypertension, which concluded: "The Joint UK Societies did not recommend the use of renal denervation for treatment of resistant hypertension in routine clinical practice but remains committed to supporting research activity in this field."<sup>46</sup>

## ***Eighth Joint National Committee***

The Eighth Joint National Committee (2014), which was appointed to provide recommendations on hypertension treatment, published an evidence-based guideline for the management of hypertension in adults.<sup>47</sup> This guideline did not discuss the use of renal denervation.

## ***European Society of Cardiology***

The European Society of Cardiology (2013) issued an expert consensus statement on catheter-based renal denervation which concluded that, based on the available evidence, renal denervation can be considered as a treatment option in "patients with resistant hypertension, whose blood pressure cannot be controlled by a combination of lifestyle modification and pharmacological therapy according to current guidelines."<sup>48</sup>

The statement indicated that patients should meet the following criteria before renal denervation is considered:

- "Office-based systolic BP [blood pressure]  $\geq 160$  mmHg ( $\geq 150$  mmHg in diabetes type 2)
- $\geq 3$  antihypertensive drugs in adequate dosage and combination (incl. diuretic)
- Lifestyle modification
- Exclusion of secondary hypertension
- Exclusion of pseudo-resistance using ABPM [ambulatory blood pressure monitoring] (average BP  $> 130$  mmHg or mean daytime BP  $> 135$  mmHg)
- Preserved renal function (GFR [glomerular filtration rate]  $\geq 45$  mL/min/1.73 m<sup>2</sup>)



- Eligible renal arteries: no polar or accessory arteries; no renal artery stenosis; no prior revascularization”

## Medicare National Coverage

There is no national coverage determination.

## Regulatory Status

No RFA devices have been approved by the U.S. Food and Drug Administration (FDA) for ablation of the renal sympathetic nerves as a treatment for hypertension. Several devices have been developed for this purpose and are in various stages of application for FDA approval (FDA product code: DQY):

- Symplicity™ Renal Denervation System (Medtronic). In April 2018, FDA approved an investigational device exemption pivotal trial, SPYRAL HTN. The trial will be randomized and sham-controlled.
- The EnligHTN™ Multi-Electrode Renal Denervation System (St. Jude Medical) is an RFA catheter using a 4-point multiablation basket design. In January 2014, the EnligHTN™ Renal Guiding Catheter was cleared for marketing by the FDA through the 510(k) process based on substantial equivalence to predicate devices for the following indication: percutaneous use through an introducer sheath to facilitate a pathway to introduce interventional and diagnostic devices into the renal arterial vasculature.
- The OneShot™ Renal Denervation System (Covidien) is an irrigated RFA balloon catheter, consisting of a spiral-shaped electrode surrounding a balloon. (In 2014, Covidien abandoned development of its OneShot™ Renal Denervation program.)
- The Vessix™ Renal Denervation System (Boston Scientific; formerly the V2 renal denervation system, Vessix Vascular) is a combination of a RF balloon catheter and bipolar RF generator technologies, intended to permit a lower voltage intervention.

Other RFA catheters (eg, Thermocouple Catheter™ [Biosense Webster]), used for other types of ablation procedures (eg, cardiac electrophysiology procedures), have been used off-label for RFA of the renal arteries.





## References

1. Acelajado MC, Calhoun DA. Resistant hypertension, secondary hypertension, and hypertensive crises: diagnostic evaluation and treatment. *Cardiol Clin*. Nov 2010;28(4):639-654. PMID 20937447
2. Doulas M, Papademetriou V, Douma S, et al. Benefits from treatment and control of patients with resistant hypertension. *Int J Hypertens*. Dec 22 2010;2011:318549. PMID 21234402
3. Zile MR, Little WC. Effects of autonomic modulation: more than just blood pressure. *J Am Coll Cardiol*. Mar 6 2012;59(10):910-912. PMID 22381426
4. Azizi M, Sapoval M, Gosse P, et al. Optimum and stepped care standardised antihypertensive treatment with or without renal denervation for resistant hypertension (DENERHTN): a multicentre, open-label, randomised controlled trial. *Lancet*. May 16 2015;385(9981):1957-1965. PMID 25631070
5. Courand PY, Pereira H, Del Giudice C, et al. Abdominal aortic calcifications influences the systemic and renal hemodynamic response to renal denervation in the DENERHTN (Renal Denervation for Hypertension) Trial. *J Am Heart Assoc*. Oct 10 2017;6(10). PMID 29018027
6. Gosse P, Cremer A, Pereira H, et al. Twenty-four-hour blood pressure monitoring to predict and assess impact of renal denervation: The DENERHTN Study (Renal Denervation for Hypertension). *Hypertension*. Mar 2017;69(3):494-500. PMID 28115517
7. Rosa J, Widimsky P, Tousek P, et al. Randomized comparison of renal denervation versus intensified pharmacotherapy including spironolactone in true-resistant hypertension: six-month results from the Prague-15 study. *Hypertension*. Feb 2015;65(2):407-413. PMID 25421981
8. Bhatt DL, Kandzari DE, O'Neill WW, et al. A controlled trial of renal denervation for resistant hypertension. *N Engl J Med*. Apr 10 2014;370(15):1393-1401. PMID 24678939
9. Bakris GL, Townsend RR, Liu M, et al. Impact of renal denervation on 24-hour ambulatory blood pressure: results from SYMPLICITY HTN-3. *J Am Coll Cardiol*. Sep 16 2014;64(11):1071-1078. PMID 24858423
10. Bakris GL, Townsend RR, Flack JM, et al. 12-month blood pressure results of catheter-based renal artery denervation for resistant hypertension: the SYMPLICITY HTN-3 trial. *J Am Coll Cardiol*. Apr 7 2015;65(13):1314-1321. PMID 25835443
11. Mahfoud F, Bakris G, Bhatt DL, et al. Reduced blood pressure-lowering effect of catheter-based renal denervation in patients with isolated systolic hypertension: data from SYMPLICITY HTN-3 and the Global SYMPLICITY Registry. *Eur Heart J*. Jan 7 2017;38(2):93-100. PMID 28158510
12. Kario K, Bhatt DL, Brar S, et al. Effect of catheter-based renal denervation on morning and nocturnal blood pressure: insights from SYMPLICITY HTN-3 and SYMPLICITY HTN-Japan. *Hypertension*. Dec 2015;66(6):1130-1137. PMID 26558819
13. Lu D, Wang K, Liu Q, et al. Reductions of left ventricular mass and atrial size following renal denervation: a meta-analysis. *Clin Res Cardiol*. Aug 2016;105(8):648-656. PMID 26838292
14. Flack JM, Bhatt DL, Kandzari DE, et al. An analysis of the blood pressure and safety outcomes to renal denervation in African Americans and Non-African Americans in the SYMPLICITY HTN-3 trial. *J Am Soc Hypertens*. Oct 2015;9(10):769-779. PMID 26362830
15. Esler MD, Krum H, Sobotka PA, et al. Renal sympathetic denervation in patients with treatment-resistant hypertension (The Symplicity HTN-2 Trial): a randomised controlled trial. *Lancet*. Dec 4 2010;376(9756):1903-1909. PMID 21093036
16. Esler MD, Krum H, Schlaich M, et al. Renal sympathetic denervation for treatment of drug-resistant hypertension: one-year results from the Symplicity HTN-2 randomized, controlled trial. *Circulation*. Dec 18 2012;126(25):2976-2982. PMID 23248063



17. Esler MD, Bohm M, Sievert H, et al. Catheter-based renal denervation for treatment of patients with treatment-resistant hypertension: 36 month results from the SYMPPLICITY HTN-2 randomized clinical trial. *Eur Heart J*. Jul 7 2014;35(26):1752-1759. PMID 24898552
18. Kario K, Ogawa H, Okumura K, et al. SYMPPLICITY HTN-Japan - first randomized controlled trial of catheter-based renal denervation in Asian patients. *Circ J*. Apr 2015;79(6):1222-1229. PMID 25912693
19. de Jager RL, de Beus E, Beeftink MM, et al. Impact of medication adherence on the effect of renal denervation: The SYMPATHY Trial. *Hypertension*. Apr 2017;69(4):678-684. PMID 28264922
20. de Jager RL, van Maarseveen EM, Bots ML, et al. Medication adherence in patients with apparent resistant hypertension: findings from the SYMPATHY trial. *Br J Clin Pharmacol*. Jan 2018;84(1):18-24. PMID 28815689
21. Oliveras A, Armario P, Clara A, et al. Spironolactone versus sympathetic renal denervation to treat true resistant hypertension: results from the DENERVHTA study - a randomized controlled trial. *J Hypertens*. Sep 2016;34(9):1863-1871. PMID 27327441
22. Mathiassen ON, Vase H, Bech JN, et al. Renal denervation in treatment-resistant essential hypertension. A randomized, sham-controlled, double-blinded 24-h blood pressure-based trial. *J Hypertens*. Aug 2016;34(8):1639-1647. PMID 27228432
23. Desch S, Okon T, Heinemann D, et al. Randomized sham-controlled trial of renal sympathetic denervation in mild resistant hypertension. *Hypertension*. Jun 2015;65(6):1202-1208. PMID 25824248
24. Fengler K, Heinemann D, Okon T, et al. Renal denervation improves exercise blood pressure: insights from a randomized, sham-controlled trial. *Clin Res Cardiol*. Jul 2016;105(7):592-600. PMID 26728060
25. Schneider S, Promny D, Sinnecker D, et al. Impact of sympathetic renal denervation: a randomized study in patients after renal transplantation (ISAR-denerve). *Nephrol Dial Transplant*. Nov 2015;30(11):1928-1936. PMID 26333545
26. Fadl Elmula FE, Hoffmann P, Larstorp AC, et al. Adjusted drug treatment is superior to renal sympathetic denervation in patients with true treatment-resistant hypertension. *Hypertension*. May 2014;63(5):991-999. PMID 24591332
27. Pokushalov E, Romanov A, Corbucci G, et al. A randomized comparison of pulmonary vein isolation with versus without concomitant renal artery denervation in patients with refractory symptomatic atrial fibrillation and resistant hypertension. *J Am Coll Cardiol*. Sep 25 2012;60(13):1163-1170. PMID 22958958
28. Coppolino G, Pisano A, Rivoli L, et al. Renal denervation for resistant hypertension. *Cochrane Database Syst Rev*. Feb 21 2017;2:CD011499. PMID 28220472
29. Chen XH, Kim S, Zeng XX, et al. Account for clinical heterogeneity in assessment of catheter-based renal denervation among resistant hypertension patients: subgroup meta-analysis. *Chin Med J (Engl)*. Jul 5 2017;130(13):1586-1594. PMID 28639575
30. Pappaccogli M, Covella M, Berra E, et al. Effectiveness of renal denervation in resistant hypertension: a meta-analysis of 11 controlled studies. *High Blood Press Cardiovasc Prev*. May 11 2018. PMID 29752703
31. Fadl Elmula FEM, Feng YM, Jacobs L, et al. Sham or no sham control: that is the question in trials of renal denervation for resistant hypertension. A systematic meta-analysis. *Blood Press*. Aug 2017;26(4):195-203. PMID 28443356
32. Sun D, Li C, Li M, et al. Renal denervation vs pharmacotherapy for resistant hypertension: a meta-analysis. *J Clin Hypertens (Greenwich)*. Aug 2016;18(8):733-740. PMID 26619813
33. Zhang X, Wu N, Yan W, et al. The effects of renal denervation on resistant hypertension patients: a meta-analysis. *Blood Press Monit*. Aug 2016;21(4):206-214. PMID 26901340
34. Yao Y, Zhang D, Qian J, et al. The effect of renal denervation on resistant hypertension: Meta-analysis of randomized controlled clinical trials. *Clin Exp Hypertens*. Mar 2016;38(3):278-286. PMID 27018652
35. Fadl Elmula FE, Jin Y, Yang WY, et al. Meta-analysis of randomized controlled trials of renal denervation in treatment-resistant hypertension. *Blood Press*. Oct 2015;24(5):263-274. PMID 26194721
36. Kwok CS, Loke YK, Pradhan S, et al. Renal denervation and blood pressure reduction in resistant hypertension: a systematic review and meta-analysis. *Open Heart*. Oct 2014;1(1):e000092. PMID 25332808



37. Pancholy SB, Shantha GP, Patel TM, et al. Meta-analysis of the effect of renal denervation on blood pressure and pulse pressure in patients with resistant systemic hypertension. *Am J Cardiol.* Sep 15 2014;114(6):856-861. PMID 25084693
38. Davis MI, Filion KB, Zhang D, et al. Effectiveness of renal denervation therapy for resistant hypertension: a systematic review and meta-analysis. *J Am Coll Cardiol.* Jul 16 2013;62(3):231-241. PMID 23644092
39. Shantha GP, Pancholy SB. Effect of renal sympathetic denervation on apnea-hypopnea index in patients with obstructive sleep apnea: a systematic review and meta-analysis. *Sleep Breath.* Mar 2015;19(1):29-34. PMID 24839239
40. Brandt MC, Mahfoud F, Reda S, et al. Renal sympathetic denervation reduces left ventricular hypertrophy and improves cardiac function in patients with resistant hypertension. *J Am Coll Cardiol.* Mar 6 2012;59(10):901-909. PMID 22381425
41. Mahfoud F, Cremers B, Janker J, et al. Renal hemodynamics and renal function after catheter-based renal sympathetic denervation in patients with resistant hypertension. *Hypertension.* Jun 25 2012;60(2):419-424. PMID 22733462
42. Ukena C, Mahfoud F, Kindermann I, et al. Cardiorespiratory response to exercise after renal sympathetic denervation in patients with resistant hypertension. *J Am Coll Cardiol.* Sep 6 2011;58(11):1176-1182. PMID 21884958
43. Ewen S, Mahfoud F, Linz D, et al. Effects of renal sympathetic denervation on exercise blood pressure, heart rate, and capacity in patients with resistant hypertension. *Hypertension.* Apr 2014;63(4):839-845. PMID 24420550
44. Rosendorff C, Lackland DT, Allison M, et al. Treatment of hypertension in patients with coronary artery disease: a scientific statement from the American Heart Association, American College of Cardiology, and American Society of Hypertension. *Circulation.* May 12 2015;131(19):e435-470. PMID 25829340
45. Brook RD, Rajagopalan S. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. A report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Soc Hypertens.* Mar 2018;12(3):238. PMID 29396104
46. Lobo MD, de Belder MA, Cleveland T, et al. Joint UK societies' 2014 consensus statement on renal denervation for resistant hypertension. *Heart.* Jan 2015;101(1):10-16. PMID 25431461
47. James PA, Oparil S, Carter BL, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA.* Feb 5 2014;311(5):507-520. PMID 24352797
48. Mahfoud F, Luscher TF, Andersson B, et al. Expert consensus document from the European Society of Cardiology on catheter-based renal denervation. *Eur Heart J.* Jul 2013;34(28):2149-2157. PMID 23620497

## History

Date	Comments
10/15/12	New Policy. Radiofrequency ablation of the renal sympathetic nerves is considered investigational for the treatment of resistant hypertension.
12/04/13	Replace policy. Policy updated with literature review through July 31, 2013. References 5, 6, 17-21 added. No change in policy statement. Codes 0338T and 0339T added to policy; codes 36251-36254 removed from policy; they are no specific to the procedure.
11/20/14	Annual Review. Policy updated with literature review through July 31, 2014. References 4-5, 8-9, 11-12, 16, 19, 29-36, 38-43, and 45 added. No change to policy statement.



Date	Comments
11/10/15	Annual Review. Policy updated with literature review through August 3, 2015; references 4-5, 8, 12-13, 16-17, 51, 54-55, and 57-58 added. Policy statement unchanged.
12/07/15	Update Related Policies. Removed 8.01.57 as it is archived.
11/01/16	Annual Review, approved October 11, 2016. Policy updated with literature review through July 25, 2016. References 9-11, 15, 17, 21-22, 24-26, and 37 added. Policy statements unchanged. Removed unlisted CPT code 64999 from coding section.
11/01/17	Annual Review, approved October 19, 2017. Policy updated with literature review through July 20, 2017; no references added. Policy statement unchanged.
12/01/18	Annual Review, approved November 6, 2018. Policy updated with literature review through July 2018; references 5-6, 11, 18-20, 28-29, 34-35, and 45 added. Policy statement unchanged.

**Disclaimer:** This medical policy is a guide in evaluating the medical necessity of a particular service or treatment. The Company adopts policies after careful review of published peer-reviewed scientific literature, national guidelines and local standards of practice. Since medical technology is constantly changing, the Company reserves the right to review and update policies as appropriate. Member contracts differ in their benefits. Always consult the member benefit booklet or contact a member service representative to determine coverage for a specific medical service or supply. CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). ©2018 Premera All Rights Reserved.

**Scope:** Medical policies are systematically developed guidelines that serve as a resource for Company staff when determining coverage for specific medical procedures, drugs or devices. Coverage for medical services is subject to the limits and conditions of the member benefit plan. Members and their providers should consult the member benefit booklet or contact a customer service representative to determine whether there are any benefit limitations applicable to this service or supply. This medical policy does not apply to Medicare Advantage.



**Discrimination is Against the Law**

Premera Blue Cross complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex. Premera does not exclude people or treat them differently because of race, color, national origin, age, disability or sex.

Premera:

- Provides free aids and services to people with disabilities to communicate effectively with us, such as:
  - Qualified sign language interpreters
  - Written information in other formats (large print, audio, accessible electronic formats, other formats)
- Provides free language services to people whose primary language is not English, such as:
  - Qualified interpreters
  - Information written in other languages

If you need these services, contact the Civil Rights Coordinator.

If you believe that Premera has failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, or sex, you can file a grievance with:

Civil Rights Coordinator - Complaints and Appeals  
PO Box 91102, Seattle, WA 98111  
Toll free 855-332-4535, Fax 425-918-5592, TTY 800-842-5357  
Email AppealsDepartmentInquiries@Premera.com

You can file a grievance in person or by mail, fax, or email. If you need help filing a grievance, the Civil Rights Coordinator is available to help you.

You can also file a civil rights complaint with the U.S. Department of Health and Human Services, Office for Civil Rights, electronically through the Office for Civil Rights Complaint Portal, available at <https://ocrportal.hhs.gov/ocr/portal/lobby.jsf>, or by mail or phone at: U.S. Department of Health and Human Services  
200 Independence Avenue SW, Room 509F, HHH Building  
Washington, D.C. 20201, 1-800-368-1019, 800-537-7697 (TDD)  
Complaint forms are available at <http://www.hhs.gov/ocr/office/file/index.html>.

**Getting Help in Other Languages**

**This Notice has Important Information.** This notice may have important information about your application or coverage through Premera Blue Cross. There may be key dates in this notice. You may need to take action by certain deadlines to keep your health coverage or help with costs. You have the right to get this information and help in your language at no cost. Call 800-722-1471 (TTY: 800-842-5357).

**አማርኛ (Amharic):**

ይህ ማስታወቂያ አስፈላጊ መረጃ ይዟል። ይህ ማስታወቂያ ስለ ማመልከቻዎ ወይም የ Premera Blue Cross ሽፋን አስፈላጊ መረጃ ሊኖረው ይችላል። በዚህ ማስታወቂያ ውስጥ ቁልፍ ቀዳሾች ሊኖሩ ይችላሉ። የጤና ሽፋንዎን ለመጠበቅና በአስፈላጊ እርዳታ ለማግኘት በተውሰኑ የጊዜ ገደቦች እርምጃ መውሰድ ይገባዎት ይሆናል። ይህን መረጃ እንዲያገኙ እና የለምንም ክፍያ በቋንቋዎ እርዳታ እንዲያገኙ መሰታወቅ አለዎት። በስልክ ቁጥር 800-722-1471 (TTY: 800-842-5357) ይደውሉ።

**العربية (Arabic):**

يحتوي هذا الإشعار على معلومات هامة. قد يحتوي هذا الإشعار على معلومات مهمة بخصوص طلبك أو التخطيط التي تزيد الحصول عليها من خلال Premera Blue Cross. قد تكون هناك تواريخ مهمة في هذا الإشعار. وقد تحتاج لاتخاذ إجراء في تاريخ معينة للحفاظ على تغطيتك الصحية أو المساعدة في دفع التكاليف. يحق لك الحصول على هذه المعلومات والمساعدة بلغتك دون تكبد أية تكلفة. اتصل بـ 800-722-1471 (TTY: 800-842-5357)

**中文 (Chinese):**

**本通知有重要的訊息。**本通知可能有關於您透過 Premera Blue Cross 提交的申請或保險的重要訊息。本通知內可能有重要日期。您可能需要在截止日期之前採取行動，以保留您的健康保險或者費用補貼。您有權利免費以您的母語得到本訊息和幫助。請撥電話 800-722-1471 (TTY: 800-842-5357)。

**Oromoo (Cushite):**

**Beeksisni kun odeeffannoo barbaachisaa qaba.** Beeksisti kun sagantaa yookan karaa Premera Blue Cross tiin tajaajila keessan ilaalchisee odeeffannoo barbaachisaa qabaachuu danda'a. Guyyaawwan murteessaa ta'an beeksisa kana keessatti ilaalaa. Tarii kaffaltiidhaan deeggaramuuf yookan tajaajila fayyaa keessaniif guyyaa dhumaa irratti wanti raawwattan jiraachuu danda'a. Kaffaltii irraa bilisa haala ta'een afaan keessaniin odeeffannoo argachuu fi deeggarsa argachuuf mirga ni qabaattu. Lakkoofsa bilbilaa 800-722-1471 (TTY: 800-842-5357) tii bilbilaa.

**Français (French):**

**Cet avis a d'importantes informations.** Cet avis peut avoir d'importantes informations sur votre demande ou la couverture par l'intermédiaire de Premera Blue Cross. Le présent avis peut contenir des dates clés. Vous devez peut-être prendre des mesures par certains délais pour maintenir votre couverture de santé ou d'aide avec les coûts. Vous avez le droit d'obtenir cette information et de l'aide dans votre langue à aucun coût. Appelez le 800-722-1471 (TTY: 800-842-5357).

**Kreyòl ayisyen (Creole):**

**Avi sila a gen Enfòmasyon Enpòtan ladann.** Avi sila a kapab genyen enfòmasyon enpòtan konsènan aplikasyon w lan oswa konsènan kouvèti asirans lan atravè Premera Blue Cross. Kapab genyen dat ki enpòtan nan avi sila a. Ou ka gen pou pran kèk aksyon avan sèten dat limit pou ka kenbe kouvèti asirans sante w la oswa pou yo ka ede w avèk depans yo. Se dwa w pou resewva enfòmasyon sa a ak asistans nan lang ou pale a, san ou pa gen pou peye pou sa. Rele nan 800-722-1471 (TTY: 800-842-5357).

**Deutsche (German):**

**Diese Benachrichtigung enthält wichtige Informationen.** Diese Benachrichtigung enthält unter Umständen wichtige Informationen bezüglich Ihres Antrags auf Krankenversicherungsschutz durch Premera Blue Cross. Suchen Sie nach eventuellen wichtigen Terminen in dieser Benachrichtigung. Sie könnten bis zu bestimmten Stichtagen handeln müssen, um Ihren Krankenversicherungsschutz oder Hilfe mit den Kosten zu behalten. Sie haben das Recht, kostenlose Hilfe und Informationen in Ihrer Sprache zu erhalten. Rufen Sie an unter 800-722-1471 (TTY: 800-842-5357).

**Hmoob (Hmong):**

**Tsab ntawv tshaj xo no muaj cov ntshiab lus tseem ceeb.** Tej zaum tsab ntawv tshaj xo no muaj cov ntshiab lus tseem ceeb txog koj daim ntawv thov kev pab los yog koj qhov kev pab cuam hnu ntawm Premera Blue Cross. Tej zaum muaj cov hnuv tseem ceeb uas sau rau hauv daim ntawv no. Tej zaum koj kuj yuav tau ua qee yam uas peb kom koj ua tsis pub dhau cov caij nyoog uas teev tseg rau hauv daim ntawv no mas koj thiaj yuav tau txais kev pab cuam kho mob los yog kev pab them tej nqi kho mob ntawd. Koj muaj cai kom lawv muab cov ntshiab lus no uas tau muab sau ua koj hom lus pub dawb rau koj. Hu rau 800-722-1471 (TTY: 800-842-5357).

**Iloko (Ilocano):**

**Daytoy a Pakdaar ket naglaon iti Napateg nga Impormasion.** Daytoy a pakdaar mabalin nga adda ket naglaon iti napateg nga impormasion maipanggep iti aplikasyonyo wenna coverage babaen iti Premera Blue Cross. Daytoy ket mabalin dagiti importante a petsa iti daytoy a pakdaar. Mabalin nga adda rumbeng nga aramidenyo nga addang sakbay dagiti partikular a naituding nga aldaw tapno mapagtalinaedyo ti coverage ti salun-atyto wenna tulong kadagiti gastos. Adda karbenganyo a mangala iti daytoy nga impormasion ken tulong iti bukodyo a pagsasao nga awan ti bayadanyo. Tumawag iti numero nga 800-722-1471 (TTY: 800-842-5357).

**Italiano (Italian):**

**Questo avviso contiene informazioni importanti.** Questo avviso può contenere informazioni importanti sulla tua domanda o copertura attraverso Premera Blue Cross. Potrebbero esserci date chiave in questo avviso. Potrebbe essere necessario un tuo intervento entro una scadenza determinata per consentirti di mantenere la tua copertura o sovvenzione. Hai il diritto di ottenere queste informazioni e assistenza nella tua lingua gratuitamente. Chiama 800-722-1471 (TTY: 800-842-5357).

**日本語 (Japanese):**

この通知には重要な情報が含まれています。この通知には、Premera Blue Cross の申請または補償範囲に関する重要な情報が含まれている場合があります。この通知に記載されている可能性がある重要な日付をご確認ください。健康保険や有料サポートを維持するには、特定の期日までに行動を取らなければならない場合があります。ご希望の言語による情報とサポートが無料で提供されます。800-722-1471 (TTY: 800-842-5357)までお電話ください。

**한국어 (Korean):**

본 통지서에는 중요한 정보가 들어 있습니다. 즉 이 통지서는 귀하의 신청에 관하여 그리고 Premera Blue Cross 를 통한 커버리지에 관한 정보를 포함하고 있을 수 있습니다. 본 통지서에는 핵심이 되는 날짜들이 있을 수 있습니다. 귀하의 건강 커버리지를 계속 유지하거나 비용을 절감하기 위해서 일정한 마감일까지 조치를 취해야 할 필요가 있을 수 있습니다. 귀하의 이러한 정보와 도움을 귀하의 언어로 비용 부담없이 얻을 수 있는 권리가 있습니다. 800-722-1471 (TTY: 800-842-5357) 로 전화하십시오.

**ລາວ (Lao):**

ແຈງການນີ້ມີຂໍ້ມູນສໍາຄັນ. ແຈງການນີ້ອາດຈະມີຂໍ້ມູນສໍາຄັນກ່ຽວກັບຄໍາຮ້ອງສະໝັກ ຫຼື ຄວາມຄົມຄອງປະກັນໄພຂອງທ່ານຜ່ານ Premera Blue Cross. ອາດຈະມີວັນທີ່ສໍາຄັນໃນແຈງການນີ້. ທ່ານອາດຈະຈໍາເປັນຕ້ອງດໍາເນີນການຕາມກຳນົດ ເວລາສະເພາະເພື່ອຮັກສາຄວາມຄົມຄອງປະກັນສະພາບ ຫຼື ຄວາມຊ່ວຍເຫຼືອເວັ້ນເວົ້ອງຄ່າໃຊ້ຈ່າຍຂອງທ່ານໄດ້. ທ່ານມີສິດໄດ້ຮັບຂໍ້ມູນນີ້ ແລະ ຄວາມຊ່ວຍເຫຼືອເປັນພາສາຂອງທ່ານໂດຍບໍ່ເສຍຄ່າ. ໃຫ້ໃບທາ 800-722-1471 (TTY: 800-842-5357).

**ភាសាខ្មែរ (Khmer):**

សេចក្តីជូនដំណឹងនេះមានព័ត៌មានយ៉ាងសំខាន់។ សេចក្តីជូនដំណឹងនេះប្រហែលជាមានព័ត៌មានយ៉ាងសំខាន់អំពីទម្រង់បែបបទ ឬការរៀបចំរបស់អ្នកកាមរយ: Premera Blue Cross ។ ប្រហែលជាមាន កាលបរិច្ឆេទសំខាន់នៅក្នុងសេចក្តីជូនដំណឹងនេះ។ អ្នកប្រហែលជាត្រូវការបញ្ជាក់សមត្ថភាព ដល់កិច្ចការច្នៃផ្ទះធានា ដើម្បីនឹងរក្សាទុកការធានារ៉ាប់រងអនាគតរបស់អ្នក ឬប្រាក់ដុល្លារចេញផ្លូវ។ អ្នកមានសិទ្ធិទទួលព័ត៌មាននេះ និងដុល្លារនៅក្នុងភាសារបស់អ្នកដោយមិនអស់លុយឡើយ។ សូមទូរស័ព្ទ 800-722-1471 (TTY: 800-842-5357)។

**ਪੰਜਾਬੀ (Punjabi):**

ਇਸ ਨੋਟਿਸ ਵਿਚ ਖਾਸ ਜਾਣਕਾਰੀ ਹੈ. ਇਸ ਨੋਟਿਸ ਵਿਚ Premera Blue Cross ਵਲੋਂ ਤੁਹਾਡੀ ਕਵਰੇਜ ਅਤੇ ਅਰਜੀ ਬਾਰੇ ਮਹੱਤਵਪੂਰਨ ਜਾਣਕਾਰੀ ਹੋ ਸਕਦੀ ਹੈ . ਇਸ ਨੋਟਿਸ ਨਵ ਖਾਸ ਤਾਰੀਖਾਂ ਹੋ ਸਕਦੀਆਂ ਹਨ. ਜੇਕਰ ਤੁਸੀਂ ਜਸਰਤ ਕਵਰੇਜ ਰਿੱਖਣੀ ਹੋਵੇ ਜਾਂ ਓਸ ਦੀ ਲਾਗਤ ਜਵਿੱਚ ਮਦਦ ਦੇ ਇਛੁੱਕ ਹੋ ਤਾਂ ਤੁਹਾਨੂੰ ਅੰਤਮ ਤਾਰੀਖ ਤੋਂ ਪਹਿਲਾਂ ਢੁੱਝ ਖਾਸ ਕਦਮ ਚੁੱਕਣ ਦੀ ਲੋੜ ਹੋ ਸਕਦੀ ਹੈ ,ਤੁਹਾਨੂੰ ਮੁਫਤ ਵਿੱਚ ਤੋਂ ਅਪਣੀ ਭਾਸ਼ਾ ਵਿੱਚ ਜਾਣਕਾਰੀ ਅਤੇ ਮਦਦ ਪ੍ਰਾਪਤ ਕਰਨ ਦਾ ਅਧਿਕਾਰ ਹੈ ,ਕਾਲ 800-722-1471 (TTY: 800-842-5357).

**فارسی (Farsi):**

این اعلامیه حاوی اطلاعات مهم میباشد. این اعلامیه ممکن است حاوی اطلاعات مهم درباره فرم تقاضا و یا پوشش بیمه ای شما از طریق Premera Blue Cross باشد. به تاریخ های مهم در این اعلامیه توجه نمایید. شما ممکن است برای حفظ پوشش بیمه تان یا کمک در پرداخت هزینه های درمانی تان، به تاریخ های مشخصی برای انجام کارهای خاصی احتیاج داشته باشید. شما حق این را دارید که این اطلاعات و کمک را به زبان خود به طور رایگان دریافت نمایید. برای کسب اطلاعات با شماره 800-722-1471 (کلیران TTY تماس باشماره 800-842-5357) تماس برقرار نمایید.

**Polskie (Polish):**

To ogłoszenie może zawierać ważne informacje. To ogłoszenie może zawierać ważne informacje odnośnie Państwa wniosku lub zakresu świadczeń poprzez Premera Blue Cross. Prosimy zwrócić uwagę na kluczowe daty, które mogą być zawarte w tym ogłoszeniu aby nie przekroczyć terminów w przypadku utrzymania polisy ubezpieczeniowej lub pomocy związanej z kosztami. Macie Państwo prawo do bezpłatnej informacji we własnym języku. Zadzwońcie pod 800-722-1471 (TTY: 800-842-5357).

**Português (Portuguese):**

Este aviso contém informações importantes. Este aviso poderá conter informações importantes a respeito de sua aplicação ou cobertura por meio do Premera Blue Cross. Poderão existir datas importantes neste aviso. Talvez seja necessário que você tome providências dentro de determinados prazos para manter sua cobertura de saúde ou ajuda de custos. Você tem o direito de obter esta informação e ajuda em seu idioma e sem custos. Ligue para 800-722-1471 (TTY: 800-842-5357).

**Română (Romanian):**

Prezenta notificare conține informații importante privind cererea sau acoperirea asigurării dumneavoastră de sănătate prin Premera Blue Cross. Pot exista date cheie în această notificare. Este posibil să fie nevoie să acționați până la anumite termene limită pentru a vă menține acoperirea asigurării de sănătate sau asistența provizorie la costuri. Aveți dreptul de a obține gratuit aceste informații și ajutor în limba dumneavoastră. Sunați la 800-722-1471 (TTY: 800-842-5357).

**Русский (Russian):**

Настоящее уведомление содержит важную информацию. Это уведомление может содержать важную информацию о вашем заявлении или страховом покрытии через Premera Blue Cross. В настоящем уведомлении могут быть указаны ключевые даты. Вам, возможно, потребуется принять меры к определенным предельным срокам для сохранения страхового покрытия или помощи с расходами. Вы имеете право на бесплатное получение этой информации и помощь на вашем языке. Звоните по телефону 800-722-1471 (TTY: 800-842-5357).

**Fa'asamoa (Samoan):**

Atonu ua iai i lenei fa'asilasilaga ni fa'amatalaga e sili ona taua e tatau ona e malamalama i ai. O lenei fa'asilasilaga o se fesoasoani e fa'amatala atili i ai i le tulaga o le polokalame, Premera Blue Cross, ua e tau fia maua atu i ai. Fa'amolemole, ia e iloilo fa'alelei i aso fa'apitoa olo'o iai i lenei fa'asilasilaga taua. Masalo o le'a iai ni feau e tatau ona e faia ao le'i aulia le aso ua ta'ua i lenei fa'asilasilaga ia e iai pea ma maua fesoasoani mai ai i le polokalame a le Malo olo'o e iai i ai. Olo'o iai iate oe le aia tatau e maua atu i lenei fa'asilasilaga ma lenei fa'matalaga i legagana e te malamalama i ai aunoa ma se togiga tupe. Vili atu i le telefoni 800-722-1471 (TTY: 800-842-5357).

**Español (Spanish):**

Este Aviso contiene información importante. Es posible que este aviso contenga información importante acerca de su solicitud o cobertura a través de Premera Blue Cross. Es posible que haya fechas clave en este aviso. Es posible que deba tomar alguna medida antes de determinadas fechas para mantener su cobertura médica o ayuda con los costos. Usted tiene derecho a recibir esta información y ayuda en su idioma sin costo alguno. Llame al 800-722-1471 (TTY: 800-842-5357).

**Tagalog (Tagalog):**

Ang Paunawa na ito ay naglalaman ng mahalagang impormasyon tungkol sa iyong aplikasyon o pagsakop sa pamamagitan ng Premera Blue Cross. Maaaring may mga mahalagang petsa dito sa paunawa. Maaring mangailangan ka na magsagawa ng hakbang sa ilang mga itinakdang panahon upang mapanatili ang iyong pagsakop sa kalusugan o tulong na walang gastos. May karapatan ka na makakuha ng ganiitong impormasyon at tulong sa iyong wika ng walang gastos. Tumawag sa 800-722-1471 (TTY: 800-842-5357).

**ไทย (Thai):**

ประกาศนี้มีข้อมูลสำคัญ ประกาศนี้อาจมีข้อมูลที่สำคัญเกี่ยวกับกาการสมัครหรือขอบเขตประกันสุขภาพของคุณผ่าน Premera Blue Cross และอาจมีกำหนดการในประกาศนี้ คุณอาจจะต้องดำเนินการภายในกำหนดระยะเวลาที่แน่นอนเพื่อจะรักษาการประกันสุขภาพของคุณหรือการช่วยเหลือที่มีค่าใช้จ่าย คุณมีสิทธิที่จะได้รับข้อมูลและความช่วยเหลือนี้ในภาษาของคุณโดยไม่มีค่าใช้จ่าย โทร 800-722-1471 (TTY: 800-842-5357)

**Український (Ukrainian):**

Це повідомлення містить важливу інформацію. Це повідомлення може містити важливу інформацію про Ваше звернення щодо страховального покриття через Premera Blue Cross. Зверніть увагу на ключові дати, які можуть бути вказані у цьому повідомленні. Існує імовірність того, що Вам треба буде здійснити певні кроки у конкретні кінцеві строки для того, щоб зберегти Ваше медичне страхування або отримати фінансову допомогу. У Вас є право на отримання цієї інформації та допомоги безкоштовно на Вашій рідній мові. Дзвоніть за номером телефону 800-722-1471 (TTY: 800-842-5357).

**Tiếng Việt (Vietnamese):**

Thông báo này cung cấp thông tin quan trọng. Thông báo này có thông tin quan trọng về đơn xin tham gia hoặc hợp đồng bảo hiểm của quý vị qua chương trình Premera Blue Cross. Xin xem ngày quan trọng trong thông báo này. Quý vị có thể phải thực hiện theo thông báo đúng trong thời hạn để duy trì bảo hiểm sức khỏe hoặc được trợ giúp thêm về chi phí. Quý vị có quyền được biết thông tin này và được trợ giúp bằng ngôn ngữ của mình miễn phí. Xin gọi số 800-722-1471 (TTY: 800-842-5357).