

MEDICAL POLICY – 7.01.519

Treatment of Varicose Veins/Venous Insufficiency

BCBSA Ref. Policy: 7.01.124

Effective Date: **Feb. 5, 2021**

Last Revised: Oct. 13, 2020


Replaces: 7.01.55, 7.01.76,
7.01.515, and
7.01.124

RELATED MEDICAL POLICIES:

10.01.514 Cosmetic and Reconstructive Services

Select a hyperlink below to be directed to that section.

[POLICY CRITERIA](#) | [DOCUMENTATION REQUIREMENTS](#) | [CODING](#)
[RELATED INFORMATION](#) | [EVIDENCE REVIEW](#) | [REFERENCES](#) | [HISTORY](#)

 Clicking this icon returns you to the hyperlinks menu above.

Introduction

Varicose veins are swollen, twisted veins that are visible just below the skin. They most often occur in the legs but can develop in other parts of the body. Veins have valves that keep the blood moving toward the heart. If the valves are weak or damaged, blood can pool in the veins. It's this pooling that causes the veins to swell and appear twisted. Most varicose veins do not result in physical symptoms. On occasion, they can cause symptoms such as pain, an achy feeling, bleeding, or skin ulcers (sores). Varicose veins usually can be treated without surgery by activities such as exercising, raising the legs, or wearing compression stockings. This policy describes when varicose vein surgery or other procedures may be considered medically necessary.

Note: The Introduction section is for your general knowledge and is not to be taken as policy coverage criteria. The rest of the policy uses specific words and concepts familiar to medical professionals. It is intended for providers. A provider can be a person, such as a doctor, nurse, psychologist, or dentist. A provider also can be a place where medical care is given, like a hospital, clinic, or lab. This policy informs them about when a service may be covered.

Policy Coverage Criteria

Procedures	Medical Necessity
<p>Vein ablation procedures:</p> <ul style="list-style-type: none"> • Surgery (ligation and stripping) • Endovenous thermal ablation (radiofrequency or laser) • Chemical ablation (microfoam sclerotherapy Varithena) • Cyanoacrylate adhesive (VenaSeal™ Closure System) <p>(36465, 36466, 36475, 36476, 36478, 36479, 36482, 36483, 37700, 37718, 37722, 37735, 37780, 37785)</p>	<p>Surgery (ligation and stripping), endovenous thermal ablation (radiofrequency or laser), chemical ablation (microfoam sclerotherapy, eg, Varithena), or cyanoacrylate adhesive (eg, VenaSeal) may be considered medically necessary for treatment of symptomatic varicose veins (<u>great or small saphenous veins</u>)/venous insufficiency when the following criteria have been met:</p> <ul style="list-style-type: none"> • There is moderate to severe (greater than 0.5 second) saphenous reflux documented on venous studies and CEAP [Clinical-Etiology-Anatomy-Pathophysiology] class C2 or greater (see Definition of Terms) <p>AND</p> <ul style="list-style-type: none"> • There is documentation of ONE or more of the following four indications: <ul style="list-style-type: none"> ○ Ulceration secondary to venous stasis, or ○ Recurrent superficial thrombophlebitis, or ○ Hemorrhage or recurrent bleeding episodes from a ruptured superficial varicosity, or ○ Persistent pain, swelling, itching, burning, or other symptoms are associated with saphenous reflux along with both of the following: <ul style="list-style-type: none"> ▪ These symptoms significantly interfere with activities of daily living, and ▪ Conservative management, including compression therapy (see Definition of Terms) for at least 3 months, has not improved the symptoms <p>Note: If compression therapy is successful, it should be continued for as long as it is working. Failure of conservative therapy is defined as pain, swelling, itching, burning, or other symptoms associated with vein reflux, despite conservative therapy, severe enough to require daily pain medicines and causing inability to manage daily activities at home or at work.</p>



Procedures	Medical Necessity
	<p>Treatment of <u>great or small saphenous veins</u> by surgery, endovenous thermal ablation (radiofrequency or laser), or chemical ablation (microfoam sclerotherapy, eg, Varithena), or cyanoacrylate adhesive (eg, VenaSeal) that does not meet the criteria described above is considered not medically necessary.</p> <p>Surgery (ligation and stripping), endovenous thermal ablation (radiofrequency or laser), chemical ablation (microfoam sclerotherapy (eg, Varithena), or cyanoacrylate adhesive (VenaSeal) may be considered medically necessary for treatment of symptomatic <u>accessory saphenous veins</u> /venous insufficiency when ALL of the following criteria have been met:</p> <ul style="list-style-type: none"> • Incompetence of the accessory saphenous vein is isolated <p>AND</p> <ul style="list-style-type: none"> • There is moderate to severe (greater than 0.5 second) accessory saphenous reflux documented on venous studies <p>AND</p> <ul style="list-style-type: none"> • There is documentation of ONE or more of the following four indications: <ul style="list-style-type: none"> ○ Ulceration secondary to venous stasis, or ○ Recurrent superficial thrombophlebitis, or ○ Hemorrhage or recurrent bleeding episodes from a ruptured superficial varicosity, or ○ Persistent pain, swelling, itching, burning, or other symptoms are associated with saphenous reflux along with both of the following: <ul style="list-style-type: none"> ▪ These symptoms significantly interfere with activities of daily living, and ▪ Conservative management, including compression therapy (see Definition of Terms) for at least 3 months, has not improved these symptoms. <p>Note: If compression therapy is successful, it should be continued for as long as it is working. Failure of conservative therapy is defined as pain, swelling, itching, burning, or other symptoms associated with vein reflux, despite conservative therapy, severe enough to require daily pain</p>



Procedures	Medical Necessity
	<p>medicines and causing inability to manage daily activities at home or at work</p> <p>Concurrent treatment of the accessory saphenous veins along with the great or small saphenous veins may be considered medically necessary when criteria are met for each vein and there is documentation of anatomy showing that the accessory saphenous vein discharged directly into the common femoral vein.</p> <p>Treatment of <u>accessory saphenous veins</u> by surgery, endovenous thermal ablation (radiofrequency or laser), chemical ablation (microfoam sclerotherapy (eg, Varithena), or cyanoacrylate adhesive that does not meet the criteria described above is considered not medically necessary.</p> <p>Endovenous thermal ablation (radiofrequency or laser) of <u>tributary veins</u> is considered investigational</p>
<p>Vein ablation procedures:</p> <ul style="list-style-type: none"> • Stab avulsion • Hook phlebectomy • Sclerotherapy • TIPP* <p>(36470, 36471, 37765, 37766, 37799)</p> <p>Note: *TIPP-Transilluminated powered phlebectomy</p>	<p>Phlebectomy (stab avulsion, hook phlebectomy, or TIPP) or initial* sclerotherapy as a component of the treatment of <u>symptomatic varicose tributaries</u> when performed either at the same time or following prior treatment (surgical, radiofrequency, or laser) of the saphenous veins is considered medically necessary.</p> <p>Sclerotherapy of <u>isolated tributary veins</u> without prior or concurrent treatment of saphenous veins is considered investigational.</p> <p>Note: *Initial sclerotherapy for these indications is limited to a maximum of 3 sclerotherapy treatment sessions per leg.</p> <p>For sclerotherapy treatment requests after ablation (surgical, radiofrequency, or laser) of the <u>saphenous veins</u>, when performed for the treatment of residual or recurrent symptoms, there must be 3 months of conservative management, including compression therapy (see Definition</p>



Procedures	Medical Necessity
	<p>of Terms) after the most recent varicose vein procedure, which has not successfully treated the patient’s symptoms.</p> <p>Note: If compression therapy is successful, it should be continued for as long as it is working. Failure of conservative therapy is defined as pain, swelling, itching, burning, or other symptoms associated with vein reflux, despite conservative therapy, severe enough to require daily pain medicines and causing inability to manage daily activities at home or at work.</p> <p>Any other techniques, other than stab avulsion, hook phlebectomy, sclerotherapy (liquid or foam), or TIPP, for the treatment of <u>symptomatic varicose tributaries</u>, when performed either at the same time or following prior treatment of saphenous veins, is considered investigational.</p> <p>Sclerotherapy techniques, other than microfoam sclerotherapy (eg, Varithena), of <u>great, small, or accessory saphenous veins</u> is considered investigational.</p> <p>Sclerotherapy of <u>perforator veins</u> is considered investigational.</p> <p>Stab avulsion, hook phlebectomy, or TIPP of <u>perforator, great or small saphenous, or accessory saphenous veins</u> is considered investigational.</p>
Ultrasound guidance	<p>It should be noted that the bulk of the literature discussing the role of ultrasound guidance refers to sclerotherapy of the saphenous vein, as opposed to the varicose tributaries. When ultrasound guidance is used to guide sclerotherapy of the varicose tributaries, it would be considered either not medically necessary or incidental to the injection procedure.</p>
<p>Vein ablation procedures:</p> <ul style="list-style-type: none"> • Surgery (ligation)/SEPS • Thermal ablation (radiofrequency, laser) <p>(37500, 37760, 37761)</p>	<p>Surgical ligation (including subfascial endoscopic perforator surgery [SEPS] or endovenous thermal ablation (radiofrequency or laser) of incompetent <u>perforator veins</u> may be considered medically necessary as a treatment of leg ulcers associated with chronic venous insufficiency when ALL of the following criteria have been met:</p>



Procedures	Medical Necessity
	<ul style="list-style-type: none"> • There is documented perforator reflux (greater than 0.5 second) on venous studies; And • The superficial saphenous veins (great, small, accessory saphenous and symptomatic varicose tributaries) have been previously eliminated; And • Ulcers have not resolved following combined superficial vein treatment and compression therapy (see Definition of Terms) for at least 3 months; And • The venous insufficiency is not secondary to deep venous thromboembolism <p>Note: If compression therapy is successful, it should be continued for as long as it is working. Failure of conservative therapy is defined as pain, swelling, itching, burning, or other symptoms associated with vein reflux, despite conservative therapy, severe enough to require daily pain medicines and causing inability to manage daily activities at home or at work</p> <p>Ligation or ablation of incompetent perforator veins performed concurrently with superficial venous surgery is not medically necessary.</p>
Telangiectasia (36468)	Treatment, by any method, of small telangiectasia such as spider veins (1mm or less), superficial reticular veins (1-2 mm), angiomas, and hemangiomas is considered cosmetic.

Treatment	Investigational
<p>Other techniques</p> <p>(0524T, 36473, 36474, 37799)</p>	<p>The following techniques are considered investigational for the treatment of any vein(s):</p> <ul style="list-style-type: none"> • Endovenous cryoablation • Mechanochemical ablation (eg, MOCA, ClariVein™ Catheter) <p>Note: This list may not be all inclusive.</p>

Documentation Requirements

Review of this procedure requires the provider to specifically indicate the following in the clinical notes submitted with the request:



Documentation Requirements

- Specific procedures to be performed, including the CPT code (s)

AND

- Specific vein to be treated for each CPT code and whether the proposed treatment is to the left leg, right leg, or both legs

Additionally, written documentation in the medical record must include:

1. CPT codes for the procedures being requested, and the name of the vein to be treated with each CPT code
2. Copy of the venous studies (ultrasound studies evaluating blood flow in the veins)
3. CEAP (Clinical-Etiology-Anatomy-Pathophysiology) class C2 or greater (Clinical classification, large varicose veins)
4. One or more indications:
 - Ulceration secondary to venous stasis (leg ulcers caused by poor blood flow in the veins)
 - Recurrent superficial thrombophlebitis (recurrent blood clots or inflammation in a small vein near the surface of the skin)
 - Hemorrhage or recurrent bleeding episodes from a ruptured superficial varicosity (bleeding from a varicose vein)
 - Persistent pain, swelling, itching, burning, or other symptoms are associated with saphenous reflux (pooling of the blood caused by valves that do not work correctly)

AND

- These symptoms significantly interfere with activities of daily living; and
- Conservative management including compression therapy for at least 3 months has not improved the symptoms

Coding

Code	Description
CPT	
0524T	Endovenous catheter directed chemical ablation with balloon isolation of incompetent extremity vein, open or percutaneous, including all vascular access, catheter manipulation, diagnostic imaging, imaging guidance and monitoring
36465	Injection of non-compounded foam sclerosant with ultrasound compression maneuvers to guide dispersion of the injectate, inclusive of all imaging guidance and monitoring; single incompetent extremity truncal vein (eg, great saphenous vein, accessory saphenous vein) (Varithena)



Code	Description
36466	Injection of non-compounded foam sclerosant with ultrasound compression maneuvers to guide dispersion of the injectate, inclusive of all imaging guidance and monitoring; multiple incompetent truncal veins (eg, great saphenous vein, accessory saphenous vein), same leg (Varithena)
36468	Single or multiple injections of sclerosing solutions, spider veins (telangiectasia); limb or trunk
36470	Injection of sclerosing solution; single vein
36471	Injection of sclerosing solution; multiple veins, same leg
36473	Endovenous ablation therapy of incompetent vein, extremity, inclusive of all imaging guidance and monitoring, percutaneous, mechanochemical; first vein treated (eg, MOCA ClariVein)
36474	Endovenous ablation therapy of incompetent vein, extremity, inclusive of all imaging guidance and monitoring, percutaneous, mechanochemical; subsequent vein(s) treated in a single extremity, each through separate access sites (List separately in addition to code for primary procedure) (eg, MOCA ClariVein)
36475	Endovenous ablation therapy of incompetent vein, extremity, inclusive of all imaging guidance and monitoring, percutaneous, radiofrequency; first vein treated
36476	Endovenous ablation therapy of incompetent vein, extremity, inclusive of all imaging guidance and monitoring, percutaneous, radiofrequency; second and subsequent veins treated in a single extremity, each through separate access sites (List separately in addition to code for primary procedure)
36478	Endovenous ablation therapy of incompetent vein, extremity, inclusive of all imaging guidance and monitoring, percutaneous, laser; first vein treated
36479	Endovenous ablation therapy of incompetent vein, extremity, inclusive of all imaging guidance and monitoring, percutaneous, laser; second and subsequent veins treated in a single extremity, each through separate access sites (List separately in addition to code for primary procedure)
36482	Endovenous ablation therapy of incompetent vein, extremity, by transcatheter delivery of a chemical adhesive (eg, cyanoacrylate) remote from the access site, inclusive of all imaging guidance and monitoring, percutaneous; first vein treated (VenaSeal)
36483	Endovenous ablation therapy of incompetent vein, extremity, by transcatheter delivery of a chemical adhesive (eg, cyanoacrylate) remote from the access site, inclusive of all imaging guidance and monitoring, percutaneous; subsequent vein(s) treated in a single extremity, each through separate access sites (List separately in addition to code for primary procedure) (VenaSeal)
37500	Vascular endoscopy, surgical, with ligation of perforator veins, subfascial (SEPS)
37700	Ligation and division of long saphenous vein at saphenofemoral junction, or distal interruptions
37718	Ligation, division, and stripping, short saphenous vein
37722	Ligation, division, and stripping, long (greater) saphenous veins from saphenofemoral junction to knee or below



Code	Description
37735	Ligation and division and complete stripping of long and short saphenous veins with radical excision of ulcer and skin graft and/or interruption of communicating veins of lower leg, with excision of deep fascia
37760	Ligation of perforator veins, subfascial, radical (Linton type) including skin graft, when performed, open, 1 leg
37761	Ligation of perforator vein(s), subfascial, open, including ultrasound guidance, when performed, 1 leg
37765	Stab phlebectomy of varicose veins, 1 extremity; 10-20 stab incisions
37766	Stab phlebectomy of varicose veins, 1 extremity; more than 20 stab incisions
37780	Ligation and division of short saphenous vein at saphenopopliteal junction (separate procedure)
37785	Ligation, division, and/or excision of varicose vein cluster(s), one leg
37799	Unlisted procedure, vascular surgery (TRIVEX System)

Note: CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). HCPCS codes, descriptions and materials are copyrighted by Centers for Medicare Services (CMS).

Related Information

Definition of Terms

Accessory saphenous veins: Veins that travel in parallel with the greater and lesser saphenous veins and are generally 2 to 2.5 mm in their normal state.

Clinical-Etiologic-Anatomic-Pathophysiology (CEAP) classification system: The standard classification of venous disease considers the clinical, etiologic, anatomic, and pathologic (CEAP) characteristics of venous insufficiency. See table below.

Class	Clinical Classification
C0	No visible or palpable signs of venous disease
C1	Telangiectasies or reticular veins
C2	Varicose veins
C3	Edema
C4a	Pigmentation and eczema
C4b	Lipodermatosclerosis and atrophie blanche
C5	Healed venous ulcer
C6	Active venous ulcer



Class	Clinical Classification
S	Symptoms including ache, pain, tightness, skin irritation, heaviness, muscle cramps, and other complaints attributable to venous dysfunction
A	Asymptomatic

Adapted from: <https://www.veinforum.org/wp-content/uploads/2018/03/Revised-CEAP-Classification-May-2004.pdf> Accessed July 2019.

Compression Therapy: Compression hosiery or stockings are often the first line of treatment for varicose veins. Compression stockings are generally thought to be effective in a range of 20 mmHg to 40mmHg. The Society for Vascular Surgery/American Venous Forum recommends compression therapy of 20-30 mmHg for patients with symptomatic varicose veins. Compression stockings of 20-30mmHg are available over the counter without a prescription.

Greater/long saphenous vein: Superficial vein running the entire length of the leg and is generally 3 mm in its normal state. A typical GSV contains an average of 7 valves throughout its entire length, and it is the most common superficial vein to develop venous reflux.

Lesser (small)/short saphenous veins: Superficial vein of the calf and is generally 2.5 mm in its normal state. The small saphenous vein originates at the back of the ankle near the outer malleous bone, and usually runs up the back of the lower leg to the popliteal vein behind the knee.

Moderate to severe reflux: In current practice, most vascular laboratories consider the presence of venous flow reversal for greater than 0.5 second with proximal compression, Valsalva maneuver, or distal compression and release to represent pathologic reflux.

Perforator veins: These connect superficial veins to deep veins. They contain one-way valves to direct the blood from the superficial system to the deep system and are generally less than 3mm in their normal state.

Stab avulsion: This technique is also known as stab/hook phlebectomy. Stab avulsion results in removal of the varicose veins through incisions that are 2-3 mm in length. The veins are hooked with a tiny hook-like instrument and pulled out. The wounds are closed with tapes, not sutures, and the leg is wrapped in elastic compression support. Once healed, the incision sites are almost invisible.

Telangiectasia/spider veins: Very small (≤ 1 mm in diameter) thread veins found commonly just under the surface of the skin, usually not distorting skin or surrounding tissues. These are not considered varicose veins.



Tributary veins: Veins that empty into a larger vein.

Evidence Review

Description

A variety of treatment modalities are available to treat varicose veins/venous insufficiency, including surgery, thermal ablation, sclerotherapy, mechanochemical ablation (MOCA), cyanoacrylate adhesive (CAC), and cryotherapy. The application of each modality is influenced by the severity of the symptoms, type of vein, source of venous reflux, and the use of other (prior or concurrent) treatments.

Background

Venous Reflux/Venous Insufficiency

The venous system of the lower extremities consists of the superficial veins (this includes the great and small saphenous and accessory, or duplicate, veins that travel in parallel with the great and small saphenous veins), the deep system (popliteal and femoral veins), and perforator veins that cross through the fascia and connect the deep and superficial systems. One-way valves are present within all veins to direct the return of blood up the lower limb. Because the venous pressure in the deep system is generally greater than that of the superficial system, valve incompetence at any level may lead to backflow (venous reflux) with pooling of blood in superficial veins. Varicose veins with visible varicosities may be the only sign of venous reflux, although itching, heaviness, tension, and pain may also occur. Chronic venous insufficiency secondary to venous reflux can lead to thrombophlebitis, leg ulcerations, and hemorrhage. The CEAP classification of venous disease considers the clinical, etiologic, anatomic, and pathologic characteristics of venous insufficiency, ranging from class 0 (no visible sign of disease) to class 6 (active ulceration).

Treatment

Treatment of venous reflux/venous insufficiency seeks to reduce abnormal pressure transmission from the deep to the superficial veins. Conservative medical treatment consists of elevation of



the extremities, graded compression, and wound care when indicated. Conventional surgical treatment consists of identifying and correcting the site of reflux by ligation of the incompetent junction followed by stripping of the vein to redirect venous flow through veins with intact valves. While most venous reflux is secondary to incompetent valves at the saphenofemoral or saphenopopliteal junctions, reflux may also occur at incompetent valves in the perforator veins or the deep venous system. The competence of any single valve is not static and may be pressure-dependent. For example, accessory saphenous veins may have independent saphenofemoral or saphenopopliteal junctions that become incompetent when the great or small saphenous veins are eliminated, and blood flow is diverted through the accessory veins.

Treatment of Saphenous Veins and Tributaries

Saphenous veins include the great and small saphenous and accessory saphenous veins that travel in parallel with the great or small saphenous veins. Tributaries are veins that empty into a larger vein. Treatment of venous reflux typically includes the following:

- Identification by preoperative Doppler ultrasonography of the valvular incompetence
- Control of the most proximal point of reflux, traditionally by suture ligation of the incompetent saphenofemoral or saphenopopliteal junction
- Removal of the superficial vein from circulation, eg, by stripping of the great and/or small saphenous veins
- Removal of varicose tributaries (at the time of the initial treatment or subsequently) by stab avulsion (phlebectomy) or injection sclerotherapy.

Minimally invasive alternatives to ligation and stripping have been investigated. They include forms of sclerotherapy, cyanoacrylate adhesive, , and thermal ablation using cryotherapy, high frequency radio waves (200–300 kHz), or laser energy.

Thermal Ablation

Radiofrequency ablation (RFA) is performed using a specially designed catheter inserted through a small incision in the distal medial thigh to within 1 to 2 cm of the saphenofemoral junction. The catheter is slowly withdrawn, closing the vein. Laser ablation is performed similarly; a laser fiber is introduced into the great saphenous vein under ultrasound guidance; the laser is activated and slowly removed along the course of the saphenous vein. Cryoablation uses



extreme cold. The objective of endovenous techniques is to injure the vessel, causing retraction and subsequent fibrotic occlusion of the vein. Technical developments since thermal ablation procedures were initially introduced include the use of perivenous tumescent anesthesia, which allows successful treatment of veins larger than 12 mm in diameter and helps to protect adjacent tissue from thermal damage during treatment of the small saphenous vein.

Sclerotherapy

The objective of sclerotherapy is to destroy the endothelium of the target vessel by injecting an irritant solution (either a detergent, osmotic solution, or chemical irritant), ultimately occluding the vessel. Treatment success depends on accurate injection of the vessel, an adequate injectate volume and concentration of sclerosant, and compression. Historically, larger veins and very tortuous veins were not considered good candidates for sclerotherapy due to technical limitations. Technical improvements in sclerotherapy have included the routine use of Duplex ultrasound to target refluxing vessels, luminal compression of the vein with anesthetics, and a foam/sclerosant injectate in place of liquid sclerosant. Foam sclerosants are produced by forcibly mixing a gas (eg, air or carbon dioxide) with a liquid sclerosant (eg, polidocanol or sodium tetradecyl sulfate). Physician-compounded foam is produced at the time of treatment. A commercially available microfoam sclerosant with a proprietary gas mix is available that is proposed to provide smaller and more consistent bubble size than what is produced with physician-compounded sclerosant foam.

Endovenous Mechanochemical Ablation (MCA)

Endovenous mechanochemical ablation uses both sclerotherapy and mechanical damage to the lumen. Following ultrasound imaging, a disposable catheter with a motor drive is inserted into the distal end of the target vein and advanced to the saphenofemoral junction. As the catheter is pulled back, a wire rotates at 3500 rpm within the lumen of the vein, abrading the lumen. At the same time, a liquid sclerosant (sodium tetradecyl sulfate) is infused near the rotating wire. It is proposed that mechanical ablation allows for better efficacy of the sclerosant, and results in less pain and risk of nerve injury without the need for the tumescent anesthesia used with endovenous thermal ablation techniques (radiofrequency ablation and endovenous laser ablation).



Cyanoacrylate Adhesive

A cyanoacrylate adhesive is a clear, free-flowing liquid that polymerizes in the vessel via an anionic mechanism (ie, polymerizes into a solid material on contact with body fluids or tissue). The adhesive is gradually injected along the length of the vein in conjunction with ultrasound and manual compression. The acute coaptation halts blood flow through the vein until the implanted adhesive becomes fibrotically encapsulated and establishes chronic occlusion of the treated vein. Cyanoacrylate glue has been used as a surgical adhesive and sealant for a variety of indications, including gastrointestinal bleeding, embolization of brain arteriovenous malformations, and to seal surgical incisions or other skin wounds.

Transilluminated Powered Phlebectomy (TIPP)

Transilluminated powered phlebectomy (TIPP) is an alternative to stab avulsion or hook phlebectomy. This procedure uses two instruments: an illuminator, which also provides irrigation, and a resector, which has an oscillating tip and suction pump. Following removal of the saphenous vein, the illuminator is introduced via a small incision in the skin and tumescence solution (anesthetic and epinephrine) is infiltrated along the course of the varicosity. The resector is then inserted under the skin from the opposite direction, and the oscillating tip is placed directly beneath the illuminated veins to fragment and loosen the veins from the supporting tissue. Irrigation from the illuminator is used to clear the vein fragments and blood through aspiration and additional drainage holes. The illuminator and resector tips may then be repositioned, thereby reducing the number of incisions needed when compared with stab avulsion or hook phlebectomy. It has been proposed that transilluminated powered phlebectomy might decrease surgical time, decrease complications such as bruising, and lead to a faster recovery than established procedures.

Treatment of Perforator Veins

Perforator veins cross through the fascia and connect the deep and superficial venous systems. Incompetent perforating veins were originally treated with an open surgical procedure, called the Linton procedure, which involved a long medial calf incision to expose all posterior, medial, and paramedial perforators. While this procedure was associated with healing of ulcers, it was largely abandoned due to a high incidence of wound complications. The Linton procedure was subsequently modified by using a series of perpendicular skin flaps instead of a longitudinal skin flap to provide access to incompetent perforator veins in the lower part of the leg. The modified



Linton procedure may occasionally be used to close incompetent perforator veins that cannot be reached by less invasive procedures.

Subfascial endoscopic perforator surgery is a less-invasive surgical procedure for the treatment of incompetent perforators and has been reported since the mid-1980s. Guided by Duplex ultrasound scanning, small incisions are made in the skin, and the perforating veins are clipped or divided by endoscopic scissors. The surgery can be performed as an outpatient procedure. Endovenous ablation of incompetent perforator veins with sclerotherapy and radiofrequency ablation has also been reported.

Summary of Evidence

Saphenous Veins

For individuals who have varicose veins/venous insufficiency and saphenous vein reflux who receive endovenous thermal ablation (radiofrequency or laser), the evidence includes randomized controlled trials (RCTs) and systematic reviews of controlled trials. The relevant outcomes are symptoms, change in disease status, morbid events, quality of life (QOL), and treatment-related morbidity (TRM). There are a number of large RCTs and systematic reviews of RCTs assessing endovenous thermal ablation of the saphenous veins. Comparison with the standard of ligation and stripping at 2- to 5-year follow-up has supported the use of both endovenous laser ablation and radiofrequency ablation (RFA). Evidence has suggested that ligation and stripping lead to more neovascularization, while thermal ablation leads to more recanalization, resulting in similar clinical outcomes for endovenous thermal ablation and surgery. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have varicose veins/venous insufficiency and saphenous vein reflux who receive microfoam sclerotherapy, the evidence includes RCTs. The relevant outcomes are symptoms, change in disease status, morbid events, quality of life, and treatment-related morbidity. For physician-compounded sclerotherapy, there is high variability in success rates and some reports of serious adverse events. By comparison, rates of occlusion with the microfoam sclerotherapy (polidocanol 1%) approved by the Food and Drug Administration are similar to those reported for endovenous laser ablation or stripping. Results of a noninferiority trial of physician-compounded sclerotherapy have indicated that once occluded, recurrence rates at two years are similar to those of ligation and stripping. Together, this evidence indicates that the more consistent occlusion with the microfoam sclerotherapy preparation will lead to recurrence



rates similar to ligation and stripping in the longer term. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

Based on the available evidence, clinical input obtained in 2015, and clinical practice guidelines, the use of endovenous RFA, endovenous laser ablation, and microfoam sclerotherapy are considered to improve outcomes when used in the saphenous veins. For treatment of saphenous tributaries at the same time or following treatment of the saphenous vein, stab avulsion, hook phlebectomy, sclerotherapy, or transilluminated powered phlebectomy improve outcomes.

For individuals who have varicose veins/venous insufficiency and saphenous vein reflux who receive mechanochemical ablation, the evidence includes four RCTs with 6 months to two years results that compared MOCA to thermal ablation, a prospective cohort with follow-up out to 5 years, and retrospective case series. The relevant outcomes are symptoms, change in disease status, morbid events, quality of life, and treatment-related morbidity. Mechanochemical ablation is a combination of liquid sclerotherapy with mechanical abrasion. A potential advantage of this procedure compared with thermal ablation is that mechanochemical ablation does not require multiple needle sticks with tumescent anesthesia and may result in less pain during the procedure. Results to date have been mixed regarding a reduction in intraprocedural pain compared to thermal ablation procedures. Occlusion rates at 6 mo to 2 year from RCTs indicate lower anatomic success rates compared to thermal ablation, but a difference in clinical outcomes at these early time points has not been observed. Experience with other endoluminal ablation procedures suggests that lower anatomic success in the short term is associated with recanalization and clinical recurrence between 2 to 5 years. The possibility of later clinical recurrence is supported by a prospective cohort study with 5-year follow-up following treatment with MOCA. However, there have been improvements in technique since the cohort study was begun, and clinical progression is frequently observed with venous disease. Because of these limitations of the single arm studies, longer follow-up in the more recently conducted RCTs is needed to establish the efficacy and durability of this procedure compared with the criterion standard of thermal ablation. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have varicose veins/venous insufficiency and saphenous vein reflux who receive cyanoacrylate adhesive (CAC), the evidence includes two RCTs and a prospective cohort. The relevant outcomes are symptoms, change in disease status, morbid events, quality of life, and treatment-related morbidity. Evidence includes a multicenter noninferiority trial with follow-up through 36 months, an RCT with follow-up through 24 months, and a prospective cohort with 30-month follow-up. The short-term efficacy of VenaSeal CAC has been shown to be noninferior to RFA at up to 36 months. At 24 and 36 months the study had greater than 20%



loss to follow-up, but loss to follow-up was similar in the 2 groups at the long-term follow-up and is not expected to influence the comparative results. A second RCT (n=525) with the same active CAC ingredient (N-butyl cyanoacrylate) that is currently available outside of the U.S. found no significant differences in vein closure between CAC and thermal ablation controls at 24-month follow-up. The CAC procedure and return to work were shorter and pain scores were lower compared to thermal ablation, although the subjective pain scores may have been influenced by differing expectations in this study. A prospective cohort reported high closure rates at 30 months. Overall, results indicate that outcomes from CAC are at least as good as thermal ablation techniques, the current standard of care. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have varicose veins/venous insufficiency and saphenous vein reflux who receive cryoablation, the evidence includes RCTs and multicenter series. The relevant outcomes are symptoms, change in disease status, morbid events, quality of life, and treatment-related morbidity. Results from a recent RCT of cryoablation have indicated that this therapy is inferior to conventional stripping. Studies showing a benefit on health outcomes are needed. The evidence is insufficient to determine the effects of the technology on health outcomes.

Varicose Tributary Veins

For individuals who have varicose tributary veins who receive ablation (stab avulsion, sclerotherapy, or phlebectomy) of tributary veins, the evidence includes RCTs and systematic reviews of RCTs. The relevant outcomes are symptoms, change in disease status, morbid events, quality of life, and treatment-related morbidity. The literature has shown that sclerotherapy is effective for treating tributary veins following occlusion of the saphenofemoral or saphenopopliteal junction and saphenous veins. No studies have been identified comparing RFA or laser ablation of tributary veins with standard procedures (microphlebectomy and/or sclerotherapy). Transilluminated powered phlebectomy is effective at removing varicosities; outcomes are comparable to available alternatives such as stab avulsion and hook phlebectomy. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

Perforator Veins

For individuals who have perforator vein reflux who receive ablation (eg, subfascial endoscopic perforator surgery) of perforator veins, the evidence includes RCTs and systematic reviews of



RCTs. The relevant outcomes are symptoms, change in disease status, morbid events, quality of life, and treatment-related morbidity. The literature has indicated that the routine ligation or ablation of incompetent perforator veins is not necessary for the treatment of varicose veins/venous insufficiency at the time of superficial vein procedures. However, when combined superficial vein procedures and compression therapy have failed to improve symptoms (ie, ulcers), treatment of perforator vein reflux may be as beneficial as an alternative (eg, deep vein valve replacement). Comparative studies are needed to determine the most effective method of ligating or ablating incompetent perforator veins. Subfascial endoscopic perforator surgery is possibly as effective as the Linton procedure with a reduction in adverse events. Although only one case series has been identified showing an improvement in health outcomes, endovenous ablation with specialized laser or radiofrequency probes has been shown to effectively ablate incompetent perforator veins with a potential decrease in morbidity compared with surgical interventions. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

Ongoing and Unpublished Clinical Trials

Some currently ongoing and unpublished trials that might influence this policy are listed in [Table 1](#).

Table 1. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
Ongoing			
NCT03392753	Mechanochemical Ablation Compared to Cyanoacrylate Adhesive	180	Dec 2020
NCT037222134	Mechanochemical Ablation vs Thermal Ablation in Patients with Great Saphenous Vein Insufficiency	132	Dec 2020
NTR4613^a	Mechanochemical endovenous ablation versus radiofrequency ablation in the treatment of primary small saphenous vein insufficiency (MESSI trial)	160	Apr 2020
NCT01936168	Mechanochemical endovenous ablation versus radiofrequency ablation in the treatment of primary great saphenous vein incompetence (MARADONA)	213	Dec 2020
NCT03835559	Randomized Controlled Trial Comparing the Clinical Outcomes After Cyanoacrylate Closure and Surgical Stripping for Incompetent Saphenous Veins	146	Feb 2021



NCT No.	Trial Name	Planned Enrollment	Completion Date
NCT02627846	A Randomised Clinical Trial Comparing Endovenous Laser Ablation and Mechanochemical Ablation (ClariVein®) in the Management of Superficial Venous Insufficiency (LAMA)	140	Sep 2025

NCT: national clinical trial. NTR: Netherlands Trial Registry.

^a Denotes industry-sponsored or cosponsored trial

Clinical Input from Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

In response to requests, input was received from 4 physician specialty societies while this policy was under review in 2015. There was no agreement on the need to treat varicose tributaries to improve functional outcomes in the absence of saphenous vein disease. Input was also mixed on the use of mechanochemical ablation and cyanoacrylate adhesive.

Practice Guidelines and Position Statements

Society for Vascular Surgery and American Venous Forum

The Society for Vascular Surgery and the American Venous Forum (2011) published joint clinical practice guidelines.⁵⁵ [Table 2](#) provides the recommendations.

Table 2. Guidelines on Management of Varicose Veins and Associated Chronic Venous Diseases

Recommendation	Grade ^a	SOR	QOE
Compression therapy for venous ulcerations and varicose veins			
Compression therapy is recommended as the primary treatment to aid healing of venous ulceration	1B	Strong	Moderate



Recommendation	Grade^a	SOR	QOE
To decrease the recurrence of venous ulcers, ablation of the incompetent superficial veins in addition to compression therapy is recommended	1A	Strong	High
Use of compression therapy for patients with symptomatic varicose veins is recommended	2C	Weak	Low
Compression therapy as the primary treatment if the patient is a candidate for saphenous vein ablation is not recommended	1B	Strong	Moderate
Treatment of the incompetent great saphenous vein			
Endovenous thermal ablation (radiofrequency or laser) is recommended over <ul style="list-style-type: none"> • Chemical ablation with foam or • High ligation and stripping due to reduced convalescence and less pain and morbidity. Cryostripping is a technique that is new in the United States, and it has not been fully evaluated.	1B 1B	Strong Strong	Moderate Moderate
Varicose tributaries			
Phlebectomy or sclerotherapy are recommended to treat varicose tributaries	1B	Strong	Moderate
Transilluminated powered phlebectomy using lower oscillation speeds and extended tumescence is an alternative to traditional phlebectomy	2C	Weak	Low
Perforating vein incompetence			
Selective treatment of perforating vein incompetence in patients with simple varicose veins is not recommended	1B	Strong	Moderate
Treatment of pathologic perforating veins (outward flow of ≥ 500 ms duration, with a diameter of ≥ 3.5 mm) located underneath healed or active ulcers (CEAP class C5-C6) is recommended	2B	Weak	Moderate

QOE: quality of evidence; SOR: strength of recommendation.

^a Grading: strong = 1 or weak = 2, based on a level of evidence that is either high quality = A, moderate quality = B, or low quality = C.

Society of Interventional Radiography

In 2019, the Society of Interventional Radiography along with Cardiovascular Interventional Radiological Society of Europe, American College of Phlebology, and Canadian Interventional Radiology Association, published consensus quality improvement guidelines for the treatment of lower extremity superficial venous insufficiency using endovenous thermal ablation.⁵⁶

In a joint statement, American Venous Forum and Society of Interventional Radiography (2007) recommended reporting standards for endovenous ablation for the treatment of venous insufficiency.⁵⁷ They recommended that reporting in clinical studies should include the symptoms of venous disease, history of the disease and prior treatment, the presence of major comorbidities, and any exclusion criteria. It was noted that potential candidates for endovenous ablation might include patients with reflux in an incompetent great saphenous vein or smaller saphenous vein or a major tributary branch of the great or smaller saphenous veins such as the



anterior thigh circumflex vein, posterior thigh circumflex vein, or anterior accessory greater saphenous vein. The presence of reflux in these veins is important to document using duplex ultrasound imaging, and the ultrasound criteria used to define reflux should be indicated. It was also stated that in current practice, most vascular laboratories consider the presence of venous flow reversal for greater than 0.5 to 1.0 second with proximal compression, Valsalva maneuver, or distal compression and release to represent pathologic reflux.

American Vein and Lymphatic Society

In 2015, the American Vein and Lymphatic Society (AVL, previously named the American College of Phlebology) published guidelines on the treatment of superficial vein disease.⁵⁸ AVL gave a Grade 1 recommendation based on high quality evidence that compression is an effective method for the management of symptoms, but when patients have a correctable source of reflux definitive treatment should be offered unless contraindicated. AVL recommends against a requirement for compression therapy when a definitive treatment is available. AVL gave a strong recommendation based on moderate quality evidence that endovenous thermal ablation is the preferred treatment for saphenous and accessory saphenous vein incompetence, and gave a weak recommendation based on moderate quality evidence that mechanochemical ablation may also be used to treat venous reflux.

In 2017, AVL published guidelines on the treatment of refluxing accessory saphenous veins.³⁵ The College gave a Grade 1 recommendation based on level C evidence that patients with symptomatic incompetence of the accessory saphenous veins be treated with endovenous thermal ablation or sclerotherapy to reduce symptomatology. The guidelines noted that although accessory saphenous veins may drain into the great saphenous vein before it drains into the common femoral vein, they can also empty directly into the common femoral vein.

National Institute for Health and Care Excellence

In 2013, The National Institute for Health and Care Excellence (NICE) updated its guidance on ultrasound-guided foam sclerotherapy for varicose veins.⁵⁹ NICE stated that:

“1.1 Current evidence on the efficacy of ultrasound-guided foam sclerotherapy for varicose veins is adequate. The evidence on safety is adequate, and provided that patients are warned of the small but significant risks of foam embolization (see section 1.2), this procedure may be used with normal arrangements for clinical governance, consent and audit.



1.2 During the consent process, clinicians should inform patients that there are reports of temporary chest tightness, dry cough, headaches and visual disturbance, and rare but significant complications including myocardial infarction, seizures, transient ischaemic attacks and stroke.”

The NICE (2016) revised its guidance on endovenous mechanochemical ablation, concluding that “Current evidence on the safety and efficacy of endovenous mechanochemical ablation for varicose veins appears adequate to support the use of this procedure....”⁵⁵.

The NICE (2013) published guidance on the diagnosis and management of varicose veins in the leg.⁶⁰ The NICE (2015) published a technology assessment on the clinical effectiveness and cost-effectiveness of foam sclerotherapy, endovenous laser ablation and surgery for varicose veins.⁵⁷ Five-year trial results are currently being evaluated.

Medicare National Coverage

There is no national coverage determination.

Regulatory Status

In 2015, the VenaSeal® Closure System (Sapheon, a part of Medtronic) was approved by the U.S. Food and Drug Administration (FDA) through the premarket approval (P140018) process for the permanent closure of clinically significant venous reflux through endovascular embolization with coaptation. The VenaSeal® Closure System seals the vein using a cyanoacrylate adhesive agent. FDA product code: PJQ.

In 2013, Varithena™ (formerly Varisolve®), a sclerosant microfoam made with a proprietary gas mix, was approved by the FDA under a new drug application (205-098) for the treatment of incompetent great saphenous veins, accessory saphenous veins and visible varicosities of the great saphenous vein system above and below the knee.

The following devices were cleared for marketing by the FDA through the 501 (k) process for endovenous treatment of superficial vein reflux:

- In 1999, the VNUS® Closure™ system, a radiofrequency device, was cleared by the FDA through the 510(k) process for “endovascular coagulation of blood vessels in patients with superficial vein reflux.” In 2005, the VNUS RFS™ and RFS*flex*™ devices were cleared by the FDA for “use in vessel and tissue coagulation including, treatment of incompetent (ie, refluxing) perforator and tributary veins.” In 2008, the modified VNUS® ClosureFAST™



Intravascular Catheter was cleared by the FDA through the 510(k) process. FDA product code: GEI.

- In 2002, the Diomed 810 nm surgical laser and EVLT™ (endovenous laser therapy) procedure kit were cleared by the FDA through the 510(k) process, "... for use in the endovascular coagulation of the great saphenous vein of the thigh in patients with superficial vein reflux." FDA product code: GEX.
- In 2005, a modified Erbe Erbokryo® cryosurgical unit (Erbe USA) was approved by the FDA for marketing. A variety of clinical indications are listed, including cryostripping of varicose veins of the lower limbs. FDA product code: GEH.
- In 2003, the Trivex® system (InaVein), a device for transilluminated powered phlebectomy (TIPP) was cleared by the FDA through the 510(k) process for "ambulatory phlebectomy procedures for the resection and ablation of varicose veins." FDA product code: DNQ.
- In 2008, the ClariVein® Infusion Catheter (Vascular Insights) was cleared by the FDA through the 510(k) process (K071468) for mechanochemical ablation. The FDA determined that this device was substantially equivalent to the Trellis® Infusion System (K013635) and the Slip-Cath® Infusion Catheter (K882796). The system includes an infusion catheter, motor drive, stopcock and syringe and is intended for the infusion of physician-specified agents in the peripheral vasculature. FDA product code: KRA.

References

1. O'Meara S, Cullum NA, Nelson EA. Compression for venous leg ulcers. *Cochrane Database Syst Rev*. Jan 21 2009(1):CD000265. PMID 19160178
2. O'Meara S, Cullum N, Nelson EA, et al. Compression for venous leg ulcers. *Cochrane Database Syst Rev*. Nov 14 2012;11:CD000265. PMID 23152202
3. Shingler S, Robertson L, Boghossian S, et al. Compression stockings for the initial treatment of varicose veins in patients without venous ulceration. *Cochrane Database Syst Rev*. Nov 9 2011;11(11):CD008819. PMID 22071857
4. Howard DP, Howard A, Kothari A, et al. The role of superficial venous surgery in the management of venous ulcers: a systematic review. *Eur J Vasc Endovasc Surg*. Oct 2008;36(4):458-465. PMID 18675558
5. O'Donnell TF, Jr. The present status of surgery of the superficial venous system in the management of venous ulcer and the evidence for the role of perforator interruption. *J Vasc Surg*. Oct 2008;48(4):1044-1052. PMID 18992425
6. Jones L, Braithwaite BD, Selwyn D, et al. Neovascularisation is the principal cause of varicose vein recurrence: results of a randomised trial of stripping the long saphenous vein. *Eur J Vasc Endovasc Surg*. Nov 1996;12(4):442-445. PMID 8980434
7. Rutgers PH, Kitslaar PJ. Randomized trial of stripping versus high ligation combined with sclerotherapy in the treatment of the incompetent greater saphenous vein. *Am J Surg*. Oct 1994;168(4):311-315. PMID 7943585



8. Nesbitt C, Bedenis R, Bhattacharya V, et al. Endovenous ablation (radiofrequency and laser) and foam sclerotherapy versus open surgery for great saphenous vein varices. *Cochrane Database Syst Rev.* Jul 30 2014;7(7):CD005624. PMID 25075589
9. Paravastu SC, Horne M, Dodd PD. Endovenous ablation therapy (laser or radiofrequency) or foam sclerotherapy versus conventional surgical repair for short saphenous varicose veins. *Cochrane Database Syst Rev.* Nov 29 2016;11:CD010878. PMID 27898181
10. Brittenden J, Cotton SC, Elders A, et al. A randomized trial comparing treatments for varicose veins. *N Engl J Med.* Sep 25 2014;371(13):1218-1227. PMID 25251616
11. Rass K, Frings N, Glowacki P, et al. Comparable effectiveness of endovenous laser ablation and high ligation with stripping of the great saphenous vein: two-year results of a randomized clinical trial (RELACS study). *Arch Dermatol.* Jan 2012;148(1):49-58. PMID 21931012
12. Rass K, Frings N, Glowacki P, et al. Same Site Recurrence is More Frequent After Endovenous Laser Ablation Compared with High Ligation and Stripping of the Great Saphenous Vein: 5 year Results of a Randomized Clinical Trial (RELACS Study). *Eur J Vasc Endovasc Surg.* Nov 2015; 50(5): 648-56. PMID 26319476
13. Christenson JT, Gueddi S, Gemayel G, et al. Prospective randomized trial comparing endovenous laser ablation and surgery for treatment of primary great saphenous varicose veins with a 2-year follow-up. *J Vasc Surg.* Nov 2010;52(5):1234-1241. PMID 20801608
14. Biemans AA, Kockaert M, Akkersdijk GP, et al. Comparing endovenous laser ablation, foam sclerotherapy, and conventional surgery for great saphenous varicose veins. *J Vasc Surg.* Sep 2013;58(3):727-734 e721. PMID 23769603
15. van der Velden SK, Biemans AA, De Maeseneer MG, et al. Five-year results of a randomized clinical trial of conventional surgery, endovenous laser ablation and ultrasound-guided foam sclerotherapy in patients with great saphenous varicose veins. *Br J Surg.* Sep 2015;102(10):1184-1194. PMID 26132315
16. Wallace T, El-Sheikha J, Nandhra S, et al. Long-term outcomes of endovenous laser ablation and conventional surgery for great saphenous varicose veins. *Br J Surg.* Dec 2018; 105(13): 1759-1767. PMID 30132797
17. Theivacumar NS, Darwood RJ, Gough MJ. Endovenous laser ablation (EVLA) of the anterior accessory great saphenous vein (AAGSV): abolition of sapheno-femoral reflux with preservation of the great saphenous vein. *Eur J Vasc Endovasc Surg.* Apr 2009;37(4):477-481. PMID 19201621
18. Hamann SAS, Giang J, De Maeseneer MGR, et al. Editor's Choice - Five Year results of great saphenous vein treatment: a meta-analysis. *Eur J Vasc Endovasc Surg.* Dec 2017;54(6):760-770. PMID 29033337
19. Vahaaho S, Mahmoud O, Halmesmaki K, et al. Randomized clinical trial of mechanochemical and endovenous thermal ablation of great saphenous varicose veins. *Br J Surg.* Apr 2019; 106(5): 548-554. PMID 30908611
20. Shadid N, Ceulen R, Nelemans P, et al. Randomized clinical trial of ultrasound-guided foam sclerotherapy versus surgery for the incompetent great saphenous vein. *Br J Surg.* Aug 2012;99(8):1062-1070. PMID 22627969
21. Lam YL, Lawson JA, Toonder IM, et al. Eight-year follow-up of a randomized clinical trial comparing ultrasound-guided foam sclerotherapy with surgical stripping of the great saphenous vein. *Br J Surg.* May 2018; 105(6): 692-698. PMID 29652081
22. U.S. Food and Drug Administration, Center for Drug Evaluation and Research. Summary Review: 205098 Varithena. 2013; https://www.accessdata.fda.gov/drugsatfda_docs/nda/2013/205098Orig1s000SumR.pdf Accessed October 2020.
23. Todd KL, 3rd, Wright D, for the Vanish-Investigator Group. The VANISH-2 study: a randomized, blinded, multicenter study to evaluate the efficacy and safety of polidocanol endovenous microfoam 0.5% and 1.0% compared with placebo for the treatment of saphenofemoral junction incompetence. *Phlebology.* Oct 2014;29(9):608-618. PMID 23864535
24. Vasquez M, Gasparis AP, Varithena 017 Investigator G. A multicenter, randomized, placebo-controlled trial of endovenous thermal ablation with or without polidocanol endovenous microfoam treatment in patients with great saphenous vein incompetence and visible varicosities. *Phlebology.* May 2017;32(4):272-281. PMID 26957489



25. Bootun R, Lane T, Dharmarajah B, et al. Intra-procedural pain score in a randomised controlled trial comparing mechanochemical ablation to radiofrequency ablation: The Multicentre Venefit versus ClariVein(R) for varicose veins trial. *Phlebology*. Feb 2016;31(1):61-65. PMID 25193822
26. Lane T, Bootun R, Dharmarajah B, et al. A multi-centre randomised controlled trial comparing radiofrequency and mechanical occlusion chemically assisted ablation of varicose veins - Final results of the Venefit versus Clarivein for varicose veins trial. *Phlebology*. Mar 2017;32(2):89-98. PMID 27221810
27. Lam YL, Toonder IM, Wittens CH. Clarivein(R) mechano-chemical ablation an interim analysis of a randomized controlled trial dose-finding study. *Phlebology*. Apr 2016;31(3):170-176. PMID 26249150
28. Holeywijn S, van Eekeren RRJP, Vahl A, et al. Two-year results of a multicenter randomized controlled trial comparing Mechanochemical endovenous Ablation to RADiOfrequeNcy Ablation in the treatment of primary great saphenous vein incompetence (MARADONA trial). *J Vasc Surg Venous Lymphat Disord*. May 2019; 7(3): 364-374. PMID 31000063
29. Mohamed AH, Leung C, Wallace T, et al. A Randomized Controlled Trial of Endovenous Laser Ablation Versus Mechanochemical Ablation With ClariVein in the Management of Superficial Venous Incompetence (LAMA Trial). *Ann Surg*. Jan 21 2020. PMID 31977509
30. Thierens N, Holeywijn S, Vissers WH, et al. Five-year outcomes of mechano-chemical ablation of primary great saphenous vein incompetence. *Phlebology*. May 2020; 35(4): 255-261. PMID 31291849
31. U.S. Food and Drug Administration. VenaSeal Closure System. PMA P140018. 2015; <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfma/pma.cfm?id=P140018> Accessed October 2020.
32. Morrison N, Gibson K, McEnroe S, et al. Randomized trial comparing cyanoacrylate embolization and radiofrequency ablation for incompetent great saphenous veins (VeClose). *J Vasc Surg*. Apr 2015;61(4):985- 994. PMID 25650040
33. Gibson K, Ferris B. Cyanoacrylate closure of incompetent great, small and accessory saphenous veins without the use of post-procedure compression: Initial outcomes of a post-market evaluation of the VenaSeal System (the WAVES Study). *Vascular*. Apr 2017;25(2):149-156. PMID 27206470
34. Klem TM, Schnater JM, Schutte PR, et al. A randomized trial of cryo stripping versus conventional stripping of the great saphenous vein. *J Vasc Surg*. Feb 2009;49(2):403-409. PMID 19028042
35. Gibson K, Khilnani N, Schul M, et al. American College of Phlebology Guidelines - Treatment of refluxing accessory saphenous veins. *Phlebology*. Aug 2017; 32(7): 448-452. PMID 27738242
36. Morrison, NN, Kolluri, RR, Vasquez, MM, Madsen, MM, Jones, AA, Gibson, KK. Comparison of cyanoacrylate closure and radiofrequency ablation for the treatment of incompetent great saphenous veins: 36-Month outcomes of the VeClose randomized controlled trial. *Phlebology*, 2018 Nov 8;268355518810259:268355518810259. PMID 30403154
37. Eroglu E, Yasim A. A Randomised Clinical Trial Comparing N-Butyl Cyanoacrylate, Radiofrequency Ablation and Endovenous Laser Ablation for the Treatment of Superficial Venous Incompetence: Two Year Follow up Results. *Eur J Vasc Endovasc Surg*. Oct 2018; 56(4): 553-560. PMID 30042039
38. Morrison N, Gibson K, Vasquez M, et al. VeClose trial 12-month outcomes of cyanoacrylate closure versus radiofrequency ablation for incompetent great saphenous veins. *J Vasc Surg Venous Lymphat Disord*. May 2017; 5(3): 321-330. PMID 28411697
39. Eroglu E, Yasim A, Ari M, et al. Mid-term results in the treatment of varicose veins with N-butyl cyanoacrylate. *Phlebology*. Dec 2017; 32(10): 665-669. PMID 28669248
40. Zierau UT. Sealing veins with the VenaSeal Sapheon Closure System: results for 795 treated truncal veins after 1000 days. *Vasomed*. 2015;27:124-127.
41. Disselhoff BC, der Kinderen DJ, Kelder JC, et al. Randomized clinical trial comparing endovenous laser with cryostripping for great saphenous varicose veins. *Br J Surg*. Oct 2008;95(10):1232-1238. PMID 18763255
42. Disselhoff BC, der Kinderen DJ, Kelder JC, et al. Five-year results of a randomized clinical trial comparing endovenous laser ablation with cryostripping for great saphenous varicose veins. *Br J Surg*. Aug 2011;98(8):1107-1111. PMID 21633948



43. Tisi PV, Beverley C, Rees A. Injection sclerotherapy for varicose veins. *Cochrane Database Syst Rev*. Oct 18 2006(4):CD001732. PMID 17054141
44. Leopardi D, Hoggan BL, Fitridge RA, et al. Systematic review of treatments for varicose veins. *Ann Vasc Surg*. Mar 2009;23(2):264-276. PMID 19059756
45. El-Sheikha J, Nandhra S, Carradice D, et al. Clinical outcomes and quality of life 5 years after a randomized trial of concomitant or sequential phlebectomy following endovenous laser ablation for varicose veins. *Br J Surg*. Aug 2014;101(9):1093-1097. PMID 24916467
46. Yamaki T, Hamahata A, Soejima K, et al. Prospective randomised comparative study of visual foam sclerotherapy alone or in combination with ultrasound-guided foam sclerotherapy for treatment of superficial venous insufficiency: preliminary report. *Eur J Vasc Endovasc Surg*. Mar 2012;43(3):343-347. PMID 22230599
47. Michaels JA, Campbell WB, Brazier JE, et al. Randomised clinical trial, observational study and assessment of cost-effectiveness of the treatment of varicose veins (REACTIV trial). *Health Technol Assess*. Apr 2006;10(13):1- 196, iii-iv. PMID 16707070
48. Luebke T, Brunkwall J. Meta-analysis of transilluminated powered phlebectomy for superficial varicosities. *J Cardiovasc Surg (Torino)*. Dec 2008; 49(6): 757-64. PMID 19043390
49. Chetter IC, Mylankal KJ, Hughes H, et al. Randomized clinical trial comparing multiple stab incision phlebectomy and transilluminated powered phlebectomy for varicose veins. *Br J Surg*. Feb 2006; 93(2): 169-74. PMID 16432820
50. Tenbrook JA, Jr., lafrati MD, O'Donnell T F, Jr., et al. Systematic review of outcomes after surgical management of venous disease incorporating subfascial endoscopic perforator surgery. *J Vasc Surg*. Mar 2004;39(3):583-589. PMID 14981453
51. van Gent WB, Catarinella FS, Lam YL, et al. Conservative versus surgical treatment of venous leg ulcers: 10- year follow up of a randomized, multicenter trial. *Phlebology*. Mar 2015;30(1 Suppl):35-41. PMID 25729066
52. Blomgren L, Johansson G, Dahlberg-Akerman A, et al. Changes in superficial and perforating vein reflux after varicose vein surgery. *J Vasc Surg*. Aug 2005; 42(2): 315-20. PMID 16102633
53. Lin ZC, Loveland PM, Johnston RV, et al. Subfascial endoscopic perforator surgery (SEPS) for treating venous leg ulcers. *Cochrane Database Syst Rev*. Mar 03 2019; 3: CD012164. PMID 30827037
54. Luebke T, Brunkwall J. Meta-analysis of subfascial endoscopic perforator vein surgery (SEPS) for chronic venous insufficiency. *Phlebology*. Feb 2009;24(1):8-16. PMID 19155335
55. National Institute for Health and Care Excellence (NICE). Endovenous mechanochemical ablation for varicose veins [IPG557]. 2016; <https://www.nice.org.uk/guidance/ipg557> Accessed October 2020.
56. 56. Khilnani NM, Grassi CJ, Kundu S, et al. Multi-society consensus quality improvement guidelines for the treatment of lower-extremity superficial venous insufficiency with endovenous thermal ablation from the Society of Interventional Radiology, Cardiovascular Interventional Radiological Society of Europe, American College of Phlebology and Canadian Interventional Radiology Association. *J Vasc Interv Radiol*. Jan 2010; 21(1): 14-31. PMID 20123189
57. Brittenden J, Cotton SC, Elders A, et al. Clinical effectiveness and cost-effectiveness of foam sclerotherapy, endovenous laser ablation and surgery for varicose veins: results from the Comparison of LAser, Surgery and foam Sclerotherapy (CLASS) randomised controlled trial. *Health Technol Assess*. Apr 2015;19(27):1-342. PMID 25858333
58. American College of Phlebology. Superficial venous disease. 2015; <https://www.myavls.org/assets/pdf/VaricoseVeinGuidelines3.9.15.pdf> . Accessed October 2020
59. National Institute for Health and Care Excellence (NICE). Ultrasound-guided foam sclerotherapy for varicose veins [IPG440] 2013; <https://www.nice.org.uk/guidance/ipg440> Accessed October 2020.
60. National Institute for Health and Care Excellence (NICE). Varicose veins: diagnosis and management [CG168]. 2013; <https://www.nice.org.uk/guidance/cg168> Accessed October 2020.



History

Date	Comments
12/11/12	Replace policy. Formatting changed to now organize procedures by type of vein. Policy statement extensively revised with additional medically necessary criteria for each procedure. Sclerotherapy now considered medically necessary only for accessory and tributary veins. Policy on hold for provider notification; the effective date is May 19, 2013.
04/08/13	Policy statement clarification. In criteria for sclerotherapy of accessory saphenous veins when not done concurrently with other listed treatments, the moderate to severe saphenous reflux criterion has been removed. To clarify the definition of "symptomatic" for treatment of tributaries not done concurrently with other listed treatments, specific criteria added for stab avulsion, hook phlebectomy, sclerotherapy or transilluminated powered phlebectomy treatment of tributaries. Tributary criterion for moderate to severe saphenous reflux, when not done concurrently with other listed treatments, has been removed.
06/14/13	Minor clarification. Hook phlebectomy clarified to indicate that it is also known as stab phlebectomy or micro-phlebectomy.
10/14/13	Replace policy. CPT codes for sclerotherapy (36468, 36469, 36470, 36471), TIPP (37765 or 37766, stab avulsion (37765, or 37799), etc. added to Policy Guidelines section.
02/27/14	Update Related Policies. Add 10.01.514.
06/09/14	Interim update. Criteria for vein size removed from policy statement. Minor edit to change definition of "moderate to severe reflux" to greater than 0.5 second. Coding update: ICD-9 and ICD-10 diagnosis and procedure codes removed – these are not utilized for adjudication.
02/10/15	Annual Review. Policy updated with literature review through September 23, 2014; reference 8-9, 18, 24 33 added and some references removed; policy statement revised to allow sclerotherapy as medically necessary when criteria are met. Information added regarding FDA approval of Varithena sclerotherapy. Documentation requirements added to Policy Guidelines.
05/12/15	Interim Update. Policy statement clarified: Treatment of tributaries by sclerotherapy, stab avulsion, stab/hook phlebectomy or transilluminated powered phlebectomy may now be considered medically necessary up to 10 days after saphenous veins have been treated. Definition of moderate to severe saphenous reflux and abbreviation "TIPP" added to the policy statement.
06/09/15	Interim Update. Policy statement clarified: TRIVEX listed as an example of TIPP. Exercises and pain relievers removed from conservative management requirement. TRIVEX manufacturer website added to Reference section. CPT codes 76942, 93970-71 and HCPCS code S2202 removed; these are not reviewed.
11/10/15	Interim Update. Policy updated with clarification to coverage criteria for the accessory saphenous vein, indicating parameters around treatment of the greater or lesser saphenous veins and the presence of reflux.



Date	Comments
01/12/16	Annual Review. Added statement that cyanoacrylate adhesive/embolization (CAE) (eg, VenaSeal closure system) and endovenous mechanochemical ablation (MOCA) (eg ClariVein) of any vein is considered investigational. The failure of compression therapy removed from the policy statements on venous stasis and thrombophlebitis. References added. Policy updated with literature review through July 7, 2015.
10/01/16	Interim Update, approved September 13, 2016. Perforator vein criteria revised: Requirement of active ulcers changed to "healed or active ulcers" and "demonstrated perforator reflux" is more clearly defined with measurements of outward flow and vein diameter.
12/01/16	Policy moved to new format. Policy statements unchanged. Added Prior Authorization Requirements section.
01/01/17	Coding update, added CPT codes 36473 and 36474 effective 1/1/17.
03/01/17	Annual Review, approved February 14, 2017. No change to policy statement. Compression Therapy added to Definition of Terms. Reference 64 added. Removed CPT code 36469 from policy as it terminated as of 01/01/2015.
11/02/17	Minor formatting edits made.
01/16/18	Minor edit, added Documentation Requirements table to the Policy Coverage Criteria section.
01/23/18	Coding update, added CPT codes 36465, 36466, 36482, and 36483 (new codes effective 1/1/18).
08/01/18	Annual Review, approved July 25, 2018. Policy updated with literature review through March 2018; references 18, 21, 24-27, and 30-31 added; references 54 and 56 updated. Policy statements unchanged.
11/01/18	Interim Review, approved October 9, 2018. Reordered and clarified policy statement criteria by procedure type rather than vein type. Added criteria for number of treatment sessions of sclerotherapy allowed and criteria for requests for sclerotherapy after ablation of the saphenous veins for the treatment of residual or recurrent symptoms.
01/01/19	Interim Review, approved December 19, 2018. Added missing wording to statement: "Phlebectomy (stab avulsion, hook phlebectomy, or TIPP) or initial* sclerotherapy as a component of the treatment of <u>symptomatic varicose tributaries</u> when performed either at the same time or following prior treatment (surgical, radiofrequency, or laser) of the saphenous veins is considered medically necessary" to clarify the intent of the statement. Added new HCPCS code 0524T (new code effective 1/1/19).
04/01/19	Annual Review, approved March 19, 2019. Policy updated with literature review through November 2018; references 16, 19, 33-34 added. Minor edits to policy statements for clarity; otherwise intent of policy statements unchanged.
08/01/19	Interim Review, approved July 11, 2019. Policy updated with literature review through March 2019, references 60, and 65-67 added. Cyanoacrylate adhesive may be considered medically necessary. A statement was added on concurrent treatment of the accessory saphenous veins.
12/01/19	Interim Review, approved November 26, 2019. Policy effective date changed to January 1, 2020.



Date	Comments
04/01/20	Delete policy, approved March 10, 2020. This policy will be deleted effective July 2, 2020, replaced with policy 10.01.530.
07/02/20	Delete policy.
11/01/20	Policy reinstated effective February 5, 2021 approved October 13, 2020. Policy updated with literature review through March 2020; references added. Policy statements unchanged

Disclaimer: This medical policy is a guide in evaluating the medical necessity of a particular service or treatment. The Company adopts policies after careful review of published peer-reviewed scientific literature, national guidelines and local standards of practice. Since medical technology is constantly changing, the Company reserves the right to review and update policies as appropriate. Member contracts differ in their benefits. Always consult the member benefit booklet or contact a member service representative to determine coverage for a specific medical service or supply. CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). ©2020 Premera All Rights Reserved.

Scope: Medical policies are systematically developed guidelines that serve as a resource for Company staff when determining coverage for specific medical procedures, drugs or devices. Coverage for medical services is subject to the limits and conditions of the member benefit plan. Members and their providers should consult the member benefit booklet or contact a customer service representative to determine whether there are any benefit limitations applicable to this service or supply. This medical policy does not apply to Medicare Advantage.



Discrimination is Against the Law

Premera Blue Cross complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex. Premera does not exclude people or treat them differently because of race, color, national origin, age, disability or sex.

Premera:

- Provides free aids and services to people with disabilities to communicate effectively with us, such as:
 - Qualified sign language interpreters
 - Written information in other formats (large print, audio, accessible electronic formats, other formats)
- Provides free language services to people whose primary language is not English, such as:
 - Qualified interpreters
 - Information written in other languages

If you need these services, contact the Civil Rights Coordinator.

If you believe that Premera has failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, or sex, you can file a grievance with:

Civil Rights Coordinator - Complaints and Appeals
PO Box 91102, Seattle, WA 98111
Toll free 855-332-4535, Fax 425-918-5592, TTY 800-842-5357
Email AppealsDepartmentInquiries@Premera.com

You can file a grievance in person or by mail, fax, or email. If you need help filing a grievance, the Civil Rights Coordinator is available to help you.

You can also file a civil rights complaint with the U.S. Department of Health and Human Services, Office for Civil Rights, electronically through the Office for Civil Rights Complaint Portal, available at <https://ocrportal.hhs.gov/ocr/portal/lobby.jsf>, or by mail or phone at: U.S. Department of Health and Human Services
200 Independence Avenue SW, Room 509F, HHH Building
Washington, D.C. 20201, 1-800-368-1019, 800-537-7697 (TDD)
Complaint forms are available at <http://www.hhs.gov/ocr/office/file/index.html>.

Getting Help in Other Languages

This Notice has Important Information. This notice may have important information about your application or coverage through Premera Blue Cross. There may be key dates in this notice. You may need to take action by certain deadlines to keep your health coverage or help with costs. You have the right to get this information and help in your language at no cost. Call 800-722-1471 (TTY: 800-842-5357).

አማርኛ (Amharic):

ይህ ማስታወቂያ አስፈላጊ መረጃ ይዟል። ይህ ማስታወቂያ ስለ ማመልከቻዎ ወይም የ Premera Blue Cross ሽፋን አስፈላጊ መረጃ ሊኖረው ይችላል። በዚህ ማስታወቂያ ውስጥ ቁልፍ ቀናት ሊኖሩ ይችላሉ። የጤና ሽፋንዎን ለማመልከት በአስፈላጊ እርዳታ ለማግኘት በተውሰኑ የጊዜ ገደቦች እርምጃ መውሰድ ይገባዎት ይሆናል። ይህን መረጃ እንዲያገኙ እና የለምንም ክፍያ በቋንቋዎ እርዳታ እንዲያገኙ መሰታወቅ አለዎት። በስልክ ቁጥር 800-722-1471 (TTY: 800-842-5357) ይደውሉ።

العربية (Arabic):

يحتوي هذا الإشعار على معلومات هامة. قد يحتوي هذا الإشعار على معلومات مهمة بخصوص طلبك أو التخطيط التي تزيد الحصول عليها من خلال Premera Blue Cross. قد تكون هناك تواريخ مهمة في هذا الإشعار. وقد تحتاج لاتخاذ إجراء في تاريخ معينه للحفاظ على تغطيتك الصحية أو للمساعدة في دفع التكاليف. يحق لك الحصول على هذه المعلومات والمساعدة بلغتك دون تكبد أية تكلفة. اتصل بـ 800-722-1471 (TTY: 800-842-5357)

中文 (Chinese):

本通知有重要的訊息。本通知可能有關於您透過 Premera Blue Cross 提交的申請或保險的重要訊息。本通知內可能有重要日期。您可能需要在截止日期之前採取行動，以保留您的健康保險或者費用補貼。您有權利免費以您的母語得到本訊息和幫助。請撥電話 800-722-1471 (TTY: 800-842-5357)。

Oromoo (Cushite):

Beeksisni kun odeeffannoo barbaachisaa qaba. Beeksisti kun sagantaa yookan karaa Premera Blue Cross tiin tajaajila keessan ilaalchisee odeeffannoo barbaachisaa qabaachuu danda'a. Guyyaawwan murteessaa ta'an beeksisa kana keessatti ilaalaa. Tarii kaffaltiidhaan deeggaramuuf yookan tajaajila fayyaa keessaniif guyyaa dhumaa irratti wanti raawwattan jiraachuu danda'a. Kaffaltii irraa bilisa haala ta'een afaan keessaniin odeeffannoo argachuu fi deeggarsa argachuuf mirga ni qabaattu. Lakkoofsa bilbilaa 800-722-1471 (TTY: 800-842-5357) tii bilbilaa.

Français (French):

Cet avis a d'importantes informations. Cet avis peut avoir d'importantes informations sur votre demande ou la couverture par l'intermédiaire de Premera Blue Cross. Le présent avis peut contenir des dates clés. Vous devez peut-être prendre des mesures par certains délais pour maintenir votre couverture de santé ou d'aide avec les coûts. Vous avez le droit d'obtenir cette information et de l'aide dans votre langue à aucun coût. Appelez le 800-722-1471 (TTY: 800-842-5357).

Kreyòl ayisyen (Creole):

Avi sila a gen Enfòmasyon Enpòtan ladann. Avi sila a kapab genyen enfòmasyon enpòtan konsènan aplikasyon w lan oswa konsènan kouvèti asirans lan atravè Premera Blue Cross. Kapab genyen dat ki enpòtan nan avi sila a. Ou ka gen pou pran kèk aksyon avan sèten dat limit pou ka kenbe kouvèti asirans sante w la oswa pou yo ka ede w avèk depans yo. Se dwa w pou resewva enfòmasyon sa a ak asistans nan lang ou pale a, san ou pa gen pou peye pou sa. Rele nan 800-722-1471 (TTY: 800-842-5357).

Deutsche (German):

Diese Benachrichtigung enthält wichtige Informationen. Diese Benachrichtigung enthält unter Umständen wichtige Informationen bezüglich Ihres Antrags auf Krankenversicherungsschutz durch Premera Blue Cross. Suchen Sie nach eventuellen wichtigen Terminen in dieser Benachrichtigung. Sie könnten bis zu bestimmten Stichtagen handeln müssen, um Ihren Krankenversicherungsschutz oder Hilfe mit den Kosten zu behalten. Sie haben das Recht, kostenlose Hilfe und Informationen in Ihrer Sprache zu erhalten. Rufen Sie an unter 800-722-1471 (TTY: 800-842-5357).

Hmoob (Hmong):

Tsab ntawv tshaj xo no muaj cov ntshiab lus tseem ceeb. Tej zaum tsab ntawv tshaj xo no muaj cov ntshiab lus tseem ceeb txog koj daim ntawv thov kev pab los yog koj qhov kev pab cuam hnu ntawm Premera Blue Cross. Tej zaum muaj cov hnuv tseem ceeb uas sau rau hauv daim ntawv no. Tej zaum koj kuj yuav tau ua qee yam uas peb kom koj ua tsis pub dhau cov caij nyoog uas teev tseg rau hauv daim ntawv no mas koj thiaj yuav tau txais kev pab cuam kho mob los yog kev pab them tej nqi kho mob ntawd. Koj muaj cai kom lawv muab cov ntshiab lus no uas tau muab sau ua koj hom lus pub dawb rau koj. Hu rau 800-722-1471 (TTY: 800-842-5357).

Iloko (Ilocano):

Daytoy a Pakdaar ket naglaon iti Napateg nga Impormasion. Daytoy a pakdaar mabalin nga adda ket naglaon iti napateg nga impormasion maipanggep iti aplikasyonyo wenna coverage babaen iti Premera Blue Cross. Daytoy ket mabalin dagiti importante a petsa iti daytoy a pakdaar. Mabalin nga adda rumbeng nga aramidenyo nga addang sakbay dagiti partikular a naituding nga aldaw tapno mapagtalinaedyo ti coverage ti salun-atyto wenna tulong kadagiti gastos. Adda karbenganyo a mangala iti daytoy nga impormasion ken tulong iti bukodyo a pagsasao nga awan ti bayadanyo. Tumawag iti numero nga 800-722-1471 (TTY: 800-842-5357).

Italiano (Italian):

Questo avviso contiene informazioni importanti. Questo avviso può contenere informazioni importanti sulla tua domanda o copertura attraverso Premera Blue Cross. Potrebbero esserci date chiave in questo avviso. Potrebbe essere necessario un tuo intervento entro una scadenza determinata per consentirti di mantenere la tua copertura o sovvenzione. Hai il diritto di ottenere queste informazioni e assistenza nella tua lingua gratuitamente. Chiama 800-722-1471 (TTY: 800-842-5357).

日本語 (Japanese):

この通知には重要な情報が含まれています。この通知には、Premera Blue Cross の申請または補償範囲に関する重要な情報が含まれている場合があります。この通知に記載されている可能性がある重要な日付をご確認ください。健康保険や有料サポートを維持するには、特定の期日までに行動を取らなければならない場合があります。ご希望の言語による情報とサポートが無料で提供されます。800-722-1471 (TTY: 800-842-5357)までお電話ください。

한국어 (Korean):

본 통지서에는 중요한 정보가 들어 있습니다. 즉 이 통지서는 귀하의 신청에 관하여 그리고 Premera Blue Cross 를 통한 커버리지에 관한 정보를 포함하고 있을 수 있습니다. 본 통지서에는 핵심이 되는 날짜들이 있을 수 있습니다. 귀하의 건강 커버리지를 계속 유지하거나 비용을 절감하기 위해서 일정한 마감일까지 조치를 취해야 할 필요가 있을 수 있습니다. 귀하의 이러한 정보와 도움을 귀하의 언어로 비용 부담없이 얻을 수 있는 권리가 있습니다. 800-722-1471 (TTY: 800-842-5357) 로 전화하십시오.

ລາວ (Lao):

ແຈງການນີ້ມີຂໍ້ມູນສໍາຄັນ. ແຈງການນີ້ອາດຈະມີຂໍ້ມູນສໍາຄັນກ່ຽວກັບຄໍາອ້ອງສະໝັກ ຫຼື ຄວາມຄົມຄອງປະກັນໄພຂອງທ່ານຜ່ານ Premera Blue Cross. ອາດຈະມີວັນທີ່ສໍາຄັນໃນແຈງການນີ້. ທ່ານອາດຈະຈໍາເປັນຕ້ອງດໍາເນີນການຕາມກຳນົດ ເວລາສະເພາະເພື່ອຮັກສາຄວາມຄົມຄອງປະກັນສະພາບ ຫຼື ຄວາມຊ່ວຍເຫຼືອເວັ້ນເວົ້ອງຄ່າໃຊ້ຈ່າຍຂອງທ່ານໄດ້. ທ່ານມີສິດໄດ້ຮັບຂໍ້ມູນນີ້ ແລະ ຄວາມຊ່ວຍເຫຼືອເປັນພາສາຂອງທ່ານໂດຍບໍ່ເສຍຄ່າ. ໃຫ້ໃບທາ 800-722-1471 (TTY: 800-842-5357).

ភាសាខ្មែរ (Khmer):

សេចក្តីជូនដំណឹងនេះមានព័ត៌មានយ៉ាងសំខាន់។ សេចក្តីជូនដំណឹងនេះប្រហែលជាមានព័ត៌មានយ៉ាងសំខាន់អំពីទម្រង់បែបបទ ឬការរៀបចំរបស់អ្នកតាមរយៈ Premera Blue Cross ។ ប្រហែលជាមាន កាលបរិច្ឆេទសំខាន់នៅក្នុងសេចក្តីជូនដំណឹងនេះ។ អ្នកប្រហែលជាត្រូវការបញ្ជាក់សមត្ថភាព ដល់កំណត់ថ្លៃជាតំបន់នានា ដើម្បីនឹងរក្សាទុកការធានារ៉ាប់រងអនាគតរបស់អ្នក ឬប្រាក់ដុល្លារចេញថ្លៃ។ អ្នកមានសិទ្ធិទទួលបានព័ត៌មាននេះ និងដុល្លារនៅក្នុងភាសារបស់អ្នកដោយមិនអស់លុយឡើយ។ សូមទូរស័ព្ទ 800-722-1471 (TTY: 800-842-5357)។

ਪੰਜਾਬੀ (Punjabi):

ਇਸ ਨੋਟਿਸ ਵਿਚ ਖਾਸ ਜਾਣਕਾਰੀ ਹੈ. ਇਸ ਨੋਟਿਸ ਵਿਚ Premera Blue Cross ਵਲੋਂ ਤੁਹਾਡੀ ਕਵਰੇਜ ਅਤੇ ਅਰਜੀ ਬਾਰੇ ਮਹੱਤਵਪੂਰਨ ਜਾਣਕਾਰੀ ਹੋ ਸਕਦੀ ਹੈ . ਇਸ ਨੋਟਿਸ ਨਵਚ ਖਾਸ ਤਾਰੀਖਾਂ ਹੋ ਸਕਦੀਆਂ ਹਨ. ਜੇਕਰ ਤੁਸੀਂ ਜਸਰਤ ਕਵਰੇਜ ਰਿੱਖਣੀ ਹੋਵੇ ਜਾਂ ਓਸ ਦੀ ਲਾਗਤ ਜਵਿੱਚ ਮਦਦ ਦੇ ਇਛੁੱਕ ਹੋ ਤਾਂ ਤੁਹਾਨੂੰ ਅੰਤਮ ਤਾਰੀਖ ਤੋਂ ਪਹਿਲਾਂ ਢੁੱਝ ਖਾਸ ਕਦਮ ਚੁੱਕਣ ਦੀ ਲੋੜ ਹੋ ਸਕਦੀ ਹੈ ,ਤੁਹਾਨੂੰ ਮੁਫਤ ਵਿੱਚ ਤੋਂ ਅਪਣੀ ਭਾਸ਼ਾ ਵਿੱਚ ਜਾਣਕਾਰੀ ਅਤੇ ਮਦਦ ਪ੍ਰਾਪਤ ਕਰਨ ਦਾ ਅਧਿਕਾਰ ਹੈ ,ਕਾਲ 800-722-1471 (TTY: 800-842-5357).

فارسی (Farsi):

این اعلامیه حاوی اطلاعات مهم میباشد. این اعلامیه ممکن است حاوی اطلاعات مهم درباره فرم تقاضا و یا پوشش بیمه ای شما از طریق Premera Blue Cross باشد. به تاریخ های مهم در این اعلامیه توجه نمایید. شما ممکن است برای حفظ پوشش بیمه تان یا کمک در پرداخت هزینه های درمانی تان، به تاریخ های مشخصی برای انجام کارهای خاصی احتیاج داشته باشید. شما حق این را دارید که این اطلاعات و کمک را به زبان خود به طور رایگان دریافت نمایید. برای کسب اطلاعات با شماره 800-722-1471 (کلیربران TTY تماس باشماره 800-842-5357) تماس برقرار نمایید.

Polskie (Polish):

To ogłoszenie może zawierać ważne informacje. To ogłoszenie może zawierać ważne informacje odnośnie Państwa wniosku lub zakresu świadczeń poprzez Premera Blue Cross. Prosimy zwrócić uwagę na kluczowe daty, które mogą być zawarte w tym ogłoszeniu aby nie przekroczyć terminów w przypadku utrzymania polisy ubezpieczeniowej lub pomocy związanej z kosztami. Macie Państwo prawo do bezpłatnej informacji we własnym języku. Zadzwońcie pod 800-722-1471 (TTY: 800-842-5357).

Português (Portuguese):

Este aviso contém informações importantes. Este aviso poderá conter informações importantes a respeito de sua aplicação ou cobertura por meio do Premera Blue Cross. Poderão existir datas importantes neste aviso. Talvez seja necessário que você tome providências dentro de determinados prazos para manter sua cobertura de saúde ou ajuda de custos. Você tem o direito de obter esta informação e ajuda em seu idioma e sem custos. Ligue para 800-722-1471 (TTY: 800-842-5357).

Română (Romanian):

Prezenta notificare conține informații importante privind cererea sau acoperirea asigurării dumneavoastră de sănătate prin Premera Blue Cross. Pot exista date cheie în această notificare. Este posibil să fie nevoie să acționați până la anumite termene limită pentru a vă menține acoperirea asigurării de sănătate sau asistența provizorie la costuri. Aveți dreptul de a obține gratuit aceste informații și ajutor în limba dumneavoastră. Sunați la 800-722-1471 (TTY: 800-842-5357).

Русский (Russian):

Настоящее уведомление содержит важную информацию. Это уведомление может содержать важную информацию о вашем заявлении или страховом покрытии через Premera Blue Cross. В настоящем уведомлении могут быть указаны ключевые даты. Вам, возможно, потребуется принять меры к определенным предельным срокам для сохранения страхового покрытия или помощи с расходами. Вы имеете право на бесплатное получение этой информации и помощь на вашем языке. Звоните по телефону 800-722-1471 (TTY: 800-842-5357).

Fa'asamoa (Samoan):

Atonu ua iai i lenei fa'asilasilaga ni fa'amatalaga e sili ona taua e tatau ona e malamalama i ai. O lenei fa'asilasilaga o se fesoasoani e fa'amatala atili i ai i le tulaga o le polokalame, Premera Blue Cross, ua e tau fia maua atu i ai. Fa'amolemole, ia e iloilo fa'alelei i aso fa'apitoa olo'o iai i lenei fa'asilasilaga taua. Masalo o le'a iai ni feau e tatau ona e faia ao le'i aulia le aso ua ta'ua i lenei fa'asilasilaga ina ia e iai pea ma maua fesoasoani mai ai i le polokalame a le Malo olo'o e iai i ai. Olo'o iai iate oe le aia tatau e maua atu i lenei fa'asilasilaga ma lenei fa'matalaga i legagana e te malamalama i ai aunoa ma se togiga tupe. Vili atu i le telefoni 800-722-1471 (TTY: 800-842-5357).

Español (Spanish):

Este Aviso contiene información importante. Es posible que este aviso contenga información importante acerca de su solicitud o cobertura a través de Premera Blue Cross. Es posible que haya fechas clave en este aviso. Es posible que deba tomar alguna medida antes de determinadas fechas para mantener su cobertura médica o ayuda con los costos. Usted tiene derecho a recibir esta información y ayuda en su idioma sin costo alguno. Llame al 800-722-1471 (TTY: 800-842-5357).

Tagalog (Tagalog):

Ang Paunawa na ito ay naglalaman ng mahalagang impormasyon tungkol sa iyong aplikasyon o pagsakop sa pamamagitan ng Premera Blue Cross. Maaaring may mga mahalagang petsa dito sa paunawa. Maaring mangailangan ka na magsagawa ng hakbang sa ilang mga itinakdang panahon upang mapanatili ang iyong pagsakop sa kalusugan o tulong na walang gastos. May karapatan ka na makakuha ng ganiitong impormasyon at tulong sa iyong wika ng walang gastos. Tumawag sa 800-722-1471 (TTY: 800-842-5357).

ไทย (Thai):

ประกาศนี้มีข้อมูลสำคัญ ประกาศนี้อาจมีข้อมูลที่สำคัญเกี่ยวกับกาการสมัครหรือขอบเขตประกันสุขภาพของคุณผ่าน Premera Blue Cross และอาจมีกำหนดการในประกาศนี้ คุณอาจจะต้องดำเนินการภายในกำหนดระยะเวลาที่แน่นอนเพื่อจะรักษาการประกันสุขภาพของคุณหรือการช่วยเหลือที่มีค่าใช้จ่าย คุณมีสิทธิที่จะได้รับข้อมูลและความช่วยเหลือนี้ในภาษาของคุณโดยไม่มีค่าใช้จ่าย โทร 800-722-1471 (TTY: 800-842-5357)

Український (Ukrainian):

Це повідомлення містить важливу інформацію. Це повідомлення може містити важливу інформацію про Ваше звернення щодо страховального покриття через Premera Blue Cross. Зверніть увагу на ключові дати, які можуть бути вказані у цьому повідомленні. Існує імовірність того, що Вам треба буде здійснити певні кроки у конкретні кінцеві строки для того, щоб зберегти Ваше медичне страхування або отримати фінансову допомогу. У Вас є право на отримання цієї інформації та допомоги безкоштовно на Вашій рідній мові. Дзвоніть за номером телефону 800-722-1471 (TTY: 800-842-5357).

Tiếng Việt (Vietnamese):

Thông báo này cung cấp thông tin quan trọng. Thông báo này có thông tin quan trọng về đơn xin tham gia hoặc hợp đồng bảo hiểm của quý vị qua chương trình Premera Blue Cross. Xin xem ngày quan trọng trong thông báo này. Quý vị có thể phải thực hiện theo thông báo đúng trong thời hạn để duy trì bảo hiểm sức khỏe hoặc được trợ giúp thêm về chi phí. Quý vị có quyền được biết thông tin này và được trợ giúp bằng ngôn ngữ của mình miễn phí. Xin gọi số 800-722-1471 (TTY: 800-842-5357).